

**A REMARK ON THE EXISTENCE OF SPHERE WITH
PRESCRIBED MEAN CURVATURE***

SHING TUNG YAU†

On 1976, the author (see the open problem section in [3]) asked the following question: if ρ is a function defined on R^3 , what is the condition on ρ so that there is a sphere in R^3 whose mean curvature is given by ρ . These were important works by Treibergs-Wei [2] and others on this question. Recently in a lecture that I gave in the Chinese University of Hong Kong, I realized that my work with R. Schoen on the existence of black Hole can be used to settle the prescribed mean curvature question for a large class of ρ .

Recall that in [1], we are given on a three dimensional manifold M , two tensors (g_{ij}, p_{ij}) where $g_{ij} > 0$ is a Riemannian metric.

Let

$$\mu = \frac{1}{2} \left[R - \sum p^{ij} p_{ij} + \left(\sum p_i^i \right)^2 \right]$$

$$J^i = \sum_j D_j \left[p^{ij} - \left(\sum p_k^k \right) g^{ij} \right]$$

where R is the scalar curvature of M .

For a domain Ω in M , we define

$$Rad(\Omega) = \sup \left\{ r : \text{for any simple closed curve } \Gamma \text{ in } \Omega \right.$$

which is homotopically trivial, Γ
does not bound a disk in a tubular
neighborhood $N_r(\Gamma)$ of radius r }

Then the major theorem in [1] says

THEOREM 1. *Let Ω be a bounded region on which $\mu - |J| \geq \Lambda > 0$ holds. Assume that $\Omega \subset \Omega_1$ where the mean curvature of $\partial\Omega_1$ (with respect to the outer normal) is strictly greater than the absolute value of the trace of $p_{ij}|_{\partial\Omega_1}$. If $Rad(\Omega) \geq \sqrt{\frac{3}{2}} \frac{\pi}{\sqrt{\Lambda}}$, then Ω_1 contains a sphere Σ such that the mean curvature of $\Sigma = \pm \text{tr}_\Sigma(p_{ij})$. Moreover, any such Σ lying within Ω has diameter at most $\frac{2\pi}{\sqrt{3}\sqrt{\Lambda}}$ and any such Σ intersecting $\partial\Omega$ has the property that $\Omega \cap \Sigma$ has everywhere within distance $\frac{2\pi}{\sqrt{3}\sqrt{\Lambda}}$ of $\partial\Omega$.*

In this note, we apply this theorem to the prescribed mean curvature problem:

Let $p_{ij} = \frac{\rho}{2} g_{ij}$. Then

$$2\mu = R + \frac{3}{2} \rho^2$$

$$J^i = (D_j \rho) g^{ij}$$

*Received July 12, 1997; accepted for publication September 8, 1997.

†Department of Mathematics, Harvard University, Cambridge, MA 02138, U.S.A. (yau@math.harvard.edu) and Department of Mathematics, Chinese University of Hong Kong, Shatin, Hong Kong.

and $\mu - |J| = \frac{1}{2}R + \frac{3}{4}\rho^2 - |D\rho|$.

Hence we have the following theorem

THEOREM 2. *Let Ω_1 be a compact domain in M so that the mean curvature of $\partial\Omega_1$ (with respect to its outer normal) is greater than $|\rho|$. Let $\Omega \subset \Omega_1$ be a subdomain on which*

$$\frac{1}{2}R + \frac{3}{4}\rho^2 - |D\rho| \geq \Lambda > 0$$

where R is the scalar curvature of M .

If $\text{Rad}(\Omega) \geq \sqrt{\frac{3}{2}\frac{\pi}{\Lambda}}$, then Ω_1 contains a sphere Σ intersecting Ω so that the mean curvature of $\Sigma = \pm\rho$. Moreover, any such Σ lying within Ω has diameter at most $\frac{2\pi}{\sqrt{3}\sqrt{\Lambda}}$ and any such Σ intersecting $\partial\Omega$ has the property that $\Omega \cap \Sigma$ lies everywhere within distance $\frac{2\pi}{\sqrt{3}\sqrt{\Lambda}}$ of $\partial\Omega$.

In [4], we also observed the following: Let $\lambda_1 \geq \lambda_2 \geq \lambda_3$ be eigenvalue of the symmetric tensor p_{ij} . Then if for some nonnegative function f and $\Omega \subset \Omega_1$ so that

$$\int_{\partial\Omega} f + \int_{\Omega} |\nabla f| < \int_{\Omega} f \min(\lambda_2 + \lambda_3, \lambda_2 + 2\lambda_3)$$

then there is a closed surface Σ such that the mean curvature of $\Sigma = \pm \text{tr}_{\Sigma}(p_{ij})$.

THEOREM 3. *Let Ω_1 be a compact domain in M so that the mean curvature of $\partial\Omega_1$, (with respect to outer normal) is greater than $|\rho|$. Suppose for some domain $\Omega \subset \Omega_1$, and for some nonnegative function f ,*

$$\int_{\partial\Omega} f + \int_{\Omega} |\nabla f| < \int_{\substack{\Omega \\ \rho > 0}} f\rho + \frac{3}{2} \int_{\substack{\Omega \\ \rho < 0}} f\rho$$

Then there is a closed surface Σ whose mean curvature $= \pm\rho$.

REMARK It is an interesting question to determine the genus of Σ .

REFERENCES

- [1] R. SCHOEN AND S. T. YAU, *The existence of a black hole due to condensation of matter*, Comm. Math. Phys. **90** (1983) 575-579.
- [2] A. TREIBERGS AND S. W. WEI, *Embedded hyperspheres with prescribed mean curvature*, J. Differential Geom. **18** (1983) 513-521.
- [3] S. T. YAU, *Applications of geometric ideas in general relativity: black holes and conserved quantity*, to appear.
- [4] S. T. YAU, *Seminar on Differential Geometry*, Princeton University Press, Princeton, 1982.