

ON QUANTUM COHOMOLOGY RINGS OF FANO MANIFOLDS AND A FORMULA OF VAFA AND INTRILIGATOR*

BERND SIEBERT[†] AND GANG TIAN[‡]

0. Introduction. Quantum multiplications on the cohomology of symplectic manifolds were first proposed by the physicist Vafa [Va2], based on Witten’s topological sigma models [Wi1]. In [RuTi], Ruan and the second named author gave a mathematical construction of quantum multiplications on cohomology groups of *positive* symplectic manifolds (cf. Chapter 1). The definition uses certain symplectic invariants, called Gromov-Witten (GW-) invariants, that were previously defined by Ruan for (semi-) positive symplectic manifolds [Ru]. A large class of such manifolds is provided by Fano manifolds (complex manifolds with ample anti-canonical bundle). Examples are low degree complete intersections and compact complex homogeneous spaces like Grassmann manifolds. If M is a Fano (or positive symplectic) manifold the quantum cohomology $QH_{[\omega]}^*(M)$ is just the cohomology space $H^*(M, \mathbb{C})$ with a (non-homogeneous) associative, graded commutative multiplication, the quantum multiplication. This multiplication depends on the choice of a (complexified) Kähler class $[\omega]$ on M . Its homogeneous part (the “weak coupling limit” $\lambda \cdot [\omega]$, $\lambda \rightarrow \infty$) is the usual cup product.

In this note we observe that quantum cohomology rings have a nice description in terms of generators and relations: If $H^*(M, \mathbb{C}) = \mathbb{C}[X_1, \dots, X_N]/(f_1, \dots, f_k)$ is a presentation of the cohomology ring (for simplicity we assume $\deg X_i$ even for the moment) then $QH_{[\omega]}^*(M) = \mathbb{C}[X_1, \dots, X_N]/(f_1^{[\omega]}, \dots, f_k^{[\omega]})$, where $f_1^{[\omega]}, \dots, f_k^{[\omega]}$ are just the polynomials f_1, \dots, f_k evaluated in the quantum ring associated to $[\omega]$ (Theorem 2.2). The $f_i^{[\omega]}$ are real-analytic in $[\omega]$ and thus have a natural analytic extension to $H^{1,1}(M)$, and the quantum cohomology rings fit together into a flat analytic family over $H^{1,1}(M)$.

As an application of this observation we shall compute the quantum cohomology of the Grassmannians. The calculations for $G(k, n)$ reduce to the single quantum product $c_k \wedge_{\mathbb{Q}} s_{n-k}$ of the top non-vanishing Chern respectively Segre class of the tautological k -bundle. We derive in Chapter 3:

THEOREM 0.1. *Let S be the tautological bundle over the Grassmann manifold $G(k, n)$ of complex k -planes in \mathbb{C}^n , $[\omega] = -\lambda \cdot c_1(S)$ a $(1, 1)$ -class on $G(k, n)$ ($\lambda \in \mathbb{R}_{>0} \Leftrightarrow [\omega]$ Kähler). Then*

$$QH_{[\omega]}^*(G(k, n)) = \mathbb{C}[X_1, \dots, X_k]/(Y_{n-k+1}, \dots, Y_{n-1}, Y_n + (-1)^k e^{-\lambda}),$$

where X_i corresponds to the i -th Chern class $c_i(S)$ and the Y_j (corresponding to the j -th Segre class of S) are given recursively by ($X_i = 0$ for $i > k$)

$$Y_j = -Y_{j-1} \cdot X_1 - \dots - Y_1 \cdot X_{j-1} - X_j.$$

This is in fact the form previously derived by Vafa using arguments from Quantum Field Theory [Va1], [Va2] (but note the sign of the quantum contribution). These

*Received September 22, 1997; accepted for publication November 25, 1997.

[†]Fakultät für Mathematik, Ruhr-Universität Bochum, D-44780 Bochum, Germany (Bernd.Siebert@ruhr-uni-bochum.de).

[‡]Department of Mathematics, MIT, Cambridge, MA 02139–4307, USA (tian@math.mit.edu).

papers and [In] also contain an intriguing formula for all GW-invariants (of the type considered here) of Grassmannians. We will see in Chapter 4 that formulas of this type occur whenever the cohomology ring has a presentation as complete intersection. In particular, we prove as Corollary 4.6

THEOREM 0.2. (Vafa-Intriligator formula) *For any Kähler class $[\omega]$ there is a finite set $C \subset \mathbb{C}^n$ and non-zero constants a_x , $x \in C$, such that for any $F \in \mathbb{C}[X_1, \dots, X_k]$*

$$\langle F \rangle_g^{[\omega]} = \sum_{x \in C} (a_x)^{g-1} \cdot F(x).$$

Here $\langle F \rangle_g^{[\omega]}$ is the genus g GW-invariant associated to F . In fact, there is a polynomial $W^{[\omega]}$ in the X_i having C as its set of critical points with a_x being (up to sign) the determinants of the Hessians at $x \in C$.

Some remarks on history and related works are in order. One of our starting points was a paper by Bertram, Daskalopoulos and Wentworth giving a mathematical formulation of the Vafa-Intriligator formula in terms of intersection theory on certain compactifications of moduli spaces of maps from a (fixed) Riemann surface to $G(k, n)$ [BDW]. They verified the formula for genus one and $k = 2$. The content of the first three chapters was known to the authors around New Year 93/94 and a preliminary version of the paper was spread in January 1994. A preprint version of the full paper was made available to the public as `alg-geom/940310`. The presentation theorem for quantum cohomology rings has been observed independently in [AsSa]. Some of the considerations of Chapter 2 appear also in [Pi]. We also realized later that the quantum product $c_k \wedge s_{n-k}$ had already been computed in [Wi3], up to a check of genericity conditions. His use of residues for the interpretation of coefficients of quantum products inspired us to the arguments in Chapter 4. The main technical result there (Proposition 4.1) about higher dimensional residues has independently been obtained in [CaDiSt].

Further partial support for the Vafa-Intriligator formula was obtained in [RRW] and [Bel]. The quantum cohomology of (absolute) flag manifolds (containing $G(k, n)$ as special case) has been calculated in [GiKi] (complete flags) and [AsSa] (general case) *assuming* the existence of an equivariant version of quantum cohomology with certain functorial properties. A description (and a proof of associativity) of the quantum multiplication for Grassmannians in terms of Schubert cycles (“quantum Pieri” and “quantum Giambelli” formulae) has later been obtained by Bertram [Be2].

By the effort of several people the quantum cohomology of generalized flag manifolds G/B have more recently been computed in complete generality, cf. [Ki] and the references given there.

The first named author thanks the DFG for support and the Courant Institute for hospitality during the academic year 1993/1994 where this work has been done. The second author is partly supported by an NSF grant and an Alfred P. Sloan Fellowship.

1. Definition of quantum multiplications. In this section, we recall the definition of quantum multiplications for positive manifolds as given in [RuTi]. The definition uses the GW-invariants from [Ru].

A symplectic manifold (M, ω) of dimension $2n$ is called *positive* if for any $f : \mathbb{P}^1 \rightarrow M$ with $[\omega](R) > 0$ it holds $c_1(M)(R) > 0$, $R = f_*[\mathbb{P}^1] \in H_2(M, \mathbb{Z})$. Example: Fano manifolds are positive with respect to a Kähler form representing $c_1(M)$. Let J

be an almost complex structure on M . A (parametrized) J -holomorphic curve on M is a smooth map $f : \Sigma \rightarrow M$ from a Riemann surface Σ satisfying

$$J \circ Df = Df \circ j,$$

as morphisms from T_Σ to f^*T_M , where j is the complex structure of Σ . This equation is a Cauchy-Riemann equation

$$\bar{\partial}_J f = 0, \quad \bar{\partial} = \frac{1}{2}(D + J \circ D \circ j).$$

For a technical reason one also needs the inhomogeneous equation $\bar{\partial}_J f = \nu$ with ν a section of $\Lambda^{0,1} \boxtimes_{\mathbb{C}} T_M$ over $\Sigma \times M$, $\Lambda^{0,1}$ the bundle of $(0, 1)$ -forms on Σ . Solutions of this equation are called perturbed or (J, ν) -holomorphic curves. The symplectic structure comes in by the assumption that J be *tamed* by ω which by definition means

$$\omega(X, JX) > 0 \quad \forall X \in T_M \setminus \{0\}.$$

Without this assumption moduli spaces of J -holomorphic curves can have rather bad non-compactness behaviour. The tameness condition is crucial in both defining GW-invariants and showing that they are invariants of the symplectic structure.

GW-invariants depend on the choice of $R \in H_2(M, \mathbb{Z})$ and integral homology classes $B_1, \dots, B_s \in H_*(M, \mathbb{Z})$ satisfying (g is the genus of Σ):

$$\sum_{i=1}^s (2n - \deg B_i) = 2c_1(M)(R) + 2n(1 - g) =: d(M, R, g). \quad (\dim)$$

Every integral homology class can be represented by so-called pseudo-manifolds (compact subspaces with singularities in real codimension at least 2, cf. [RuTi, Def.5.1]). For simplicity, we shall also use B_i to denote the pseudo-manifolds representing these homology classes. Let $t_1, \dots, t_s \in \Sigma$ be fixed points. It is shown in [RuTi] that if (J, ν) is generic the moduli space $\mathcal{M}_R(\Sigma, J, \nu)$ of (J, ν) -holomorphic maps of homology class R is smooth of dimension $d(M, R, g)$ (\dim) and the boundary of the image of the s -fold evaluation map

$$\text{ev} : \mathcal{M}_R(\Sigma, J, \nu) \rightarrow M^s, \quad f \mapsto (f(t_1), \dots, f(t_s))$$

has lower dimension. Choosing B_i transversal to the evaluation map and to the compactifying boundary there are only finitely many $f \in \mathcal{M}_R(\Sigma, J, \nu)$ with $f(t_i) \in B_i$. A GW-invariant is defined by

$$\tilde{\Phi}_{(R, \omega)}^g(B_1, \dots, B_s) := \sharp\{f \in \mathcal{M}_R(\Sigma, J, \nu) \mid f(t_i) \in B_i\},$$

the algebraic sum of such f counted with appropriate signs according to the canonically given orientation. One can prove that $\tilde{\Phi}_{(R, \omega)}^g(B_1, \dots, B_s)$ is independent of the choices of J, j, ν , and pseudo-manifolds representing B_1, \dots, B_s . As a matter of notation we set $\tilde{\Phi}_{(R, \omega)}^g(B_1, \dots, B_s) = 0$ unless the dimensions match (\dim).

To describe the compactifying set one generalizes the notion of (J, ν) -holomorphic maps to singular domains. Namely, one replaces Σ by a connected complex curve C with at most ordinary double points consisting of Σ and a number of spheres. One requires the spheres to be attached in such a way that the arithmetic genus of C still equals g . So C is obtained from Σ by attaching *trees* of spheres at points $P_a \in \Sigma$. A

(J, ν) -holomorphic map from C to M is then a continuous map $f : C \rightarrow M$ that is (J, ν) -holomorphic on Σ and J -holomorphic on each of the spheres. The compactifying set is obtained by the image of the obvious extension of the evaluation map, where the t_i may now be replaced by any point on a tree of spheres whenever the tree is attached at the original t_i .

We now make a simple but important remark: Let J be an almost complex structure such that any J -holomorphic curve with $f_*[C] = R$ is regular, i.e. $C = \Sigma$ is irreducible and the linearization of the Cauchy-Riemann operator at f is surjective. It follows from the construction of [Ru], [RuTi] that one can use exact J -holomorphic curves to compute the invariants $\tilde{\Phi}_{(R, \{\omega\})}^g$, cf. [RuTi, Rem.2.8]. In case J is integrable and f is an immersion, regularity at f is equivalent to the vanishing of $H^1(\Sigma, f^*T_M/T_\Sigma)$ [Gr, 2.1.B]. We adopt the convention that a *rational curve* in a complex manifold is a holomorphic curve (not necessarily smooth or irreducible) of arithmetic genus 1. The statement used for computation of GW-invariants in this paper is:

LEMMA 1.1. *Let (M, ω) be a Kähler manifold and $R \in H_2(M, \mathbb{Z})$ such that any rational curve $C \subset M$ homologous to R is non-singular and has $H^1(C, N_{C|X}) = 0$. Let $B_1, B_2, B_3 \subset M$ be complex submanifolds transversal to the (3-fold) evaluation map. Then*

$$\tilde{\Phi}_{(R, \{\omega\})}^0(B_1, B_2, B_3) = \sum_C (B_1 \cap C) \cdot (B_2 \cap C) \cdot (B_3 \cap C),$$

where the sum is over all rational curves C homologous to R .

Proof. Let I be the integrable complex structure of M . Then $\mathcal{M}_R(\mathbb{P}^1, I, 0)$ is nothing but the complex space $\text{Hom}_R(\mathbb{P}^1, M)$ of holomorphic maps $\mathbb{P}^1 \rightarrow M$ which is thus smooth. And by the Gromov compactness theorem and the assumption on smoothness of all rational curves representing R , $\mathcal{M}_R(\mathbb{P}^1, I, 0)$ is compact. Furthermore $(B_1 \cap C) \dots (B_3 \cap C)$ is precisely the number of ways to parametrize C by $f : \mathbb{P}^1 \rightarrow C$ with three distinguished points $t_i \in \mathbb{P}^1$ mapping to $B_i, i = 1, 2, 3$. Finally there are no signs occurring in the formula since all spaces involved are canonically oriented by their complex structures. \square

To define a quantum multiplication on $H^*(M, \mathbb{Z})$, we introduce the real-valued invariant

$$\tilde{\Phi}_{[\omega]}^g(B_1, \dots, B_s) = \sum_{R \in H_2(M, \mathbb{Z})} \tilde{\Phi}_{(R, \{\omega\})}^g(B_1, \dots, B_s) e^{-[\omega](R)}.$$

In general there might be infinitely many terms contributing to the sum (e.g. in the important and interesting case of Calabi-Yau manifolds) and one faces a non-trivial convergence problem. In the positive case, however, this sum is actually finite due to the dimension condition (dim). Let us assume M positive for the following (but see Remark 2.3). Note also that letting ω vary one gets back all the invariants $\tilde{\Phi}_{(R, \{\omega\})}^g(B_1, \dots, B_s)$ by solving some system of linear equations. The quantum multiplication $\wedge_{\mathbb{Q}}$ on $H^*(M, \mathbb{R})$ is then characterized by the equation

$$(\alpha \wedge_{\mathbb{Q}} \beta)[A] = \tilde{\Phi}_{[\omega]}^0(\alpha^\vee, \beta^\vee, A),$$

where $\alpha, \beta \in H^*(M, \mathbb{Z})$ and $^\vee$ means Poincaré-dual. In terms of a basis $\{A_i\}$ for $H_*(M, \mathbb{Z})$ modulo torsion and with $\{\alpha_i\}$ the Poincaré-dual basis of $H^*(M, \mathbb{Z})$ we may

state this more explicitly as

$$\alpha_i \wedge_Q \alpha_j = \sum_{k,l} \eta^{lk} \tilde{\Phi}_{[\omega]}^0(A_i, A_j, A_k) \alpha_l,$$

with $(\eta^{lk})_{kl}$ inverse to the intersection matrix $(\eta_{ij}) = (A_i \cdot A_j)$ ($\sum_j \eta_{ij} \eta^{jk} = \delta_{ik}$). The associativity of the quantum multiplication is highly non-trivial and shown by a careful analysis of the degeneration of rational curves in [RuTi] (following an idea of Witten).

An obvious, whence decisive feature of this definition, that we are going to exploit, is that its homogeneous part of highest degree reduces to the cup product

$$\alpha_i \wedge \alpha_j = \sum_{k,l} \eta^{lk} \cdot (A_i \cdot A_j \cdot A_k) \alpha_l.$$

2. A presentation for quantum cohomology. For a commutative ring A and degrees d_i attached to variables X_i we denote by $A\langle X_1, \dots, X_n \rangle$ the free graded commutative A -algebra with generators X_i of degree d_i , i.e. with $X_i X_j = (-1)^{d_i \cdot d_j} X_j X_i$. If m of the X_i have odd degree this is isomorphic to $\Lambda^\bullet A^m \otimes S^\bullet A^{n-m}$. We will call elements of this algebra ordered polynomials (since in addition to the coefficients one has to select an order among the factors in a monomial to determine its sign).

Let (M, ω) be a positive symplectic manifold and

$$H^*(M, \mathbb{Q}) = \mathbb{Q}\langle X_1, \dots, X_n \rangle / (f_1, \dots, f_k)$$

be a homogeneous presentation of the cohomology ring, $f_i = \sum_{|J|=\deg f_i} a_{iJ} X^J$. We use multiindex notation $J = (j_1, \dots, j_n)$, $X^J = X_1^{j_1} \wedge \dots \wedge X_n^{j_n}$, $|J| = \sum_{i=1}^n j_i d_i$ etc. Denote by \wedge_Q the product in the quantum cohomology $QH_{[\omega]}^*(M)$. To distinguish clearly between calculations in $H^*(M)$ and in $QH_{[\omega]}^*(M)$, we use a hat to mark elements of the quantum cohomology. So $\hat{}$ might be thought of as a \mathbb{C} -linear map $\mathbb{C}\langle X_1, \dots, X_n \rangle \rightarrow QH_{[\omega]}^*(M)$. An ordered polynomial f may be evaluated on the generators of the quantum cohomology. We write $f(\hat{X}_1, \dots, \hat{X}_n)$ as in the commutative case. For instance, $\hat{X}^J := \hat{X}_1 \wedge_Q \dots \wedge_Q \hat{X}_1 \wedge_Q \dots \wedge_Q \hat{X}_n \wedge_Q \dots \wedge_Q \hat{X}_n$ with \hat{X}_ν occurring j_ν -often (note the difference to $(X^J)^\wedge$!).

LEMMA 2.1. $\hat{X}_1, \dots, \hat{X}_n$ generate $QH_{[\omega]}^*(M)$.

Proof. By induction on the degree. So assume $\hat{X}_1, \dots, \hat{X}_n$ generate $QH_{[\omega]}^*(M)$ up to degree $d - 1$. We want to show that the monomials $(X^J)^\wedge$, $|J| = d$, can be written as linear combinations of (quantum) products of the \hat{X}_ν . But by definition of the multiplication in $QH_{[\omega]}^*(M)$

$$\hat{X}^J = (X^J)^\wedge + \sum_{|I|<d} a_I (X^I)^\wedge,$$

and by induction hypothesis $(X^I)^\wedge = \sum_{K \leq I} b_{IK} \hat{X}^K$, so

$$(X^J)^\wedge = \hat{X}^J - \sum_{\substack{K \leq I \\ |I|<d}} a_I b_{IK} \hat{X}^K. \quad \square$$

Next we evaluate $f_i(\hat{X}_1, \dots, \hat{X}_n)$, i.e. in the quantum cohomology ring. Since f_i is a relation in the cohomology ring, the term of top-degree $\deg f_i$ vanishes. By

the lemma we find an ordered polynomial $g_i^{[\omega]}$ (depending on $[\omega]$) in n variables (say T_1, \dots, T_n , $\deg T_i = \deg X_i$) of lower degree with

$$f_i(\hat{X}_1, \dots, \hat{X}_n) = g_i^{[\omega]}(\hat{X}_1, \dots, \hat{X}_n) \quad \text{in } QH_{[\omega]}^*(M).$$

Thus $f_i^{[\omega]}(T_1, \dots, T_n) := f_i(T_1, \dots, T_n) - g_i^{[\omega]}(T_1, \dots, T_n) \in \mathbb{C}\langle T_1, \dots, T_n \rangle$ is a non-trivial relation between $\hat{X}_1, \dots, \hat{X}_n$.

PROPOSITION 2.2. $QH_{[\omega]}^*(M) = \mathbb{C}\langle T_1, \dots, T_n \rangle / (f_1^{[\omega]}, \dots, f_k^{[\omega]})$.

Proof. Let $\mathcal{J} \subset \mathbb{C}\langle T_1, \dots, T_n \rangle$ be the ideal of relations between the $\hat{X}_1, \dots, \hat{X}_n$. Then $(f_1^{[\omega]}, \dots, f_k^{[\omega]}) \subset \mathcal{J}$ and $QH_{[\omega]}^*(M) = \mathbb{C}\langle T_1, \dots, T_n \rangle / \mathcal{J}$ by the lemma. Let $F \in \mathcal{J} \setminus \{0\}$. Expand $F = F_d + F'$ with $F_d \neq 0$ (weighted) homogeneous of degree d , $d > 0$, and $\deg F' < d$. Then

$$F_d(\hat{X}_1, \dots, \hat{X}_n) = -F'(\hat{X}_1, \dots, \hat{X}_n)$$

in quantum cohomology, for F is a relation. But the highest degree ($\deg = d$) contribution to $F_d(\hat{X}_1, \dots, \hat{X}_n)$ is just $(F_d(X_1, \dots, X_n))^\wedge$. Its vanishing implies $F_d \in (f_1, \dots, f_k)$, i.e. $F_d = \varphi(f_1, \dots, f_k)$, φ a polynomial in k variables. Thus

$$\varphi(f_1^{[\omega]}, \dots, f_k^{[\omega]}) = F_d + F'' \quad \text{in } \mathbb{C}\langle T_1, \dots, T_n \rangle$$

with $\deg F'' < d$, and we may write $F = \varphi(f_1^{[\omega]}, \dots, f_k^{[\omega]}) + F' - F''$, $\deg F' - F'' < d = \deg F$. Proceeding by induction on the degree we finally see $F \in (f_1^{[\omega]}, \dots, f_k^{[\omega]})$, i.e. $\mathcal{J} = (f_1^{[\omega]}, \dots, f_k^{[\omega]})$ as claimed. \square

REMARK 2.3. 1) The proposition is constructive. The induction process shows that a particular coefficient of the perturbed relations $f^{[\omega]}$ may involve several GW-invariants. So the actual computation of $f^{[\omega]}$ is not always easy, even if one knows a good deal about the GW-invariants of M . The converse problem of determining GW-invariants from the presentation can be dealt with by Gröbner basis methods readily accessible to computer algebra programs, cf. Chapter 4.

2) In the positive case $c_1(M) > 0$ the contributions with $R \neq 0$ give rise to terms of lower degree by the dimension condition, so we were able to fix $[\omega]$ and argue by considerations on the degree. Modulo convergence problems mentioned at the end of the previous chapter, Proposition 2.2 remains valid in the semi-positive case. Arguments involving the degree can be replaced by linear independence of terms $e^{-\lambda \cdot t}$ for various λ in the algebra $\mathbb{C}\{t\}$.

Another approach, which works without any positivity condition, is to use formal power series over $H_2(M, \mathbb{Z})$ (cf. also [Pi]): Let $\mathcal{RC}(M) \subset H$ be the monoid generated by classes of J -holomorphic rational curves. Since J is also ω' -tame for ω' a symplectic form sufficiently close to ω and $\omega'(\mathcal{RC}(M)) \subset \mathbb{R}_{>0}$, $\mathcal{RC}(M)$ is a strictly convex cone. Multiplication on the group $N := \mathbb{Z} \llbracket \mathcal{RC}(M) \rrbracket$ of formal power series over $\mathcal{RC}(M)$ is thus well-defined. Then $H^*(M, \mathbb{Z}) \otimes_{\mathbb{Z}} N$ appears as natural domain of definition of quantum products for more general manifolds. Our arguments provide a presentation of the quantum cohomology ring as quotient of $\mathbb{Q}\langle T_1, \dots, T_n \rangle \otimes_{\mathbb{Q}} N$ by ordered polynomials with coefficients in N . Note also that in the positive case $QH_{[\omega]}^*(M)$ may be viewed as homeomorphic image of $H^*(M, \mathbb{Z}) \otimes_{\mathbb{Z}} \mathbb{Z}[\mathcal{RC}(M)] \subset H^*(M, \mathbb{Z}) \otimes_{\mathbb{Z}} N$ by sending $R \in H_2(M, \mathbb{Z})$ to $e^{-[\omega](R)}$. \square

3. Grassmann manifolds. As is well-known (e.g. [Fu, Ex. 14.6.6]), the cohomology ring (in fact even the Chow ring) of the Grassmann variety $G(k, n)$ of k -planes in \mathbb{C}^n has a presentation

$$H^*(G(k, n), \mathbb{Q}) = \mathbb{Q}[c_1, \dots, c_k] / (s_{n-k+1}, \dots, s_n),$$

with c_i corresponding to the Chern classes of the tautological k -bundle S and s_j to its Segre classes viewed as polynomials in c_1, \dots, c_k via

$$\begin{aligned} (1 + c_1 + \dots + c_k)(1 + s_1 + s_2 + \dots) &= 1, \\ \text{i.e. } s_j &= -s_{j-1}c_1 - \dots - s_1c_{j-1} - c_j. \end{aligned} \tag{*}$$

Note that s_j is also the j -th Chern class of the universal quotient bundle Q , which has rank $n - k$, so $s_j = 0$ in $H^*(G(k, n))$ for $j > n - k$. In fact, as a polynomial in c_i, s_j lies in the relation ideal for $j > n$ by the recursion formula (*). Under the canonical isomorphism $\Phi : G(k, n) \simeq G(n - k, n), \Lambda \mapsto (\mathbb{C}^n / \Lambda)^*$, S corresponds to the dual of the universal quotient bundle Q' on $G(n - k, n)$ and Q to the dual of the tautological $(n - k)$ -bundle S' . Thus c_i and s_j exchange (up to sign) their roles and one might as well write

$$H^*(G(k, n), \mathbb{Q}) = \mathbb{Q}[s_1, \dots, s_{n-k}] / (c_{k+1}, \dots, c_n),$$

this time with c_i polynomials in s_1, \dots, s_{n-k} (this presentation is actually better adapted to Schubert calculus, i.e. geometry, see below). These remarks are made to emphasize the symmetry between c_i and s_j . Note also that the generators all have even degree, so we need not worry about questions of sign.

Schubert calculus (e.g. [Fu, § 14.7]) provides a basis of $H^*(G(k, n))$ as \mathbb{Q} -vector space (indeed a basis of integral homology/cohomology as \mathbb{Z} -module), indexed by tuples $(\lambda_1, \dots, \lambda_k), n - k \geq \lambda_1 \geq \dots \geq \lambda_k \geq 0$, via

$$\{\lambda_1, \dots, \lambda_k\} := \det(s_{\lambda_i + j - i})_{1 \leq i, j \leq k},$$

i.e. by evaluating the Schur polynomial (S-function) associated to $(\lambda_1, \dots, \lambda_k)$ on the Segre classes of the tautological k -bundle ($s_j = 0$ for $j \notin \{0, \dots, n - k\}$). $\{\lambda_1, \dots, \lambda_k\}$ is (weighted) homogeneous of degree $2 \sum \lambda_i$. The Chern respectively Segre classes are given by

$$\begin{aligned} c_i &= (-1)^i \{1, \dots, 1, 0, \dots, 0\} \quad (\text{"1" } i\text{-times}), \\ s_j &= \{j, 0, \dots, 0\} \end{aligned}$$

(for c_i see [Fu, Lemma 14.5.1] or do an easy induction). The connection to classical Schubert calculus is given by Poincaré-duality. In fact, $\{\lambda_1, \dots, \lambda_k\} \cap [G(k, n)]$ may be represented by the Schubert varieties

$$\Omega_{\underline{V}}(n - k + 1 - \lambda_1, \dots, n - k + i - \lambda_i, \dots, n - \lambda_k).$$

To define the latter one has to fix a flag $\underline{V} = (V_1, \dots, V_n), V_1 \subset \dots \subset V_n = \mathbb{C}^n$ of linear subspaces of $\mathbb{C}^n, \dim V_i = i$. Then for $(a_1, \dots, a_k), 0 \leq a_1 < \dots < a_k \leq n$

$$\Omega_{\underline{V}}(a_1, \dots, a_k) := \{\Lambda \in G(k, n) \mid \dim \Lambda \cap V_{a_i} \geq i, 1 \leq i \leq k\}.$$

The homology class $(a_1, \dots, a_n) := [\Omega_{\underline{V}}(a_1, \dots, a_k)]$ is independent of the choice of flag. We write $\{\underline{\lambda}\}^\vee := (n - k + 1 - \lambda_1, \dots, n - k + i - \lambda_i, \dots, n - \lambda_k) \in H_*(G(k, n))$.

What is important for us is that $[G(k, n)]$ is Poincaré-dual to $\{0, \dots, 0\} = 1$ (trivial) and that the class of a point $[*]$ is Poincaré-dual to $\{n - k, \dots, n - k\} = s_{n-k}^k = (-1)^{n-k} c_k^{n-k}$ (which makes sense as $\dim G(k, n) = k(n - k)$, but is less trivial: What is the Poincaré-dual of $c_1^{k(n-k)}$?). The intersection of classes of complementary dimension is especially easy: $|\lambda| + |\mu| = k(n - k)$, then (“duality theorem”):

$$(\{\lambda\} \wedge \{\mu\}) [G] = \{\lambda\} \cap \{\mu\}^\vee = \{\lambda\}^\vee \cdot \{\mu\}^\vee = \begin{cases} 1 & , \text{ if } \underline{\mu} = \underline{\lambda}^* \\ 0 & , \text{ otherwise.} \end{cases}$$

with $\underline{\lambda}^* := (n - k - \lambda_k, \dots, n - k - \lambda_1)$. In particular, we get $\eta_{\underline{\lambda}\underline{\mu}} = \delta_{\underline{\lambda}\underline{\mu}}$ for the intersection matrix. This ends our collection of facts concerning Grassmannians.

We thus see that $H^{1,1}(G(k, n))$ is spanned by the single class $\{1, 0, \dots, 0\} = s_1 = -c_1$. The ample generator is $s_1 = c_1(Q)$ for $s_1 = \iota^* c_1(\mathcal{O}_{\mathbb{P}(\Lambda^k \mathbb{C}^n)}(1))$ where $\iota : G(k, n) \hookrightarrow \mathbb{P}(\Lambda^k \mathbb{C}^n)$ is the Plücker embedding. Dually, $H_2(G(k, n), \mathbb{Z})$ is spanned by the class of a line $\{n - k, \dots, n - k, n - k - 1\}^\vee = (1, \dots, k - 1, k + 1) =: [L]$. Since $T_{G(k, n)} \simeq \text{Hom}(S, Q) \simeq S^* \otimes Q$, as one easily verifies by using the standard local coordinates on $G(k, n)$, $c_1(G(k, n)) = \text{rk}(Q) \cdot c_1(S^*) + \text{rk}(S^*) \cdot c_1(Q) = n \cdot s_1$.

So $G(k, n)$ is a Fano manifold of index n with Picard group infinite cyclic of rank one and it makes sense to talk about its quantum cohomology in the version presented in Chapter 1. The product in $QH_{[\omega]}^*(G(k, n))$ reads

$$\widehat{\{\lambda\}} \wedge_Q \widehat{\{\mu\}} = \sum_{R=d \cdot [L], d \geq 0} \sum_{\nu} \tilde{\Phi}_R^0(\{\lambda\}^\vee, \{\mu\}^\vee, \{\nu\}^\vee) \widehat{\{\nu^*\}} e^{-[\omega](R)}. \tag{**}$$

Recall from Chapter 1 that $\tilde{\Phi}_{d \cdot [L]}^0(\{\lambda\}^\vee, \{\mu\}^\vee, \{\nu\}^\vee) = 0$ unless $|\lambda| + |\mu| + |\nu| = k(n - k) + c_1(G(k, n))(d \cdot [L]) = k(n - k) + d \cdot n$. In calculating $s_{n-k+i}(\hat{c}_1, \dots, \hat{c}_k)$ (i.e. in $QH_{[\omega]}^*(G(k, n))$), \hat{c}_i the generators corresponding to c_i as in Chapter 2) we have $|\lambda| + |\mu| \leq n$, and “=” only in the case of s_n . But $|\nu| \leq k(n - k) = \dim G(k, n)$, so we get no quantum contributions except in case of s_n with $|\nu| = k(n - k)$ and $d = 1$. Thus writing $\hat{c} = (\hat{c}_1, \dots, \hat{c}_k)$ we obtain $s_{n-k+1}(\hat{c}) = \dots = s_{n-1}(\hat{c}) = 0$ in $QH_{[\omega]}^*(G(k, n))$ and

$$\begin{aligned} s_n(\hat{c}) &= -\hat{c}_1 s_{n-1}(\hat{c}) - \dots - \hat{c}_{k-1} s_{n-k+1}(\hat{c}) - \hat{c}_k s_{n-k}(\hat{c}) \\ &= -\hat{c}_k s_{n-k}(\hat{c}) = -\tilde{\Phi}_{[L]}^0(c_k^\vee, s_{n-k}^\vee, [*]) \cdot e^{-[\omega](L)}, \end{aligned}$$

where for the last equality we have used (**). By Proposition 2.2, to prove Theorem 0.1, we are left with

PROPOSITION 3.1. $\tilde{\Phi}_{[L]}^0(c_k^\vee, s_{n-k}^\vee, [*]) = (-1)^k$.

For the proposition we first have to classify holomorphic curves homologous to $[L] = (1, \dots, k - 1, k + 1)$.

LEMMA 3.2. *Let $C \subset G(k, n)$ be a (rational) curve homologous to $(1, \dots, k - 1, k + 1)$. Then C is a Schubert variety $\Omega_V(1, \dots, k - 1, k + 1)$, i.e. there are linear subspaces $U \subset W \subset \mathbb{C}^n$, $\dim U = k - 1$, $\dim W = k + 1$, with $C = \{\Lambda \in G(k, n) \mid U \subset \Lambda \subset W\}$.*

Proof. Since $s_1[C] = 1$ by the duality theorem, $\deg \iota(C) = 1$, so the image of C under the Plücker embedding $\iota : G(k, n) \rightarrow \mathbb{P}(\Lambda^k \mathbb{C}^n)$ is a linear \mathbb{P}^1 . The image of $G(k, n)$ consists precisely of (rays of) decomposable vectors $v_1 \wedge \dots \wedge v_k = \iota(\langle v_1, \dots, v_k \rangle) \in \Lambda^k \mathbb{C}^n$. Locally, the vectors v_1, \dots, v_k may be chosen to vary smoothly with $\Lambda \in G(k, n)$: In fact, the standard (affine) coordinate neighbourhood of $\Lambda \in G(k, n)$ is

$\text{Hom}(\Lambda, \mathbb{C}^n/\Lambda)$ with $\Psi : \text{Hom}(\Lambda, \mathbb{C}^n/\Lambda) \rightarrow \mathbf{G}(k, n)$, $\varphi \mapsto \langle v + \varphi(v) \mid v \in \Lambda \rangle$ (in particular $0 \in \text{Hom}(\Lambda, \mathbb{C}^n/\Lambda)$ corresponds to Λ). Now fixing a basis v_1, \dots, v_k of Λ , $\iota \circ \Psi$ may be represented (lifted to $\Lambda^k \mathbb{C}^n$) by

$$\varphi \mapsto (v_1 + \varphi(v_1)) \wedge \dots \wedge (v_k + \varphi(v_k)) .$$

Thus choosing $\Lambda' \in C$ sufficiently close to $\Lambda \in C$ and letting e_1, \dots, e_l be a basis of $\Lambda \cap \Lambda'$, completed by e_{l+1}, \dots, e_k and e'_{l+1}, \dots, e'_k to a basis of Λ and Λ' respectively, we have for $t \in \mathbb{C}$ small

$$e_1 \wedge \dots \wedge e_k + t \cdot e_1 \wedge \dots \wedge e_l \wedge e'_{l+1} \wedge \dots \wedge e'_k = v_1(t) \wedge \dots \wedge v_k(t)$$

with $v_i(0) = e_i$ by construction. Taking $\left. \frac{d}{dt} \right|_{t=0}$ yields

$$e_1 \wedge \dots \wedge e_l \wedge e'_{l+1} \wedge \dots \wedge e'_k = \dot{v}_1(0) \wedge e_2 \wedge \dots \wedge e_k + \dots + e_1 \wedge \dots \wedge e_{k-1} \wedge \dot{v}_k(0) .$$

As one sees by expanding $\dot{v}_i(0)$ in terms of a basis of \mathbb{C}^n containing $\{e_1, \dots, e_k, e'_{l+1}, \dots, e'_k\}$ we may gather the linearly independent terms with and without e_k to form two equations. The left-hand side of the equation above belongs to the latter ($\Lambda \neq \Lambda' \Rightarrow l < k$), so we get

$$e_1 \wedge \dots \wedge e_l \wedge e'_{l+1} \wedge \dots \wedge e'_k = e_1 \wedge \dots \wedge e_{k-1} \wedge (\dot{v}_k(0) - \lambda \cdot e_k) ,$$

where $\lambda \in \mathbb{C}$ is chosen in such a way that $\dot{v}_k(0) - \lambda \cdot e_k$ lies in the span of the basis vectors different from e_k . By linear independence of wedge products of a basis of \mathbb{C}^n this shows $l = k - 1$ and $e'_k = \dot{v}_k(0) - \lambda e_k$. In view of the linearity of $\iota(C)$ we conclude

$$\iota(C) = \left\{ [t \cdot e_1 \wedge \dots \wedge e_{k-1} \wedge e'_k + u \cdot e_1 \wedge \dots \wedge e_{k-1} \wedge e_k] \mid [t : u] \in \mathbb{P}^1 \right\} ,$$

so C is the Schubert variety $\Omega_{\underline{V}}(1, \dots, k - 1, k + 1)$ belonging to a flag \underline{V} with $V_{k-1} = \langle e_1, \dots, e_{k-1} \rangle =: U$ and $V_{k+1} = \langle e_1, \dots, e_{k-1}, e_k, e'_k \rangle =: W$. \square

LEMMA 3.3. Let $A_1 = \Omega_{\underline{V}^1}(n - k, n - k + 1, \dots, n - 1) = \{1, \dots, 1\}^\vee = (-1)^k c_k^\vee$, $A_2 = \Omega_{\underline{V}^2}(1, n - k + 2, \dots, n) = \{n - k, 0, \dots, 0\}^\vee = s_{n-k}^\vee$, $A_3 = \{*\} = \Omega_{\underline{V}^3}(1, \dots, k) = \{n - k, \dots, n - k\}^\vee$, where $\underline{V}^1, \underline{V}^2, \underline{V}^3$ are three transversal flags, i.e. $\dim V_i^1 \cap V_j^2 \cap V_k^3 = \max\{0, i + j + k - 2n\}$ for any i, j, k .

Then there is exactly one rational curve C homologous to $[L]$ and with $C \cap A_i \neq \emptyset$, $i = 1, 2, 3$. Moreover, $C \cdot A_1 = C \cdot A_2 = C \cdot A_3 = 1$.

The lemma can be proved by doing intersection theory on the flag manifold $F(k - 1, k + 1; n)$ which parametrizes rational curves of minimal degree by the preceding lemma, and using the two obvious maps $\pi : F(k - 1, k + 1; n) \rightarrow \mathbf{G}(k, n)$ and $p : F(k - 1, k + 1; n) \rightarrow \mathbf{G}(k, n)$ (calculate $(p_* \pi^*[A_1]) \cdot (p_* \pi^*[A_2]) \cdot (p_* \pi^*[A_3])$). This method might be appropriate for more general A_1, A_2, A_3 , yet in our case an explicit linear algebra argument is simpler and even more enlightening.

Proof. We need to find three k -planes $\Lambda^1, \Lambda^2, \Lambda^3$ and subspaces $U, W \subset \mathbb{C}^n$, $\dim U = k - 1$, $\dim W = k + 1$ with

1. $U \subset \Lambda^1 \cap \Lambda^2 \cap \Lambda^3 \subset \Lambda^1 + \Lambda^2 + \Lambda^3 \subset W$.
2. $\dim \Lambda^1 \cap V_{n-k+(i-1)}^1 \geq i$, $i = 1, \dots, k$.
3. $V_1^2 \subset \Lambda^2$.
4. $\Lambda^3 = V_k^3$,

(1) says that $\Lambda^1, \Lambda^2, \Lambda^3$ lie on the rational curve C defined by U and W according to Lemma 3.2, whereas (2)–(4) rephrase the conditions $\Lambda_i = A_i \cap C$, $i = 1, 2, 3$. We now use transversality of the flags V^1, V^2, V^3 : From (3), (4) and (1) we readily deduce $W = V_k^3 + V_1^2$, and (2) with $i = k$ shows $\Lambda^1 \subset V_{n-1}^1$, so by (4) and (1) we get $U = V_k^3 \cap V_{n-1}^1$. This choice of U, W implies $\Lambda^1 = W \cap V_{n-1}^1$, $\Lambda^2 = U + V_1^2$, $\Lambda^3 = V_k^3$. Conversely, these $\Lambda^1, \Lambda^2, \Lambda^3$ fulfill (1)–(4). \square

Proof of Proposition 3.1. Let I be the integrable complex structure on $G(k, n)$. Consider the evaluation map

$$ev : \Sigma \times \mathcal{M}_{[L]}(\mathbb{P}^1, I, 0) \longrightarrow G(k, n) \times G(k, n) \times G(k, n).$$

Since the tangent bundle of $G(k, n)$ is generated by global sections, so is $N_{L/G(k, n)}$ for any line L . Hence $H^1(N_{L/G(k, n)}) = 0$ for any such line and $\mathcal{M}_{[L]}(\mathbb{P}^1, I, 0) = \text{Hom}_{[L]}(\mathbb{P}^1, M)$ is a manifold of expected dimension $n + k(n - k)$ (in fact a $\text{PGL}(2)$ -bundle over $F(k - 1, k + 1; n)$). By the preceding lemma the preimage under ev of $A_1 \times A_2 \times A_3$ for any A_i belonging to transversal flags consists of exactly one $f \in \mathcal{M}_{[L]}(\mathbb{P}^1, I, 0)$. But letting the transversal flags vary, the variation of $ev(f) \cap A_1 \times A_2 \times A_3$ sweeps out the normal directions of $A_1 \times A_2 \times A_3$ in $G(k, n)^3$ as follows explicitly from the proof of Lemma 3.3. \square

This finishes the proof of Theorem 0.1.

REMARK 3.4. (ON ALGEBRAICITY) While we use the symplectic definition of GW-invariants here, our arguments are essentially algebraic. In fact, the computation of the invariant $\tilde{\Phi}_{[L]}^0(c_k^\vee, s_{n-k}^\vee, [*])$ is done by ordinary intersection theory on a moduli space of algebraic lines in $G(k, n)$. And in a joint work of the second named author with J. Li it is shown by purely algebraic methods that for homogeneous varieties the “naive”, algebraic, genus zero invariants, defined by intersection theory on algebraic moduli spaces, also obey associativity [LiTi] (and coincide with the symplectic GW-invariants). So these symplectic GW-invariants are “enumerative” in the sense that they count numbers of actual holomorphic curves.

For the Grassmannians analogous results have been given independently by [Be1]. In this paper a different compactification of $\text{Hom}(\Sigma, G(k, n))$ is used (by a quot scheme). The author also shows enumerativeness of the invariants for curves of higher genus and sufficiently high degree, and the induction formula for the genus (Proposition 4.4 below).

More recently, an enumerative theory has been worked out by Behrend and Manin for genus zero invariants and so-called *convex* projective manifolds by using Kontsevich’s very pretty concept of “stable maps” [BeMa]. A smooth variety M is defined to be convex if $H^1(\mathbb{P}^1, \varphi^*T_M) = 0$ for any $\varphi \in \text{Hom}(\mathbb{P}^1, M)$. This includes homogeneous varieties for these have globally generated tangent bundles.

In all these examples $\text{Hom}(\mathbb{P}^1, M)$ is smooth of the expected dimension. The boundary being of lower dimension, intersection theory on any compactification (or of the image of the moduli space in M^s) gives the same result. \square

4. Formulae of Vafa-Intriligator type. The main results of [RuTi] show how to compute higher GW-invariants (i.e. with more than three entries or for higher genus Riemann surfaces) from the genus 0 three-point functions inductively. Namely, for $g > 0$

$$\tilde{\Phi}_{[\omega]}^g(B_1, \dots, B_s) = \sum_{i, j} \eta^{ij} \tilde{\Phi}_{[\omega]}^{g-1}(B_1, \dots, B_s, A_i, A_j),$$

(η_{ij}) the intersection matrix with respect to a basis $\{A_i\}$ of $H^*(M, \mathbb{Q})$. More invariantly, the right-hand side is the trace with respect to η of the bilinear form

$$H^*(M, \mathbb{C}) \times H^*(M, \mathbb{C}) \longrightarrow \mathbb{C}, \quad (B', B'') \longmapsto \tilde{\Phi}_{[\omega]}^{g-1}(B_1, \dots, B_s, B', B'').$$

Secondly, for $g = 0$ and $1 < r < s - 1$ (otherwise trivial) we have the *composition law*

$$\tilde{\Phi}_{[\omega]}^0(B_1, \dots, B_s) = \sum_{i,j} \eta^{ij} \tilde{\Phi}_{[\omega]}^0(B_1, \dots, B_r, A_i) \cdot \tilde{\Phi}_{[\omega]}^0(A_j, B_{r+1}, \dots, B_s),$$

a trace with respect to η as well. For $s = 4$ and $r = 2$ this equation states the associativity of quantum products. Our goal in this section is to give a closed formula for higher invariants in terms of the relations $f_1^{[\omega]}, \dots, f_k^{[\omega]}$ of the quantum cohomology ring.

Putting all $\tilde{\Phi}_{[\omega]}^g$ together for different s we get a \mathbb{C} -linear map

$$\langle \rangle_g : \mathbb{C}\langle X_1, \dots, X_n \rangle \longrightarrow \mathbb{C}, \quad X_1^{\nu_1} \dots X_n^{\nu_n} \longmapsto \tilde{\Phi}_{[\omega]}^g(X_1^\vee, \dots, X_1^\vee, \dots, X_n^\vee, \dots, X_n^\vee),$$

the Poincaré-dual X_i^\vee of X_i occurring ν_i -times. But from the composition law, letting α_i denote the Poincaré-dual of A_i ,

$$\hat{X}_1^{\nu_1} \wedge_{\mathbb{Q}} \dots \wedge_{\mathbb{Q}} \hat{X}_n^{\nu_n} = \sum_{ij} \eta^{ij} \tilde{\Phi}_{[\omega]}^0(X_1^\vee, \dots, X_1^\vee, \dots, X_n^\vee, A_i) \hat{\alpha}_j,$$

so $\langle X_1^{\nu_1} \dots X_n^{\nu_n} \rangle_0 = \tilde{\Phi}_{[\omega]}^0(X_1^\vee, \dots, X_1^\vee, \dots, X_n^\vee, [M])$ is nothing but the coefficient of the class $[\Omega]$ of the normalized volume form in $\hat{X}_1^{\nu_1} \wedge_{\mathbb{Q}} \dots \wedge_{\mathbb{Q}} \hat{X}_n^{\nu_n}$. That is, $\langle \rangle_0$ factorizes in the following way:

$$\mathbb{C}\langle X_1, \dots, X_n \rangle \longrightarrow \mathbb{C}\langle X_1, \dots, X_n \rangle / (f_1^{[\omega]}, \dots, f_k^{[\omega]}) \simeq H_{[\omega]}^*(M) \xrightarrow{\text{top}} H^{2n}(M, \mathbb{C}),$$

when $H^{2n}(M, \mathbb{C})$ is identified with \mathbb{C} by sending $[\Omega]$ to 1. In case $n = k$ and the generators have even degree, i.e. $H^*(M, \mathbb{C})$ is a (commutative) complete intersection ring, one can use higher dimensional residues to express this map more explicitly (we refer the reader to [GrHa] and [Ts] for the general facts on residues to be used):

Recall that the residue in $a \in \mathbb{C}$ of $F \in \mathbb{C}[X_1, \dots, X_k]$ with respect to a polynomial mapping $g = (g_1, \dots, g_k) : \mathbb{C}^k \rightarrow \mathbb{C}^k$ with $g^{-1}(0)$ finite is defined by

$$\text{res}_g(a; F) := \frac{1}{(2\pi i)^k} \int_{\Gamma_a^\varepsilon} \frac{F}{g_1 \cdots g_k} dX_1 \dots dX_k,$$

with $\Gamma_a^\varepsilon = \{x \in U(a) \mid |g_i(x)| = \varepsilon\}$, $U(a)$ a neighbourhood of a with $g^{-1}(0) \cap U(a) = \{a\}$ and ε so small that Γ_a^ε lies relatively compact in $U(a)$. Γ_a^ε is smooth for almost all ε by Sard's Theorem and has a canonical orientation by the k -form $d(\arg g_1) \wedge \dots \wedge d(\arg g_k)|_{\Gamma_a^\varepsilon}$. (This local residue of course makes sense for holomorphic g and $F \in \mathcal{O}_a$, but the polynomial case, to which the general case may easily be reduced, is sufficient for our purposes). Note that the hypothesis of $g^{-1}(0)$ being finite is equivalent to the saying that the ring $\mathbb{C}[X_1, \dots, X_k]/(g_1, \dots, g_k)$ is Artinian, i.e. finite dimensional as vector space over \mathbb{C} . We define the total residue

$$\text{Res}_g(F) := \sum_{a \in g^{-1}(0)} \text{res}_g(a; F),$$

which is also known as Grothendieck residue symbol $\left(\frac{F}{g_1, \dots, g_k}\right)$ in the context of duality theory in algebraic geometry [Ha]. Let $J = \det\left(\frac{\partial g_i}{\partial X_j}\right)$ be the Jacobian of g . For regular values y of g

$$\text{Res}_{g=y}(F) = \sum_{x \in g^{-1}(y)} \left(\frac{F}{J}\right)(x) =: \text{tr}\left(\frac{F}{J}\right)(0).$$

Therefore $\text{tr}(F/J)$ extends holomorphically to a neighbourhood of 0 (the extension will be denoted $\text{tr}(F/J)$ as well) and

$$\text{Res}_g(F) = \text{tr}\left(\frac{F}{J}\right)(0).$$

One abstract feature in our setting is that we have weights d_i associated to X_i and that our relations $f_i^{[\omega]}$ form a Gröbner basis of the relation ideal with respect to these weights, i.e.

$$(\text{In } f_1^{[\omega]}, \dots, \text{In } f_k^{[\omega]}) = (f_1, \dots, f_k) = \text{In}(f_1^{[\omega]}, \dots, f_k^{[\omega]}),$$

where “In” means taking initial forms with respect to the weight order. This is trivial in our case since we started with the homogeneous generators f_1, \dots, f_k of the relation ideal in a presentation of $H^*(M, \mathbb{C})$. In such situations one can describe the residue map algebraically as follows:

PROPOSITION 4.1. *Let $R = \mathbb{C}[X_1, \dots, X_k]/(g_1, \dots, g_k)$ be Artinian with $\{g_i\}$ a Gröbner basis of the relation ideal with respect to weights d_i of X_i . Put $N := \sum_i \deg g_i - \sum_i d_i$, $R_{<N} := \{F \in \mathbb{C}[X_1, \dots, X_k] \mid \deg F < N\}/(g_1, \dots, g_k)$ and $J = \det\left(\frac{\partial g_i}{\partial X_j}\right)$. Then*

$$R = R_{<N} \oplus \mathbb{C} \cdot J,$$

and the total residue map $\text{Res}_g : \mathbb{C}[X_1, \dots, X_k] \rightarrow \mathbb{C}$ factorizes via the projection onto the second factor as follows

$$\mathbb{C}[X_1, \dots, X_k] \xrightarrow{\text{can}} R \xrightarrow{\text{pr}_2} \mathbb{C} \cdot J \longrightarrow \mathbb{C},$$

where the last map sends J to $\dim_{\mathbb{C}} R$.

Proof. To check the normalization we observe

$$\text{Res}_g(J) = (\text{tr}_g 1)(0) = \text{degree of } g \text{ over } 0 = \dim_{\mathbb{C}} R.$$

The last equality is generally true by flatness if the covering space is Cohen-Macaulay, cf. eg. [Fi]. Next, it is well known that the residue vanishes on elements of (g_1, \dots, g_k) . The claim thus reduces to $\ker(\text{Res}_g)/(g_1, \dots, g_k) = R_{<N}$.

If $\deg F < N$ then $F dX_1 \dots dX_k/g_1 \dots g_k$ extends to a rational differential form φ on the weighted projective space $V = \mathbb{P}_{(1, d_1, \dots, d_k)}$ with polar divisor $D_1 + \dots + D_k \in \text{Div}(V)$, D_i the natural extension of the divisor (g_i) to V . The point of course is that the divisor at infinity $V \setminus \mathbb{C}^k$ is not a polar divisor of φ . We claim $|D_1| \cap \dots \cap |D_k| \subset \mathbb{C}^k$. In fact, the restriction of the homogenization of g_i to $V \setminus \mathbb{C}^k \simeq \mathbb{P}_{(d_1, \dots, d_k)}$ is just $\text{In } g_i$ and $V(\text{In } g_1, \dots, \text{In } g_k) = \{0\} \in \mathbb{C}^k$. The latter follows because otherwise $\dim \text{Spec } \mathbb{C}[X_1, \dots, X_k]/(\text{In } g_1, \dots, \text{In } g_k) > 0$ by homogeneity. But from the Gröbner basis property we have

$$\dim_{\mathbb{C}} \mathbb{C}[X_1, \dots, X_k]/(\text{In } g_1, \dots, \text{In } g_k) = \dim_{\mathbb{C}} \mathbb{C}[X_1, \dots, X_k]/(g_1, \dots, g_k).$$

We may thus desingularize V (at infinity) without violating the discreteness of $|D_1| \cap \dots \cap |D_k|$. We keep the notations for the pulled-back objects. Now the global residue theorem tells that on the compact manifold V the sum of the local residues of φ with respect to D_1, \dots, D_k equals zero (the local residue $\text{res}_g(a; F)$ makes sense on manifolds by noting that it depends only on the associated rational differential form $F dX_1 \dots dX_k / g_1 \dots g_k$ and the divisors $(g_1), \dots, (g_k)$). This proves $\text{Res}_g(F) = 0$ in case $\text{deg } F < N$.

The second case is F homogeneous of degree $> N$. We show $F \in (\text{In } g_1, \dots, \text{In } g_k)$. This is an easy generalization of a theorem of Macaulay (cf. e.g. [Ts]) to the weighted situation. Namely, let Q be any homogeneous polynomial and set $P = F \cdot G$. With $Q := \text{In}(g_1) \dots \text{In}(g_k)$ we have

$$\frac{P}{Q} dX_1 \dots dX_k = (\text{deg } P - \text{deg } Q + \sum_i d_i)^{-1} d\sigma = (\text{deg } P - N)^{-1} d\sigma$$

with

$$\sigma = \frac{P}{Q} \sum_{j=1}^k (-1)^{j-1} d_j \cdot X_j dX_1 \dots \widehat{dX_j} \dots dX_k,$$

where $\widehat{}$ means that this entry is to be left out. This is a simple check using the weighted Euler formula $\sum_j d_j X_j \frac{\partial H}{\partial X_j} = \text{deg}(H) \cdot P$, H weighted homogeneous (same proof as usual). Thus

$$\text{Res}_{\text{In } g}(F \cdot G) = \sum_a \int_{\Gamma_a^\varepsilon} d\sigma = 0$$

for all (homogeneous) $G \in \mathbb{C}[X_1, \dots, X_k]$. But this implies $F \in (\text{In } g_1, \dots, \text{In } g_k)$ by the “duality theorem”, cf. [Ts] (this is Poincaré duality in the case of cohomology rings). Modulo (g_1, \dots, g_k) this means that we may reduce F to lower degree. So proceeding by induction N turns out to be “top-degree” in R in that all elements of R can be represented by polynomials of degree $\leq N$.

What remains to be checked is that for F homogeneous of degree N , either $\text{Res}_g(F) \neq 0$ or $F/(g_1, \dots, g_k) \in R_{<N}$. In fact, if $\text{Res}_g(F) = 0$ then $\text{Res}_{\text{In}(g)}(F \cdot G) = 0 \forall G \in \mathbb{C}[X_1, \dots, X_k]$, because for G homogeneous of positive degree this has just been shown. So again $F \in (\text{In } g_1, \dots, \text{In } g_k)$, i.e. modulo (g_1, \dots, g_k) , F may be represented by a polynomial of degree $< N$. \square

REMARK 4.2. The decomposition $R = R_{<N} \oplus \langle J \rangle$ is *not* canonical but rather depends on the particular presentation of R . This may be seen either in elementary terms from the transformation formula for residues or as manifestation of the choice of an isomorphism $\text{Ext}_V^k(\mathcal{O}_Z, \Omega_V^k) \simeq H^0(V, \mathcal{O}_Z)$ in the duality morphism

$$\text{Ext}_V^k(\mathcal{O}_Z, \Omega_V^k) \times H^0(V, \mathcal{O}_Z) \longrightarrow \mathbb{C}$$

induced by the global residue. For quantum cohomology rings and weightings coming from cohomology, however, $R_{<N} = \bigoplus_{d < N} H^d(M, \mathbb{C})$, $N = \dim_{\mathbb{C}} M$ and $\langle Y \rangle = H^{2N}(M, \mathbb{C})$, so the decomposition *has* an invariant meaning in this case. \square

In the quantum cohomology ring the top-degree class $\mathbb{C} \cdot J$ is thus spanned by the class $[\Omega]$ of the volume form. Let $F_{[\Omega]}$ be a polynomial of degree N representing $[\Omega]$ (modulo $(f_i^{[\omega]})$ or modulo (f_i) , this will yield the same result), and

set $c = 1/\text{Res}_{f^{[\omega]}}(F_{[\Omega]})$. By the interpretation of $\langle F \rangle_0$ as coefficient of $[\Omega]$ of $F(\hat{X}_1, \dots, \hat{X}_k) \in QH_{[\omega]}^*(M)$ we conclude (for $H^*(M, \mathbb{C})$ a complete intersection)

$$\langle F \rangle_0 = c \cdot \text{Res}_{f^{[\omega]}}(F) = c \cdot \text{tr}_{f^{[\omega]}} \left(\frac{F}{J} \right) (0).$$

To incorporate the higher genus case we prove

LEMMA 4.3. *For notations as in Proposition 4.1 and $F \in \mathbb{C}[X_1, \dots, X_k]$ let B_F be the bilinear form $R \times R \rightarrow \mathbb{C}$, $(\alpha, \beta) \mapsto \text{Res}_g(F \cdot G_\alpha \cdot G_\beta)$ (with $G_\alpha, G_\beta \in \mathbb{C}[X_1, \dots, X_k]$ representing α, β — this is well-defined), $\eta = B_1$ the “intersection form”. Then*

$$\text{tr}_\eta B_F = \text{Res}_g(F \cdot J).$$

Proof. We give a basis-free, geometric proof. By definition $\text{tr}_\eta B_F$ is the trace of the endomorphism $\mu_F : R \rightarrow R$ of multiplication by (the class of) F . Putting $Z = \text{Spec } R$ we have $R = \bigoplus_{z \in g^{-1}(0)} \mathcal{O}_{Z,z}$. For $z \in g^{-1}(0)$, $F - F(z)$ is nilpotent in $\mathcal{O}_{Z,z}$ (for it has value 0 in z), so $\mu_{F-F(z)}|_{\mathcal{O}_{Z,z}}$ has trace 0 and

$$\text{tr}(\mu_F|_{\mathcal{O}_{Z,z}}) = \text{tr}(\mu_{F-F(z)}|_{\mathcal{O}_{Z,z}}) + \text{tr}(\mu_{F(z)}|_{\mathcal{O}_{Z,z}}) = F(z) \cdot \dim_{\mathbb{C}} \mathcal{O}_{Z,z}.$$

Furthermore, $\dim_{\mathbb{C}} \mathcal{O}_{Z,z} = \text{deg}_z(g)$, the local mapping degree of g at z , so summing up we get

$$\text{tr}_\eta B_F = \sum_{z \in g^{-1}(0)} \text{deg}_z(g) \cdot F(z) = \text{tr}_g(F),$$

which is nothing but $\text{Res}_g(F \cdot J)$ as claimed. \square

In view of the reduction formula to lower genus stated above we conclude a mathematical version of Witten’s “handle gluing formula” in topological QFT [Wi2].

PROPOSITION 4.4. *Let $H^*(M, \mathbb{Q}) = \mathbb{Q}[X_1, \dots, X_k]/(f_1, \dots, f_k)$ (the commutative complete intersection case) and $f_i^{[\omega]}$ the induced relations in $QH_{[\omega]}^*(M)$ as in Proposition 2.2. Put $J = \det \left(\frac{\partial f_i^{[\omega]}}{\partial X_j} \right)$. Then*

$$\langle F \rangle_g = \langle J \cdot F \rangle_{g-1}$$

holds for all $F \in \mathbb{C}[X_1, \dots, X_k]$. \square

In other words, multiplication by J acts as “attaching a handle” to our Riemann surface.

We summarize our considerations in the following main result of the present chapter:

THEOREM 4.5. *Assumptions as in the preceding proposition then for all $F \in \mathbb{C}[X_1, \dots, X_k]$ the following holds*

$$\langle F \rangle_g = c \cdot \text{Res}_{f^{[\omega]}}(J^g \cdot F) = c \cdot \lim_{\substack{y \rightarrow 0 \\ y \text{ regular} \\ \text{value of } f^{[\omega]}}} \text{tr}_{f^{[\omega]}}(J^{g-1} \cdot F)(y),$$

with $c = 1/\text{Res}_{f^{[\omega]}}(F_{[\Omega]})$, $F_{[\Omega]}$ a polynomial representing the class $[\Omega] \in QH_{[\omega]}^*(M)$ of the normalized volume form. \square

We emphasize that for $g > 0$ or $0 \in \mathbb{C}^k$ a regular value of $f^{[\omega]}$ the right-hand side has the form $\sum_{\nu} a_{\nu} F(y_{\nu})$ with $g^{-1}(0) = \{y_{\nu}\}$ and constants a_{ν} independent of F !

Note also that 0 is a regular value of $f^{[\omega]}$ iff $\dim_{\mathbb{C}} \mathcal{O}_{Z,z} = 1$ for all $z \in (f^{[\omega]})^{-1}(0)$, $Z = \text{Spec } \mathbb{C}[X_1, \dots, X_k]/(f_1^{[\omega]}, \dots, f_k^{[\omega]})$, which is if and only if

$$\dim_{\mathbb{C}} H^*(M, \mathbb{C}) = \# \left(f^{[\omega]} \right)^{-1} (0)$$

(“ \geq ” always).

We claim that this is the case for $G(k, n)$: We have a basis of Schubert classes $\{\lambda_1, \dots, \lambda_k\}$ parametrized by sequences $n - k \geq \lambda_1 \geq \dots \geq \lambda_k \geq 0$. The latter are in 1-1 correspondence with subsets $\{\lambda_k + 1, \lambda_{k-1} + 2, \dots, \lambda_1 + k\}$ of $\{1, \dots, n\}$, so $\dim_{\mathbb{C}} H^*(G(k, n), \mathbb{C}) = \binom{n}{k}$.

To find $\binom{n}{k}$ distinct elements in $(f^{[\omega]})^{-1}(0)$, we use a description of the cohomology ring coming from the study of the corresponding Landau-Ginzburg model in physics [Va1]: There is a polynomial W in the X_i (“Landau-Ginzburg potential”) so that in the notations of Theorem 0.1

$$Y_{n+1-i}(X_1, \dots, X_k) = \frac{\partial W}{\partial X_i},$$

and then $f_i^{[\omega]} = \frac{\partial W^{[\omega]}}{\partial X_i}$ with $W^{[\omega]} := W + (-1)^k e^{-[\omega](L)} \cdot X_1$. W has a simple description in terms of Chern roots, i.e. after composition with the $k!$ -fold branched covering $\Sigma : \mathbb{C}^k \rightarrow \mathbb{C}^k$

$$\Sigma(\lambda) = (-\sigma_1(\lambda), \sigma_2(\lambda), \dots, (-1)^k \sigma_k(\lambda)),$$

σ_i the elementary symmetric polynomials:

$$W \circ \Sigma = -\frac{1}{n+1} \sum_i \lambda_i^{n+1}.$$

Thus

$$W^{[\omega]} \circ \Sigma = -\sum_{i=1}^k \left(\frac{\lambda_i^{n+1}}{n+1} - (-1)^k e^{-[\omega](L)} \cdot \lambda_i \right).$$

This all is an easy formal consequence of the algebraic relations between Chern and Segre classes on one side and Chern roots on the other, cf. [BDW] for a mathematical account.

Now in a *non-degenerate* value $\Sigma(\lambda)$ of Σ , $W^{[\omega]}$ is degenerate iff

$$\frac{(\partial W^{[\omega]} \circ \Sigma)}{\partial \lambda_i}(\lambda) = 0$$

for all i . Non-degeneracy of Σ in λ is equivalent to $\lambda_i \neq \lambda_j$ for $i \neq j$. Solving the equation means

$$\lambda_i^n = (-1)^k e^{-[\omega](L)}, \quad i = 1, \dots, k.$$

This system of equations has exactly $n \cdot (n - 1) \cdots (n - k + 1)$ solutions with pairwise distinct λ_i . The fiber of Σ over the regular value $\Sigma(\underline{\lambda})$ consists of the $k!$ permutations of $\{\lambda_1, \dots, \lambda_k\}$, so we get precisely $\binom{n}{k}$ distinct elements of $(f^{[\omega]})^{-1}(0)$, as wanted. Applying the last theorem we obtain

THEOREM 4.6. (Formula of Vafa and Intriligator) *(Notations as in Theorem 0.1) For any $[\omega] \in H^{1,1}(G(k, n))$ the set $C^{[\omega]}$ of critical points of $W^{[\omega]}$ is finite and all of these are non-degenerate. Moreover, for any $F \in \mathbb{C}[X_1, \dots, X_k]$ the following formula for the genus g GW-invariants of $G(k, n)$ holds:*

$$\langle F \rangle_g^{[\omega]} = (-1)^{n+k(k-1)/2} \sum_{\underline{x} \in C^{[\omega]}} \det \left(\frac{\partial^2 W^{[\omega]}}{\partial X_i \partial X_j} \right) (\underline{x}) \cdot F(\underline{x}). \quad \square$$

Proof. What is left is a check of the normalization. This amounts to calculate the residue of $[\Omega] = (-1)^{n-k} X_k^{n-k}$ with respect to $f^{[\omega]}$. We set $a = (-1)^k e^{[\omega](L)}$. Using standard properties of residues one gets

$$\text{Res}_{f^{[\omega]}} \left((-1)^{n-k} X_k^{n-k} \right) = \frac{(-1)^n}{k!} \text{Res}_{\underline{\lambda}^n - \underline{a}} \left(\prod_{i=1}^k \lambda_i^{n-k} \prod_{i < j} (\lambda_i - \lambda_j)^2 \right),$$

where

$$\underline{\lambda}^n - \underline{a} = (\lambda_1^n - a, \dots, \lambda_k^n - a) = -\frac{\partial(W^{[\omega]} \circ \Sigma)}{\partial \lambda_i} = -f^{[\omega]} \circ \frac{\partial \Sigma}{\partial \lambda_i}.$$

In fact, $X_k^{n-k} \circ \Sigma(\underline{\lambda}) = (\prod \lambda_i)^{n-k}, \prod_{i < j} (\lambda_i - \lambda_j)^2$ is the squared Jacobian of Σ and $k!$ is the degree of Σ . Since terms of degree less than $k(n - 1)$ modulo $(\lambda_i^n - a)$ have vanishing residue (Proposition 4.1) only the term $\prod_{i=1}^k \lambda_i^{k-1}$ from the expansion of $\prod_{i < j} (\lambda_i - \lambda_j)^2$ contributes. The coefficient of this term is $(-1)^{k(k-1)/2} \cdot k!$ as one sees by writing $\prod_{i < j} (\lambda_i - \lambda_j)^2$ as determinant of a product of a Vandermonde-matrix with its transposed. What is left is a multiple of the Jacobian $n^k \prod \lambda_i^{n-1}$ of $\underline{\lambda}^n - \underline{a}$, the residue of which is known to be the degree n^k of $\underline{\lambda}^n - \underline{a}$. Putting everything together we get the claimed normalization. \square

REFERENCES

[AsSa] A. ASTASHKEVICH AND V. SADOV: *Quantum cohomology of partial flag manifolds* $F_{n_1 \dots n_k}$, *Comm. Math. Phys.* **170** (1995) 503–528.
 [BeMa] K. BEHREND, Y. MANIN: *Stacks of stable maps and Gromov-Witten invariants*, *Duke Math. J.* **85** (1996) 1–60.
 [Be1] A. BERTRAM: *Towards a Schubert calculus for maps from a Riemann surface to a Grassmannian*, *Internat. J. Math.* **5** (1994) 811–825.
 [Be2] A. BERTRAM: *Quantum Schubert calculus*, preprint **alg-geom 9410024**, to appear in: *Adv. Math.*
 [BDW] A. BERTRAM, G. DASKALOPOULOS AND R. WENTWORTH: *Gromov invariants for holomorphic maps from Riemann surfaces to Grassmannians*, *J. Amer. Math. Soc.* **9** (1996) 529–571.
 [CaDiSt] E. CATTANI, A. DICKENSTEIN, B. STURMFELS: *Computing multidimensional residues*, in: *Proceedings of MEGA 1994*.
 [Fi] G. FISCHER: *Complex analytic geometry*, LNM 538, Springer 1976.
 [Fu] W. FULTON: *Intersection theory*, Springer 1984.
 [GiKi] A. GIVENTAL AND B. KIM: *Quantum cohomology of flag manifolds and Toda lattices*, *Comm. Math. Phys.* **168** (1995) 609–641.

- [Gr] M. GROMOV: *Pseudo holomorphic curves in symplectic manifolds*, Inv. Math. **82** (1985) 307–347.
- [GrHa] P. GRIFFITHS AND J. HARRIS: *Principles of algebraic geometry*, Wiley 1978.
- [Ha] R. HARTSHORNE: *Residues and duality*, LNM 20, Springer 1966.
- [In] K. INTRILIGATOR: *Fusion Residues*, Mod. Phys. Lett. A **38** (1991) 3543–3556.
- [Ki] B. KIM: *Quantum cohomology of flag manifolds G/B and quantum Toda lattices*, preprint `alg-geom 9607001`.
- [LiTi] J. LI, G. TIAN: *Quantum cohomology of homogeneous manifolds*, J. Alg. Geom. **6** (1997) 269–307.
- [Pi] S. PIUNIKHIN: *Quantum and Floer cohomology have the same ring structure*, preprint `hep-th 9401130`.
- [RRW] M.S. RAFI, J. ROSENTHAL AND X. WANG: *Degree of the generalized Plücker embedding of a quot scheme and quantum cohomology*, preprint `hep-th 9402011`.
- [Ru] Y. RUAN: *Topological sigma model and Donaldson type invariants in Gromov theory*, Duke Math. Journ. **83** (1996) 461–500.
- [RuTi] Y. RUAN, G. TIAN: *A mathematical theory of quantum cohomology*, Journ. Diff. Geom. **42** (1995) 259–367.
- [Ts] A.K. TSIKH: *Multidimensional residues and their applications*, Transl. of Math. Monographs **103**, AMS 1992.
- [Va1] C. VAFA: *Topological Landau-Ginzburg models*, Mod. Phys. Lett. A **6** (1991) 337–346.
- [Va2] C. VAFA: *Topological Mirrors and Quantum Rings*, in: Essays on Mirror Manifolds (ed. S.-T. Yau), International Press 1992, 96–119.
- [Wi1] E. WITTEN: *Topological sigma models*, Comm. Math. Phys. **118** (1988), 353–386.
- [Wi2] E. WITTEN: *On the structure of the topological phase of two-dimensional gravity*, Nucl. Phys. B **340** (1990) 281–332.
- [Wi3] E. WITTEN: *The Verlinde algebra and the cohomology of the Grassmannian*, preprint `hep-th 9312104`.