

## THE SYMPLECTIC FLOER HOMOLOGY OF THE FIGURE EIGHT KNOT\*

WEIPING LI†

**Abstract.** In this paper, we compute the symplectic Floer homology of the figure eight knot. This provides first nontrivial knot with trivial symplectic Floer homology.

**1. Introduction.** In [4], we generalized the Casson-Lin invariant [6] to the symplectic theory point of view. Our symplectic Floer homology of knots serves a new invariant for knots, and its Euler characteristic is half of the signature of knots.

We showed that the symplectic Floer homology of the unknotted knot is trivial in [4]. The natural question arises as whether there is a nontrivial knot with trivial symplectic Floer homology. We answer this question in this paper by computing the symplectic Floer homology of the figure eight knot.

Although we know that the signature of the figure eight knot  $4_1 = \overline{\sigma_1\sigma_2^{-1}\sigma_1\sigma_2^{-1}}$  is zero, the signature does not suffice to give the information of our finer invariant - the symplectic Floer homology. For the square knot, we computed in [5] that the symplectic Floer homology is nontrivial even though its signature is zero. Our main result is the following.

**THEOREM.** *The symplectic Floer homology of the figure eight knot  $4_1 = \overline{\sigma_1\sigma_2^{-1}\sigma_1\sigma_2^{-1}}$  is*

$$HF_i^{sym}(4_1) = CF_i^{sym}(4_1) = 0, \quad \text{for all } i \in \mathbf{Z}_{2N}.$$

To our knowledge, this is the first trivial symplectic Floer homology for the nontrivial knot. It is still an open question if there is a non-homotopy 3-sphere with trivial instanton Floer homology. We wish to build the relation between our symplectic Floer homology of knots [4] and the instanton Floer homology of homology 3-spheres [1] through the Dehn surgery technique. Using the calculation of the figure eight knot, we hope to unfold the question about vanishing instanton Floer homology for a nontrivial homology 3-sphere.

### 2. The symplectic Floer homology.

**2.1. The symplectic Floer homology of braids.** We briefly recall our definition of the Floer homology of braids in this subsection. See [4] for more details.

For any knot  $K = \overline{\beta}$  with  $\beta \in B_n$ , the braid group, the space  $\mathcal{R}(S^2 \setminus K)^{[i]}$  can be identified with the space of  $2n$  matrices  $X_1, \dots, X_n, Y_1, \dots, Y_n$  in  $SU(2)$  satisfying

$$(2.1) \quad \text{tr}(X_i) = \text{tr}(Y_i) = 0, \quad \text{for } i = 1, \dots, n,$$

$$(2.2) \quad X_1 \cdot X_2 \cdots X_n = Y_1 \cdot Y_2 \cdots Y_n.$$

\*Received February 22, 1999; accepted for publication June 24, 1999.

†Department of Mathematics, Oklahoma State University, Stillwater, Oklahoma 74078-0613, U.S.A. (wli@math.okstate.edu).

Note that  $\pi_1(S^2 \setminus K)$  is generated by  $m_{x_i}, m_{y_i} (i = 1, 2, \dots, n)$  with one relation  $\prod_{i=1}^n m_{x_i} = \prod_{i=1}^n m_{y_i}$ . There is a unique reducible conjugacy class of representations  $s_K : \pi_1(S^3 \setminus K) \rightarrow U(1)$  such that

$$s_K([m_{x_i}]) = s_K([m_{y_i}]) = \begin{bmatrix} i & 0 \\ 0 & -i \end{bmatrix}.$$

Let  $\mathcal{R}^*(S^2 \setminus K)^{[i]}$  be the subset of  $\mathcal{R}(S^2 \setminus K)^{[i]}$  consisting of irreducible representations. Then  $\mathcal{R}^*(S^2 \setminus K)^{[i]}$  is a monotone symplectic manifold of dimension  $4n - 6$  by Lemma 2.3 in [4]. The symplectic manifold  $(M, \omega)$  is called *monotone* if  $\pi_2(M) = 0$  or if there exists a nonnegative  $\alpha \geq 0$  such that  $I_\omega = \alpha I_{c_1}$  on  $\pi_2(M)$ , where  $I_\omega(u) = \int_{S^2} u^*(\omega) \in \mathbf{R}$  and  $I_{c_1}(u) = \int_{S^2} u^*(c_1) \in \mathbf{Z}$  for  $u \in \pi_2(M)$ . The braid  $\beta$  induces a diffeomorphism  $\phi_\beta : \mathcal{R}^*(S^2 \setminus K)^{[i]} \rightarrow \mathcal{R}^*(S^2 \setminus K)^{[i]}$ . The induced diffeomorphism  $\phi_\beta$  is symplectic, and the fixed point set of  $\phi_\beta$  is  $\mathcal{R}^*(S^3 \setminus K)^{[i]}$  (see Lemma 2.4 in [4]).

Let  $H : \mathcal{R}^*(S^2 \setminus K)^{[i]} \times \mathbf{R} \rightarrow \mathbf{R}$  be a  $C^\infty$  time-dependent Hamiltonian function with  $H(x, s) = H(\phi_\beta(x), s + 1)$ . Let  $X_s$  be the corresponding vector field from  $\omega(X_s, \cdot) = dH_s(\cdot, s)$ , and  $\psi_s$  be the corresponding flow

$$\frac{d\psi_s}{ds} = X_s \circ \psi_s, \quad \psi_0 = id.$$

Then we have  $\psi_{s+1} \circ \phi_\beta^H = \phi_\beta \circ \psi_s$ , where  $\phi_\beta^H = \psi_1^{-1} \circ \phi_\beta$ . Let  $\Omega_{\phi_\beta}$  be the space of smooth paths  $\alpha$  in  $\mathcal{R}^*(S^2 \setminus K)^{[i]}$  such that  $\alpha(s + 1) = \phi_\beta(\alpha(s))$ . The symplectic action  $a_H : \Omega_{\phi_\beta} \rightarrow \mathbf{R}/2\alpha N\mathbf{Z}$  is given by

$$da_H(\gamma)\xi = \int_0^1 \omega(\dot{\gamma} - X_s(\gamma), \xi) ds.$$

So the critical points of  $a_H$  are the fixed points of  $\phi_\beta$ . For  $x \in \text{Fix}(\phi_\beta)$ , define  $\mu(x) = \mu_u(x, s) \pmod{2N}$ , where  $\mu_u$  is the Maslov index and  $N = N(K)$  is the minimal value of the first Chern number of the tangent bundle of  $\mathcal{R}^*(S^2 \setminus K)^{[i]}$ . The integer  $N(K)$  is a knot invariant.

Thus we have a  $\mathbf{Z}_{2N}$ -graded symplectic Floer chain complex:

$$CF_i^{\text{sym}} = \{x \in \text{Fix}(\phi_\beta) \cap \mathcal{R}^*(S^2 \setminus K)^{[i]} : \mu(x) = i\}, \quad i \in \mathbf{Z}_{2N}.$$

The following is Proposition 4.1 and Theorem 4.2 of [4].

**THEOREM 2.1.** *For a knot  $K = \bar{\beta}$  with the property that  $\pi_2(\mathcal{R}^*(S^2 \setminus K)^{[i]}) = 0$  or  $\alpha N(K) = 0$ , there is a well-defined  $\mathbf{Z}$ -graded symplectic Floer homology  $HF_*^{\text{sym}}(\phi_\beta)$ . The symplectic Floer homology  $\{HF_i^{\text{sym}}(\phi_\beta)\}_{i \in \mathbf{Z}_{2N}}$  is a knot invariant and its Euler number is half of the signature of the knot (see [4]).*

**2.2. The symplectic Floer homology of the figure eight knots.** The figure eight knot  $4_1$  has the braid representative  $\sigma_1 \sigma_2^{-1} \sigma_1 \sigma_2^{-1}$ . The knot  $4_1$  has signature zero since  $4_1$  is equivalent (by an orientation preserving homeomorphism) to its mirror image  $\overline{4_1}$ . So the figure eight knot is amphicheral. Also it is well-known that the figure eight knot is not a slice knot, and represents an element of order 2 in the knot cobordism group (see [7]).

We calculate the symplectic Floer homology of the figure eight knot by identifying the fixed points of the induced symplectic diffeomorphism in §2.1. See [3] for the  $SU(2)$  representations of the  $4_1$  knot group without trace zero condition along meridians.

Let  $\mathcal{R}^*(S^2 \setminus 4_1)^{[i]}$  be the subset of  $\mathcal{R}(S^2 \setminus 4_1)^{[i]}$  consisting of irreducible representations. Then  $\mathcal{R}^*(S^2 \setminus 4_1)^{[i]}$  can be also identified with  $(H_3 \setminus S_3)/SU(2)$  in Lin's notation [6], i.e., the set of 6-tuple  $(X_1, X_2, X_3, Y_1, Y_2, Y_3) \in SU(2)^6$  satisfying  $\text{tr}(X_j) = \text{tr}(Y_j) = 0 (j = 1, 2, 3)$  and

$$X_1 X_2 X_3 = Y_1 Y_2 Y_3.$$

By operating the conjugation on  $X_3$  and  $Y_3$ , we may assume that

$$X_3 = \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}, \quad Y_3 = \begin{pmatrix} i \cos \theta & \sin \theta \\ -\sin \theta & -i \cos \theta \end{pmatrix}, \quad 0 \leq \theta \leq \pi.$$

If  $\theta = 0$  and  $\pi$ , then we get two copies of  $(H_2 \setminus S_2)/SU(2)$  which is the pillow case (a 2-sphere with four cone points deleted [4, 6]). For  $0 < \theta < \pi$ , the identification reduces down to the following

$$X_1 X_2 \begin{pmatrix} \cos \theta & -i \sin \theta \\ -i \sin \theta & \cos \theta \end{pmatrix} = Y_1 Y_2.$$

Let  $R_\theta$  be the representations in  $\mathcal{R}^*(S^2 \setminus 4_1)^{[i]}$  satisfying the above equation. So the space  $R_\theta$  is the non-singular piece in  $\mathcal{R}^*(S^2 \setminus K)^{[i]}$ . For  $0 < \theta, \theta' < \pi$ , the space  $R_\theta$  is diffeomorphic to the space  $R_{\theta'}$ . In particular, they are all diffeomorphic to  $R_{\pi/2}$ . So  $\mathcal{R}^*(S^2 \setminus 4_1)^{[i]}$  is a generalized pillow case:

$$\mathcal{R}^*(S^2 \setminus 4_1)^{[i]} = \bigcup_{0 \leq \theta \leq \pi} R_\theta.$$

It is not easy to determine the  $\pi_2(\mathcal{R}^*(S^2 \setminus 4_1)^{[i]})$  and  $N(4_1)$ .

The fixed point set of  $\phi_{4_1}$  is  $\mathcal{R}^*(S^3 \setminus 4_1)^{[i]}$  by Lemma 2.4 in [4]. So we have, for  $\sigma = \sigma_1 \sigma_2^{-1} \sigma_1 \sigma_2^{-1}$ ,  $\text{Fix}(\phi_{4_1}) = \{(X_1, X_2, X_3) \in SU(2)^3 | \sigma(X_j) = X_j, j = 1, 2, 3\}$  up to conjugation. Let  $B_n$  be the braid group of rank  $n$  with the standard generators  $\sigma_1, \dots, \sigma_{n-1}$ , and  $F_n$  be the free group of rank  $n$  generated by  $x_1, \dots, x_n$ . Then the automorphism of  $F_n$  representing  $\sigma_k$  is given by (still denote it by  $\sigma_k$ )

$$(2.3) \quad \begin{aligned} \sigma_k : \quad & x_k \mapsto x_k x_{k+1} x_k^{-1} \\ & x_{k+1} \mapsto x_k \\ & x_l \mapsto x_l, \quad l \neq k, k+1. \end{aligned}$$

By (2.3), we compute the followings.

$$\begin{aligned}
\sigma_1\sigma_2^{-1}\sigma_1\sigma_2^{-1}(x_1) &= \sigma_1\sigma_2^{-1}\sigma_1(x_1) = \sigma_1\sigma_2^{-1}(x_1x_2x_1^{-1}) \\
&= \sigma_1(x_1x_2x_3^{-1}x_2^{-1}x_1^{-1}) \\
&= (x_1x_2x_1^{-1})x_1x_3^{-1}x_1^{-1}(x_1x_2x_1^{-1})^{-1} \\
&= x_1x_2x_3^{-1}x_2^{-1}x_1^{-1}. \\
\sigma_1\sigma_2^{-1}\sigma_1\sigma_2^{-1}(x_2) &= \sigma_1\sigma_2^{-1}\sigma_1(x_2x_3^{-1}x_2^{-1}) \\
&= \sigma_1\sigma_2^{-1}(x_1x_3^{-1}x_1^{-1}) = \sigma_1(x_1^{-1}x_2x_1) \\
&= x_1x_2^{-1}x_1x_2x_1^{-1}. \\
\sigma_1\sigma_2^{-1}\sigma_1\sigma_2^{-1}(x_3) &= \sigma_1\sigma_2^{-1}\sigma_1(x_2^{-1}) = \sigma_1\sigma_2^{-1}(x_1^{-1}) \\
&= \sigma_1(x_1) = x_1x_2x_1^{-1}.
\end{aligned}$$

Therefore the fixed point set of  $\phi_{4_1}$  is the set of points  $(X_1, X_2, X_3) \in SU(2)^3$  such that

$$\begin{aligned}
\text{tr}(X_j) &= 0, \quad j = 1, 2, 3, \\
X_1X_2X_3^{-1}X_2^{-1}X_1^{-1} &= X_1, \\
X_1X_2^{-1}X_1X_2X_1^{-1} &= X_2, \\
X_1X_2X_1^{-1} &= X_3,
\end{aligned}$$

up to conjugation. Up to conjugation, we can assume that

$$X_1 = \begin{pmatrix} i \cos \theta & \sin \theta \\ -\sin \theta & -i \cos \theta \end{pmatrix}, \quad X_2 = \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}, \quad 0 \leq \theta \leq \pi.$$

From the last equation in the above, we obtain

$$\begin{aligned}
X_1X_2X_1^{-1} &= \begin{pmatrix} i \cos \theta & \sin \theta \\ -\sin \theta & -i \cos \theta \end{pmatrix} \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix} \begin{pmatrix} -i \cos \theta & -\sin \theta \\ \sin \theta & i \cos \theta \end{pmatrix} \\
&= \begin{pmatrix} i \cos 2\theta & \sin 2\theta \\ -\sin 2\theta & -i \cos 2\theta \end{pmatrix} = X_3.
\end{aligned}$$

So the matrix  $X_3$  is completely determined by the parameter  $\theta \in [0, \pi]$ . This is, in fact, a key to complete the calculation. Now substituting  $X_3$  into the relation  $\sigma_1\sigma_2^{-1}\sigma_1\sigma_2^{-1}(X_1) = X_1$ , we have

$$\begin{aligned}
X_1X_2X_3^{-1}X_2^{-1}X_1^{-1} &= \begin{pmatrix} -\cos \theta & -i \sin \theta \\ -i \sin \theta & -\cos \theta \end{pmatrix} \begin{pmatrix} -i \cos 2\theta & -\sin 2\theta \\ \sin 2\theta & i \cos 2\theta \end{pmatrix} \begin{pmatrix} -\cos \theta & i \sin \theta \\ i \sin \theta & -\cos \theta \end{pmatrix} \\
&= \begin{pmatrix} -i \cos 4\theta & -\sin 4\theta \\ \sin 4\theta & i \cos 4\theta \end{pmatrix} = X_1.
\end{aligned}$$

This reduces to the equations

$$(2.4) \quad \cos 4\theta = -\cos \theta, \quad \sin 4\theta = -\sin \theta.$$

Similarly, we compute

$$\sigma_1\sigma_2^{-1}\sigma_1\sigma_2^{-1}(X_2) = \begin{pmatrix} i \cos 3\theta & \sin 3\theta \\ -\sin 3\theta & -i \cos 3\theta \end{pmatrix} = X_2 = \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix},$$

to get the equations

$$(2.5) \quad \cos 3\theta = 1, \quad \sin 3\theta = 0.$$

Thus the fixed point of  $\phi_{4_1}$  can be identified with

$$X_1 = \begin{pmatrix} i \cos \theta & \sin \theta \\ -\sin \theta & -i \cos \theta \end{pmatrix}, \quad X_2 = \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}, \quad X_3 = \begin{pmatrix} i \cos 2\theta & \sin 2\theta \\ -\sin 2\theta & -i \cos 2\theta \end{pmatrix}, \quad 0 \leq \theta \leq \pi,$$

subject to equations (2.4) and (2.5). Using the equations (2.5) and the angle addition formulae for sine and cosine functions with  $4\theta = 3\theta + \theta$ , (2.4) becomes

$$(2.6) \quad \sin \theta = 0, \quad \cos \theta = 0.$$

There is no solution for (2.6). Hence

$$(2.7) \quad \text{Fix}(\phi_{4_1}) = \emptyset \quad (\text{empty set}).$$

**THEOREM 2.2.** *The symplectic Floer homology of the figure eight knot  $4_1 = \sigma_1 \sigma_2^{-1} \sigma_1 \sigma_2^{-1}$  is*

$$HF_i^{\text{sym}}(4_1) = CF_i^{\text{sym}}(4_1) = 0, \quad \text{for all } i \in \mathbf{Z}_{2N}.$$

Proof: Since the  $\mathbf{Z}_{2N}$ -graded symplectic Floer chain complex  $CF_i^{\text{sym}}(4_1)$  is generated by  $\text{Fix}(\phi_{4_1})$ , the result follows from (2.7).  $\square$

**2.3. The symplectic Floer homology of knots with braid representatives in  $B_3$ .** It seems that the method in §2.2 can be adapted to knots with braid representatives in  $B_3$ . We are going to illustrate another example to show that the computation for the figure eight knot in §2.2 is quite lucky.

Let  $K = 5_2$  be the knot with 5-crossings. We have the braid representative  $\sigma_1^2 \sigma_2^2 \sigma_1^{-1} \sigma_2$  for the knot  $5_2$  (see [7]). Thus the fixed points of  $\phi_{5_2}$  can be identified, by the same method in §2.2, with the set of points  $(X_1, X_2, X_3) \in SU(2)^3$  such that

$$\begin{aligned} \text{tr}(X_j) &= 0, \quad j = 1, 2, 3, \\ X_1 X_2 X_3 X_1 X_2^{-1} X_1^{-1} X_3^{-1} X_2^{-1} X_1^{-1} &= X_1, \\ X_1 X_2 X_3^{-1} X_1^2 X_2^{-1} X_1^{-1} &= X_2, \\ X_1 X_2 X_1^{-1} X_2^{-1} X_1^{-1} &= X_3, \end{aligned}$$

up to conjugation. This follows a straightforward calculation of  $\sigma_1^2 \sigma_2^2 \sigma_1^{-1} \sigma_2(x_j) (j = 1, 2, 3)$ . Again we can compute  $X_3$  from the last equation in the above.

$$X_1 X_2 X_1^{-1} X_2^{-1} X_1^{-1} = \begin{pmatrix} -i \cos 3\theta & -\sin 3\theta \\ \sin 3\theta & i \cos 3\theta \end{pmatrix} = X_3.$$

Then  $\sigma_1^2 \sigma_2^2 \sigma_1^{-1} \sigma_2(X_j) = X_j (j = 1, 2)$  gives us

$$\begin{aligned} \sigma_1^2 \sigma_2^2 \sigma_1^{-1} \sigma_2(X_1) &= \begin{pmatrix} -i \cos 6\theta & -\sin 6\theta \\ \sin 6\theta & i \cos 6\theta \end{pmatrix} = X_1 \\ \sigma_1^2 \sigma_2^2 \sigma_1^{-1} \sigma_2(X_2) &= \begin{pmatrix} -i \cos 5\theta & -\sin 5\theta \\ \sin 5\theta & i \cos 5\theta \end{pmatrix} = X_2. \end{aligned}$$

Thus we need to solve the equations

$$(2.8) \quad \cos 6\theta = -\cos \theta, \quad \sin 6\theta = -\sin \theta, \quad \cos 5\theta = -1, \quad \sin 5\theta = 0.$$

There are three solutions of (2.8) with  $\theta = \frac{\pi}{5}, \frac{3\pi}{5}, \pi$ . Let  $\rho_j (j = 1, 2, 3)$  be the corresponding fixed points of  $\phi_{5_2}$  in  $\mathcal{R}^*(S^2 \setminus 5_2)^{[2]}$ .

By following the method in [2], for  $K = 5_2$ , we have all type I double points so that the correction term  $\mu = 0$ . Using the definition of Goeritz matrix in §1 of [2], we get the Goeritz matrix of  $5_2$ :

$$G(5_2) = \begin{pmatrix} 4 & -3 & -1 \\ -3 & 4 & -1 \\ -1 & -1 & 2 \end{pmatrix}.$$

By the theorem 6 of [2], we have

$$\text{Signature}(5_2) = \text{Signature}(G(5_2)) - \mu = 2.$$

By Theorem 2.1, the Euler characteristic of the symplectic Floer homology of  $5_2$  is one.

**PROPOSITION 2.3.** *The symplectic Floer chain complex of  $5_2$  is given by: one of the odd chain groups is generated by one of  $\rho_j (j = 1, 2, 3)$ ; even chain groups are generated by the rest two fixed points of  $\phi_{5_2}$ .*

It is nontrivial to determine the Maslov index of  $\rho_j$  and the possible Floer boundary map in order to complete the calculation.

#### REFERENCES

- [1] A. FLOER, *Instanton Homology*, in Geometry of Low-Dimensional Manifolds 1, L. M. S. Lecture Note Series 150, 1989, pp. 115–124.
- [2] C. GORDON AND R. LITHERLAND, *On the signature of a link*, Invent. Math., 47 (1978), pp. 53–69.
- [3] P. KIRK AND E. KLASSEN, *Chern-Simons invariants of 3-manifolds and representation spaces of knot groups*, Math. Ann., 287 (1990), pp. 343–367.
- [4] W. LI, *Casson-Lin's invariant and Floer homology*, J. Knot Theory and its Ramification, 6:6 (1997), pp. 851–877.
- [5] W. LI, *The symplectic Floer homology of the square knot and granny knots*, Acta Math. Sinica (Series B), 15:1 (1999), pp. 1–10.
- [6] X. S. LIN, *A knot invariant via representation spaces*, J. Diff. Geom., 35 (1992), 337–357.
- [7] D. ROLFSEN, *Knots and Links*, Publish or Perish, Inc, 1976 & 1990.