

ON THE AMPLE CONE OF AN AMPLE HYPERSURFACE *

BALÁZS SZENDRÓI†

For a smooth ample hypersurface D of dimension at least 3 in a smooth projective variety X , the restriction map $r : \text{Pic}(X) \rightarrow \text{Pic}(D)$ is an isomorphism by weak Lefschetz. In a recent paper [3], Hassett, Lin and Wang gave an example of a pair $D \subset X$ with $\dim D = 3$ such that the cone of ample divisor classes of D is strictly larger than the image under r of the cone of ample divisor classes of X ; in other words, the weak Lefschetz principle can fail for ample cones.

The purpose of this note is to point out that not only is the ample cone of a hypersurface D often larger than the ample cone of the ambient space X , in general it cannot even be computed from birational models of X . This contrasts with the conjecture of [2] (already disproved for semi-ample hypersurfaces in [5]), which predicted that such a description is possible. Note that in the special case studied in [3], the ample cone is the union of ample cones of birational models of X (Remark 2.3). However, a slight generalization of the example gives hypersurfaces with ample cone larger than the union of ample cones of all birational models of the ambient space (Corollary 3.1).

For a projective variety Y , denote by $\mathcal{N}(Y) \subset \text{Pic}_{\mathbb{R}}(Y)$ the cone of nef divisor classes; this is the closure of the ample cone and also the dual of the Mori cone by Kleiman's criterion. Facets (codimension-one faces) of $\mathcal{N}(Y)$ correspond in favourable cases to primitive (non-factorizable) contraction morphisms on Y .

1. The ambient space. I use the notation of [3]. Let $\varphi : X \rightarrow \mathbb{P}^4$ be the blowup of two distinct points $p_1, p_2 \in \mathbb{P}^4$. The line $p_1 p_2 = l' \subset \mathbb{P}^4$ has proper transform $l \subset X$. The Picard group $\text{Pic}(X) = \langle H, E_1, E_2 \rangle$ is generated by the classes of exceptional divisors E_1, E_2 of φ and the proper transform H of a general hyperplane section of \mathbb{P}^4 through p_1, p_2 . Let \mathcal{B}, \mathcal{C} be the cones in $\text{Pic}_{\mathbb{R}}(X)$ spanned by the rays $H + E_1, H + E_2, H + E_1 + E_2$ and $H, H + E_1, H + E_2$ respectively.

PROPOSITION 1.1.

- (i) *The nef cone of X is $\mathcal{N}(X) = \mathcal{B}$; the three facets of the cone correspond to the contractions of E_i , and to a small contraction $\psi : X \rightarrow \bar{X}$ contracting the single extremal curve $l \cong \mathbb{P}^1$ with $K_X \cdot l = 1$.*
- (ii) *The antiflip $\psi^+ : X^+ \rightarrow \bar{X}$ of ψ exists. The proper transform E_i^+ of E_i is isomorphic to \mathbb{P}^3 blown up in a point; $E_1^+ \cap E_2^+ \cong \mathbb{P}^2$ is the exceptional divisor of both blowups and also the small exceptional locus of ψ^+ .*
- (iii) *Let $\alpha : X^+ \dashrightarrow X$ be the flip and identify the Picard groups of X and X^+ by the map $\alpha_* : \text{Pic}(X^+) \xrightarrow{\sim} \text{Pic}(X)$ induced by proper transform on divisors composed with the isomorphisms between the Picard groups of X, X^+ and their class groups. Then $\mathcal{N}(X^+) = \mathcal{C}$. The three facets of the cone \mathcal{C} correspond to the contraction ψ^+ and two different \mathbb{P}^1 -bundle structures $\pi_i : X^+ \rightarrow Z$. The image Z is the blowup of \mathbb{P}^3 in one point. The fibres l_i of π_i are generically the lines of the original \mathbb{P}^4 through the point p_i . On E_i^+ , π_i is the identity; on E_{3-i}^+ it is the unique projection to \mathbb{P}^2 .*

*Received May 9, 2002; accepted for publication November 15, 2002.

†Mathematics Institute, University of Warwick, Coventry CV4 7AL, United Kingdom and Alfréd Rényi Institute of Mathematics, Hungarian Academy of Sciences (balazs@maths.warwick.ac.uk).

Proof. Once the statement is formulated, the proof becomes an elementary exercise; my personal favourite is toric geometry, which provides a clean and uniform proof. In the lattice $N = \mathbb{Z}^4$, the fan spanned by

$$v_0 = (-1, -1, -1, -1), v_3 = (1, 0, 0, 0), \dots, v_6 = (0, 0, 0, 1)$$

describes \mathbb{P}^4 . Barycentric subdivision of the cones $v_0v_3v_4v_5$ and $v_0v_3v_4v_6$ by the vertices

$$v_1 = (0, 0, 0, -1), v_2 = (0, 0, -1, 0)$$

leads to X . In this language, E_i are the torus-invariant divisors corresponding to the rays v_i and l is the torus orbit corresponding to the facet $v_0v_3v_4$. A short check using toric Mori theory [4] shows that there are three primitive contractions on X : the contractions of E_i and a contraction ψ with one-dimensional exceptional locus l . The extremal curves define three inequalities on ample classes which gives $\mathcal{N}(X) = \mathcal{B}$.

The fan of X^+ is obtained by replacing the cones $v_0v_1v_3v_4$ and $v_0v_2v_3v_4$ by the cones $v_0v_1v_2v_3$, $v_0v_1v_2v_4$ and $v_1v_2v_3v_4$. The new codimension-two face v_1v_2 represents the small exceptional locus $E_1^+ \cap E_2^+ \cong \mathbb{P}^2$. There are three primitive contractions again; the two \mathbb{P}^1 -bundle structures π_i come from projecting the whole fan along the vector v_i for $i = 1, 2$. The tasks of drawing the pictures and filling in the details are left to the reader. \square

2. The hypersurface. Let now $D' \subset \mathbb{P}^4$ be a smooth hypersurface of degree $d > 2$ containing p_1 and p_2 ; assume for simplicity that the line l' is either contained in D' or is at most simply tangent to D' at p_i for $i = 1, 2$ while meeting it transversally elsewhere. Let $D \subset X$ be the proper transform of D' . A simple computation shows that D is ample in X ; hence by the Lefschetz principle, the Picard groups of X and D can be identified by the restriction isomorphism $r : \text{Pic}(X) \rightarrow \text{Pic}(D)$.

PROPOSITION 2.1.

(i) If $l' \subset D'$, then

$$\mathcal{N}(D) = \mathcal{B}.$$

(ii) Assume that D' does not contain l' , but for both indices $i = 1, 2$ one of the following holds:

- a. D' contains a line through p_i , or
- b. the line l' is tangent to D' at p_{3-i} .

Then


$$\mathcal{N}(D) = \mathcal{B} \cup \mathcal{C}.$$

(iii) In all other cases, there is a strict inclusion

$$\mathcal{N}(D) \supset \mathcal{B} \cup \mathcal{C}.$$

Proof. $\mathcal{N}(D) \supset \mathcal{B}$ is clear as a restriction of an ample class from X to D is also ample. Part (i) follows immediately from Proposition 1.1(i), since every face of the nef cone of X corresponds to a contraction with D containing at least one fibre.

To prove (ii), let $\bar{D} \subset \bar{X}$, $D^+ \subset X^+$ be the proper transforms of D in \bar{X} and X^+ . As D meets l in a finite number of points, the map $\psi|_D : D \rightarrow \bar{D}$ is finite and

FIG. 2.1. Possibilities for the shape of $\mathcal{N}(D)$

birational and there is a Cartesian diagram

$$\begin{array}{ccc}
 & \tilde{D} & \\
 D & \swarrow & \searrow & D^+ \\
 & \bar{D} &
 \end{array}$$

By the conditions put on D' , the map $\sigma : \tilde{D} \rightarrow D$ is the blowup of the intersection points of l and D ; the map $\tau : \tilde{D} \rightarrow D^+$ is again finite and birational. Write $\beta : D^+ \dashrightarrow D$ for the composition $\sigma \circ \tau^{-1}$.

Take any ample divisor class A^+ on X^+ . Part (ii) will follow if I can show that the divisor class $r(\alpha_* A^+)$ is ample on D . However, using the fact that τ is finite birational,

$$r(\alpha_* A^+) = \beta_* r^+(A^+) = \sigma_* \tau^* r^+(A^+),$$

where $r^+ : \text{Pic}(Y^+) \rightarrow \text{Pic}(X^+)$ is the restriction map. As A^+ is ample on Y^+ , $r^+(A^+)$ is an ample Cartier divisor class on X^+ and hence $\tau^* r^+(A^+)$ is ample on \tilde{D} . Now the ampleness of $r(\alpha_* A^+)$ on D follows by applying Lemma 2.2 to the map $\sigma : \tilde{D} \rightarrow D$.

To prove (iii), take $A = \pi_i^* B$ where B is ample on Z ; A is a nef but non-ample divisor class in the boundary of the cone $\mathcal{N}(X^+)$. If none of the special circumstances of (i)-(ii) happen, then D^+ does not contain any fibre of π_i , so π_i restricts to D^+ as a finite morphism. Hence the restriction of A to D^+ is ample. The argument of (ii) now shows that $r(\alpha_* A)$ is ample on D and as ampleness is an open condition, (iii) follows. \square

LEMMA 2.2. *Let Y be a smooth threefold, $f : \tilde{Y} \rightarrow Y$ the blowup of a closed point. If A is an ample divisor class on \tilde{Y} , then its proper transform $B = f_* A$ is ample on Y .*

Proof. Let E be the exceptional divisor; then $f^* B = A + \lambda E$ with a non-negative integer λ . If C is a curve in Y with proper transform \tilde{C} in \tilde{Y} , then

$$B \cdot C = f^* B \cdot \tilde{C} = A \cdot \tilde{C} + \lambda E \cdot \tilde{C} > 0$$

as $A \cdot \tilde{C} > 0$ and $E \cdot \tilde{C} \geq 0$ since \tilde{C} is not contained in E . If S is any surface in X with proper transform \tilde{S} then

$$B^2 \cdot S = (A + \lambda E)^2 \tilde{S} = A \cdot (A + \lambda E) \cdot \tilde{S} \geq A^2 \cdot \tilde{S} > 0.$$

The second equality holds since $(A + \lambda E) \cdot E \cdot \tilde{S} = f^*B \cdot \tilde{S}|_E = 0$ by the projection formula, as E and \tilde{S} intersect in a curve contracted by f ; the first inequality uses $A \cdot E \cdot \tilde{S} = A \cdot \tilde{S}|_E \geq 0$. Finally in same way

$$B^3 = (A + \lambda E)^3 \geq (A + \lambda E)^2 \cdot A = (A + \lambda E) \cdot A^2 \geq A^3 > 0.$$

Hence by the Nakai–Moishezon criterion, B is ample. \square

REMARK 2.3. Note that in the special case of smooth cubics studied in [3], the first condition of Proposition 2.1(ii) is always satisfied: a smooth cubic threefold is covered by lines. Hence either $l' \subset D'$ and $\mathcal{N}(D) = \mathcal{B}$, or $l' \not\subset D'$ and $\mathcal{N}(D) = \mathcal{B} \cup \mathcal{C}$. [3, Assumptions 2.5] fixes the type of the primitive contractions on D induced by π_i : if D' contains finitely many lines through p_i and l' is not tangent to D' at p_i , then (the Stein factorization of) $\pi_i|_D$ is a small flopping contraction ($K_D \cdot l_i = 0$); otherwise it is divisorial.

3. Good hypersurfaces. In the original example of [3], as well as in all cases where the conditions of Proposition 2.1(i)-(ii) are satisfied, the ample cone of the hypersurface D has an explicit description in terms of nef cones of models of its ambient space; D is good in the following sense:

Good hypersurface: Call a hypersurface $D \subset X$ *good*, if the nef cone $\mathcal{N}(D)$ is a union of cones $r(\alpha_{i*}\mathcal{N}(X_i))$, where $\alpha_i : X_i \dashrightarrow X$ are certain birational models of X isomorphic to X in codimension one.

Anticanonical hypersurfaces in certain toric varieties were conjectured to be good by Cox and Katz [2, Conjecture 6.2.8]. By [5], this conjecture can fail if D is only semi-ample. Alas, an ample but non-anticanonical $D \subset X$ can also fail to be good.

COROLLARY 3.1. *Take any smooth degree $d > 3$ hypersurface D' in \mathbb{P}^4 with no lines through p_i and with l' not tangent at p_i for both $i = 1, 2$ and let D be its proper transform in X as before. Then $D \subset X$ is not good.*

Proof. By Proposition 2.1(iii), $\mathcal{N}(D)$ is strictly larger than $\mathcal{B} \cup \mathcal{C}$. On the other hand, X has no further birational models which are isomorphic in codimension one. For if $\alpha_i : X_i \dashrightarrow X$ is any such model, then $\alpha_{i*}A$ is a moving divisor class for A ample on X_i . However, from the explicit description of Proposition 1.1 it is clear that the moving cone of X is the interior of $\mathcal{B} \cup \mathcal{C}$. So X_i is isomorphic to X or X^+ . \square

Note that a 3-dimensional ample and anticanonical hypersurface is always good: if $D \in |-K_X|$ is ample, $r(\mathcal{N}(X)) = \mathcal{N}(D)$ by Kollár's result [1]. To the best of my knowledge, the analogous result is not known if $\deg(D) > 3$.

Acknowledgements. I thank Brendan Hassett for bringing [3] to my attention, Massimiliano Mella for pointing out an error in the first version and Miles Reid for help with Lemma 2.2.

REFERENCES

- [1] C. BORCEA, *Homogeneous vector bundles and families of Calabi-Yau threefolds II*, with an Appendix by János Kollár, in: Proc. Sympos. in Pure Math. 52 (ed. E. Bedford et al), Part 2. 83–91, AMS, Providence, RI, 1991.
- [2] D.A. COX AND S. KATZ, *Mirror symmetry and algebraic geometry*, Mathematical Surveys and Monographs 68, American Mathematical Society, Providence, 1999.
- [3] B. HASSETT, H.-W. LIN AND C.-L. WANG, *The weak Lefschetz principle is false for ample cones*, Asian J. Math., 6 (2002), pp. 95–99.
- [4] M. REID, *Decomposition of toric morphisms*, in: Algebra and Geometry, papers dedicated to I.R. Shafarevich on the occasion of his 60th birthday (eds. M. Artin and J. Tate), vol. II. 395–418, Progr. Math. 36, Birkhauser, Boston, Basel, Stuttgart, 1983.
- [5] B. SZENDRŐI, *On a conjecture of Cox and Katz*, Math. Zeitschrift, 240 (2002), pp. 233–241.

