

NEWTON SUM RULES AND MONOTONICITY PROPERTIES OF THE
ZEROS OF SCALED CO-RECURSIVE ASSOCIATED POLYNOMIALS

E. K. Ifantis, C. G. Kokologiannaki, and P. D. Siafarikas

Dedicated to Prof. Lee Lorch for his 80th birthday

ABSTRACT. Let $Q_n(x; \beta, \gamma, c)$ be the polynomials of degree n which satisfy the recurrence relation:

$$\begin{aligned} \alpha_{n+c}Q_{n+1}(x; \beta, \gamma, c) + \alpha_{n+c-1}Q_{n-1}(x; \beta, \gamma, c) + (\beta_{n+c} + \beta\delta_{n,0})Q_n(x; \beta, \gamma, c) \\ = x(1 + (\gamma - 1)\delta_{n,0})Q_n(x; \beta, \gamma, c), \end{aligned}$$

$$Q_{-1}(x; \beta, \gamma, c) = 0, \quad Q_0(x; \beta, \gamma, c) = 1.$$

In the above, β is real, $\gamma > 0$, α_{n+c} and β_{n+c} are real sequences with $\alpha_{n+c} > 0$, and $\delta_{n,0}$ is the Kronecker symbol. These polynomials are called scaled co-recursive associated polynomials. The co-recursive associated orthogonal polynomials are obtained from the above for $\gamma = 1$.

In this paper, the Newton sum rules for the k -th power of the zeros of scaled co-recursive associated orthogonal polynomials are determined in terms of the Newton sum rules of associated orthogonal polynomials. Some monotonicity properties of the zeros also are given.

1. Introduction

The orthogonal polynomials $P_n(x)$, $n \geq 0$, of degree n , with respect to a positive Borel measure on the real line with infinite mass points, can be defined by the recurrence relation:

$$\begin{aligned} \alpha_n P_{n+1}(x) + \alpha_{n-1} P_{n-1}(x) + \beta_n P_n(x) = x P_n(x) \\ P_{-1}(x) = 0, \quad P_0(x) = 1, \end{aligned} \tag{1.1}$$

where $\alpha_n > 0$ and β_n are real sequences.

The associated polynomials $P_n(x; c)$ of the above polynomials are obtained when we replace n by $n + c$ in the coefficients α_n and β_n of (1.1), i.e.,

$$\begin{aligned} \alpha_{n+c} P_{n+1}(x; c) + \alpha_{n+c-1} P_{n-1}(x; c) + \beta_{n+c} P_n(x; c) = x P_n(x; c), \\ P_{-1}(x; c) = 0, \quad P_0(x; c) = 1. \end{aligned} \tag{1.2}$$

The associated polynomials are called, associated of order c if c is an integer and numerator polynomials if $c = 1$. For particular cases of associated polynomials, many authors [3, 4, 17, 18, 37, 38, 43] gave explicitly the corresponding 4th-order differential equations. Very recently [46] for the associated polynomials of arbitrary order c which

Received September 19, 1995, revised July 9, 1996.

1991 *Mathematics Subject Classification*: 33C10.

Key words and phrases: scaled co-recursive associated polynomials, zeros, Newton sum rules, monotonicity.

satisfy (1.2), the corresponding unique differential equation was calculated with the help of a program built in Mathematica symbolic language. Results for particular cases of associated polynomials concerning explicit forms, orthogonality measures, monotonicity properties, and differential inequalities for their zeros were given in [2, 18, 19, 24, 28, 32, 43].

The co-recursive polynomials $Q_n(x)$ of $P_n(x)$, which first were introduced by T. S. Chihara [8], satisfy the same recurrence relation (1.1) with the restriction $Q_1 = \frac{x-\beta_0-\beta}{\alpha_0}$. They can be defined directly by

$$\begin{aligned} \alpha_n Q_{n+1}(x) + \alpha_{n-1} Q_{n-1}(x) + (\beta_n + \beta \delta_{n,0}) Q_n(x) &= x Q_n(x), \\ Q_{-1}(x) = 0, \quad Q_0(x) &= 1 \end{aligned} \tag{1.3}$$

where

$$\delta_{n,0} = \begin{cases} 1 & \text{for } n = 0, \\ 0 & \text{for } n \neq 0. \end{cases}$$

Orthogonal polynomials with perturbed recurrence relations play an important role not only with the Orthogonal Polynomials Theory (see [1, 8, 33, 34, 39]), but also in some other areas such as Quantum Mechanics [42], Birth and Death Processes [27], etc. Recently A. Ronveaux and F. Marcellan [39] gave explicitly the 4th-order differential equation satisfied by the co-recursive polynomials of the classical orthogonal polynomials.

The scaled co-recursive associated polynomials satisfy the recurrence relation

$$\begin{aligned} \alpha_{n+c} Q_{n+1}(x; \beta, \gamma, c) + \alpha_{n+c-1} Q_{n-1}(x; \beta, \gamma, c) \\ + (\beta_{n+c} + \beta \delta_{n,0}) Q_n(x; \beta, \gamma, c) &= x(1 + (\gamma - 1)\delta_{n,0}) Q_n(x; \beta, \gamma, c), \\ Q_{-1}(x; \beta, \gamma, c) = 0, \quad Q_0(x; \beta, \gamma, c) &= 1, \end{aligned} \tag{1.4}$$

where $\gamma > 0$. For $\gamma = 1$ these polynomials are the co-recursive associated orthogonal polynomials, which were introduced by J. Letessier in [29, 30, 31] for special cases. The case $\gamma = 0$ is also of some interest in connection with the results in [13, 14, 21, 41].

Given an orthogonal polynomial family $\{P_n(x)\}_{n=0}^\infty$, one of the most interesting problems in this subject is to study how the distribution of zeros of $P_n(x)$ is deformed when the coefficients of the recurrence relation satisfied by the orthogonal sequence are modified. It is a well-known fact that when standard modifications of the recurrence (associated, co-recursive, scaled co-recursive, co-modified, or any of its combinations) are considered, the asymptotic zero distribution of both the starting and perturbed families coincide. This means that, in this context, the finite n case is the one of interest. In this situation, one of the most natural ways of dealing with this problem is to compute the moments of the zero distribution of $P_n(x)$ or, equivalently, the Newton sum rules of their zeros, i.e., the sum of the k -th power of the zeros, $k \geq 1$, from which valuable information can be derived (see, e.g., [40, 46]).

Some of the well-known methods for determining Newton sum rules of the zeros of orthogonal polynomials [5–7, 12, 36], semiclassical polynomials [45], and associated polynomials of any order [46 and the references therein] are based on the differential equations which are satisfied by these polynomials. In the case of co-recursive associated polynomials, to the best of our knowledge, only for the Laguerre polynomials [29] and for some special cases of Jacobi polynomials [30, 31] are the differential

equations known. Another difficulty is that the known differential equations for the co-recursive associated polynomials are very complicated and difficult to handle. So methods based on the differential equations are not the natural way of dealing with these polynomials. Also some other well-known methods for determining Newton sum rules are based on the three-term recurrence relation, [9, 15, 16], on the explicit expression of the polynomials [35], and on the weight function [10, 11 and the references therein]. For a summary of all these methods, see [44].

In Section 2, we determine the Newton sum rules $\sum_{n=0}^{N-1} \omega_n^k(\beta, \gamma, c)$, $k = 1, 2, \dots$, for the zeros $\omega_n(\beta, \gamma, c)$, $n = 0, 1, 2, \dots, N - 1$, of the scaled co-recursive associated polynomials $Q_n(x; \beta, \gamma, c)$ in terms of the Newton sum rules $\sum_{n=0}^{N-1} \lambda_n^k(c)$, $k = 1, 2, \dots$, of the zeros $\lambda_n(c) \equiv \omega_n(0, 1, c)$ of the corresponding associated polynomials $P_n(x; c)$.

Some examples of Newton sum rules of the classical associated polynomials are given in Section 3.

In Section 4, we give some monotonicity properties of the zeros of the scaled co-recursive associated polynomials.

2. The main result

The method we use is general and is valid also for other cases in addition to the particular case considered in this paper. It has been used successfully by the authors in previous work [20–26]. It is well-known (see for instance [25]) that $\omega_n(\beta, \gamma, c)$, $n = 0, 1, 2, \dots, N - 1$ is a zero of the polynomial $Q_n(x; \beta, \gamma, c)$, defined by (1.4), if and only if it is an eigenvalue of the following generalized eigenvalue problem

$$(T_0 + \beta P_0)x_n = \omega_n(\beta, \gamma, c)(1 + (\gamma - 1)P_0)x_n, x_n \in H_N, \quad n = 0, 1, 2, \dots, N - 1. \quad (2.1)$$

In (2.1), $T_0 = AV^* + VA + B$ where A, B are the diagonal operators $Ae_n = \alpha_{n+c}e_n$, $Be_n = \beta_{n+c}e_n$ where $\alpha_{n+c} > 0$ and β_{n+c} are real sequences in a finite dimensional Hilbert space H_N with the orthonormal basis e_n , $n = 0, 1, 2, \dots, N - 1$ and scalar product (\cdot, \cdot) , V is the truncated shift operator ($Ve_n = e_{n+1}$, $n = 0, 1, \dots, N - 2$, $Ve_{N-1} = 0$), and V^* is its adjoint ($V^*e_n = e_{n-1}$, $n = 1, 2, \dots, N - 1$, $V^*e_0 = 0$). Finally P_0 is the projection operator on the one-dimensional space spanned by e_0 , i.e., $P_0x = (x, e_0)e_0$, $x \in H_N$. The operator C is defined by

$$Ce_n = c_n e_n, \quad (2.2)$$

where $c_0 = \gamma$ and $c_n = 1$, $n = 1, 2, \dots, N - 1$. Since $\gamma > 0$, the operator C is positive-definite, so that both C^{-1} and $C^{-1/2}$ exist.

There is a simple and important result in operator theory which finds here a useful application. This result is known for operators of trace class and Hilbert-Schmidt operators. For a symmetric operator M in a finite-dimensional real Hilbert space, e.g. the space H_N , the result can be stated as follows:

The sum $\sum_{n=0}^{N-1} (Me_n, e_n)$ is independent of the orthonormal basis e_n , $n = 0, 1, 2, \dots, N - 1$, i.e., if e_n and v_n are two orthonormal bases in H_N , then

$$\sum_{n=0}^{N-1} (Me_n, e_n) = \sum_{n=0}^{N-1} (Mv_n, v_n). \quad (2.3)$$

If $x_n, n = 0, 1, 2, \dots, N - 1$ is the complete orthonormal system of the eigenvectors of T_0 in H_N and $\omega_n(0, 1, c) \equiv \lambda_n(c)$ are the corresponding simple eigenvalues, i.e., the zeros of the polynomial $P_n(x; c)$, then for $M = T_0^k$, we find from (2.1) that

$$\sum_{n=0}^{N-1} (T_0^k e_n, e_n) = \sum_{n=0}^{N-1} (T_0^k x_n, x_n) = \sum_{n=0}^{N-1} \lambda_n^k(c). \tag{2.4}$$

From the relation,

$$\begin{aligned} T_0 e_n &= \alpha_{n+c-1} e_{n-1} + \alpha_{n+c} e_{n+1} + \beta_{n+c} e_n, \\ T_0 e_0 &= \alpha_{0+c} e_1 + \beta_{0+c} e_0, \end{aligned} \tag{2.5}$$

we get for $1 \leq k \leq 4$

$$\sum_{n=0}^{N-1} \lambda_n(c) = \sum_{n=0}^{N-1} \beta_{n+c}, \tag{2.6}$$

$$\sum_{n=0}^{N-1} \lambda_n^2(c) = 2 \sum_{n=0}^{N-2} \alpha_{n+c}^2 + \sum_{n=0}^{N-1} \beta_{n+c}^2, \tag{2.7}$$

$$\sum_{n=0}^{N-1} \lambda_n^3(c) = 3 \sum_{n=0}^{N-2} \alpha_{n+c}^2 \beta_{n+c} + 3 \sum_{n=0}^{N-2} \alpha_{n+c}^2 \beta_{n+c+1} + \sum_{n=0}^{N-1} \beta_{n+c}^3, \tag{2.8}$$

$$\begin{aligned} \sum_{n=0}^{N-1} \lambda_n^4(c) &= \sum_{n=0}^{N-1} \beta_{n+c}^4 + 2 \sum_{n=0}^{N-2} \alpha_{n+c}^4 + 4 \sum_{n=0}^{N-3} \alpha_{n+c}^2 \alpha_{n+c+1}^2 + 4 \sum_{n=0}^{N-2} \alpha_{n+c}^2 \beta_{n+c}^2 \\ &\quad + 4 \sum_{n=0}^{N-2} \alpha_{n+c}^2 \beta_{n+c+1}^2 + 4 \sum_{n=0}^{N-2} \alpha_{n+c}^2 \beta_{n+c} \beta_{n+c+1}. \end{aligned} \tag{2.9}$$

The following theorem is useful in the case of scaled co-recursive associated polynomials where the differential equations in general are unknown or very difficult to handle.

Theorem 2.1. *Let $\lambda_n(c)$ and $\omega_n(\beta, \gamma, c)$ be the zeros of the polynomials $P_n(x; c)$ and $Q_n(x; \beta, \gamma, c)$, respectively, and let k be a positive integer. Then*

$$\sum_{n=0}^{N-1} \omega_n^k(\beta, \gamma, c) = \sum_{n=0}^{N-1} \lambda_n^k(c) + (T^k e_0, e_0) + (T^k e_1, e_1) - (T_0^k e_0, e_0) - (T_0^k e_1, e_1) \tag{2.10}$$

where

$$T = C^{-1/2}(T_0 + \beta P_0)C^{-1/2} \tag{2.11}$$

and

$$T_0 = AV^* + VA + B.$$

Proof. By setting $x_n = C^{-1/2}y_n$ where $C = 1 + (\gamma - 1)P_0$, the problem (2.1) is transformed to

$$C^{-1/2}(T_0 + \beta P_0)C^{-1/2}y_n = \omega_n(\beta, \gamma, c)y_n, \quad \|y_n\| = 1.$$

From the above, we find

$$Ty_n = \omega_n(\beta, \gamma, c)y_n$$

or

$$(Ty_n, y_n) = \omega_n(\beta, \gamma, c) \quad (2.12)$$

and

$$\sum_{n=0}^{N-1} (Ty_n, y_n) = \sum_{n=0}^{N-1} (Te_n, e_n). \quad (2.13)$$

So, from (2.12) and (2.13), we get

$$\sum_{n=0}^{N-1} \omega_n(\beta, \gamma, c) = \sum_{n=0}^{N-1} (Te_n, e_n)$$

and

$$\sum_{n=0}^{N-1} \omega_n^k(\beta, \gamma, c) = \sum_{n=0}^{N-1} (T^k e_n, e_n), \quad k \geq 1,$$

or

$$\sum_{n=0}^{N-1} \omega_n^k(\beta, \gamma, c) = (T^k e_0, e_0) + (T^k e_1, e_1) + \sum_{n=2}^{N-1} (T^k e_n, e_n). \quad (2.14)$$

Since

$$T = C^{-1/2} T_0 C^{-1/2} + \beta C^{-1/2} P_0 C^{-1/2},$$

we obtain

$$\begin{aligned} T e_0 &= C^{-1/2} T_0 C^{-1/2} e_0 + \frac{\beta}{\gamma} e_0, \\ T e_1 &= C^{-1/2} T_0 C^{-1/2} e_1, \\ T e_n &= T_0 e_n, \quad 2 \leq n \leq N-1, \end{aligned}$$

and from (2.14), we get (2.10) because

$$\sum_{n=0}^{N-1} \lambda_n^k(c) = \sum_{n=0}^{N-1} (T_0^k e_n, e_n). \quad (2.15)$$

□

Remark 2.1. For $1 \leq k \leq 4$, equation (2.10) gives for $k = 1$:

$$\sum_{n=0}^{N-1} \omega_n(\beta, \gamma, c) = \sum_{n=0}^{N-1} \lambda_n(c) + \frac{\beta + \beta_{0+c}}{\gamma} - \beta_{0+c}, \quad (2.16)$$

for $k = 2$:

$$\sum_{n=0}^{N-1} \omega_n^2(\beta, \gamma, c) = \sum_{n=0}^{N-1} \lambda_n^2(c) + \frac{(\beta + \beta_{0+c})^2}{\gamma^2} + 2\alpha_{0+c}^2 \left(\frac{1}{\gamma} - 1 \right) - \beta_{0+c}^2, \quad (2.17)$$

for $k = 3$:

$$\begin{aligned} \sum_{n=0}^{N-1} \omega_n^3(\beta, \gamma, c) &= \sum_{n=0}^{N-1} \lambda_n^3(c) + \frac{(\beta + \beta_{0+c})^3}{\gamma^3} + \frac{3\alpha_{0+c}^2(\beta + \beta_{0+c})}{\gamma^2} \\ &\quad + \frac{3\alpha_{0+c}^2\beta_{1+c}}{\gamma} - \beta_{0+c}^3 - 3\alpha_{0+c}^2(\beta_{0+c} + \beta_{1+c}), \end{aligned} \tag{2.18}$$

and for $k = 4$:

$$\begin{aligned} \sum_{n=0}^{N-1} \omega_n^4(\beta, \gamma, c) &= \sum_{n=0}^{N-1} \lambda_n^4(c) + \frac{(\beta + \beta_{0+c})^4}{\gamma^4} + 2\alpha_{0+c}^4 \left(\frac{1}{\gamma^2} - 1 \right) \\ &\quad + \frac{4\alpha_{0+c}^2(\beta_{0+c} + \beta)^2}{\gamma^3} + \frac{2\alpha_{0+c}^2\beta^2}{\gamma^3} + \frac{4\alpha_{0+c}^2\beta_{1+c}^2}{\gamma} + \frac{4\alpha_{0+c}^2\beta_{1+c}}{\gamma^2}(\beta + \beta_{0+c}) \\ &\quad + 3 \left(\frac{1}{\gamma} - 1 \right) \alpha_{0+c}^2\alpha_{1+c}^2 - \beta_{0+c}^4 - 4\alpha_{0+c}^2(\beta_{0+c}^2 + \beta_{1+c}^2 + \beta_{0+c}\beta_{1+c}). \end{aligned} \tag{2.19}$$

3. Newton sum rules for the associated classical polynomials

(1) Consider the associated Laguerre polynomials $L_n^{(a)}(x; c)$, $a > -1$, $c \geq 0$, which satisfy the recurrence relation

$$\begin{aligned} (n + c + 1)L_{n+1}^{(a)}(x; c) + (n + c + a)L_{n-1}^{(a)}(x; c) \\ - (2n + 2c + a + 1)L_n^{(a)}(x; c) &= -xL_n^{(a)}(x; c), \quad n = 0, 1, 2, \dots, \\ L_{-1}^{(a)}(x; c) = 0, \quad L_0^{(a)}(x; c) &= 1. \end{aligned} \tag{3.1}$$

Setting $L_n^{(a)}(x; c) = (-1)^n \sqrt{U_n} Q_n^{(a)}(x; c)$ where U_n satisfies

$$U_{-1} = 0, \quad U_0 = 1, \quad U_n = \frac{n + c + a}{n + c} U_{n-1}, \quad n \geq 1,$$

we obtain from (3.1)

$$\begin{aligned} \sqrt{(n + c + 1)(n + c + a + 1)}Q_{n+1}^{(a)}(x; c) + \sqrt{(n + c + a)(n + c)}Q_{n-1}^{(a)}(x; c) \\ + (2n + 2c + a + 1)Q_n^{(a)}(x; c) &= xQ_n^{(a)}(x; c), \quad n \geq 1, \\ Q_{-1}^{(a)}(x; c) = 0, \quad Q_0^{(a)}(x; c) &= 1. \end{aligned} \tag{3.2}$$

The polynomials $Q_n^{(a)}(x; c)$ and the associated Laguerre polynomials $L_n^{(a)}(x; c)$ have the same zeros $\lambda_n(c)$. In this case, we have: $\alpha_{n+c}(a) = \sqrt{(n + c + 1)(n + c + a + 1)}$, $\beta_{n+c}(a) = (2n + 2c + a + 1)$, $n = 0, 1, 2, \dots, N - 1$, so, from the relations (2.6)–(2.9),

we get:

$$\sum_{n=0}^{N-1} \lambda_n(c) = N(N + 2c + a), \quad (3.3)$$

$$\sum_{n=0}^{N-1} \lambda_n^2(c) = 2N^3 + N^2(6c + 3a - 1) + N(6c^2 + 6ca - 2c + a^2 - a) - 2c^2 - 2ca, \quad (3.4)$$

$$\begin{aligned} \sum_{n=0}^{N-1} \lambda_n^3(c) &= 5N^4 + N^3(20c + 10a - 6) + N^2(30c^2 + 30ac - 18c + 6a^2 - 9a + 2) \\ &\quad + N(20c^3 + 30ac^2 - 18c^2 + 12a^2c - 18ac + 4c + a^3 - 3a^2 + 2a) \\ &\quad - (12c^3 + 18ac^2 + 6a^2c), \end{aligned} \quad (3.5)$$

and

$$\begin{aligned} \sum_{n=0}^{N-1} \lambda_n^4(c) &= 14N^5 + N^4(70c + 35a - 29) \\ &\quad + N^3(140c^2 + 140ac - 111c + 30a^2 - 58a + 22) \\ &\quad + N^2(140c^3 + 210ac^2 - 174c^2 + 90a^2c - 174ac + 66c + 10a^3 - 35a^2 + 33a - 6) \\ &\quad + N(70c^4 + 140ac^3 - 116c^3 + 90a^2c^2 + 174ac^2 - 66c^2 + 20a^3c - 70a^2c - 66ac \\ &\quad \quad - 12c + a^4 - 6a^3 + 11a^2 - 6a) \\ &\quad - (58c^4 + 116ac^3 + 70a^2c^2 + 12a^3c + 12ac). \end{aligned} \quad (3.6)$$

(2) Consider the associated Hermite polynomials defined by

$$H_{n+1}(x; c) + 2(n + c)H_{n-1}(x; c) = 2xH_n(x; c), \quad n = 0, 1, 2, \dots, \quad (3.7)$$

$$H_{-1}(x; c) = 0, \quad H_0(x; c) = 1.$$

Setting $H_n(x; c) = \sqrt{U_n}P_n(x; c)$ where U_n satisfies

$$U_{-1} = 0, \quad U_0 = 1, \quad U_n = 2(n + c)U_{n-1}, \quad n \geq 1,$$

we obtain from (3.7)

$$\sqrt{\frac{n+c+1}{2}}P_{n+1}(x; c) + \sqrt{\frac{n+c}{2}}P_{n-1}(x; c) = xP_n(x; c), \quad (3.8)$$

$$P_{-1}(x; c) = 0, \quad P_0(x; c) = 1.$$

The $P_n(x; c)$ and the associated Hermite polynomials $H_n(x; c)$ have the same zeros $\lambda_n(c)$. In this case, we have $\alpha_{n+c} = \sqrt{\frac{n+c+1}{2}}$, $\beta_{n+c} = 0$, $n = 0, 1, 2, \dots, N-1$, so, from the relations (2.6)–(2.9), we get

$$\sum_{n=0}^{N-1} \lambda_n(c) = \sum_{n=0}^{N-1} \lambda_n^3(c) = 0 \quad (3.9)$$

because $\beta_{n+c} = 0$,

$$\sum_{n=0}^{N-1} \lambda_n^2(c) = \frac{(N-1)(N+2c)}{2}, \tag{3.10}$$

and

$$\sum_{n=0}^{N-1} \lambda_n^4(c) = \frac{1}{4}(N-1)N(2N-3) + \frac{c}{2}(3N-5)(N+c). \tag{3.11}$$

(3) Consider the associated Jacobi polynomials $P_n^{(a,b)}(x; c)$, which are defined by

$$\begin{aligned} & 2(n+c+1)(n+c+a+b+1)(2n+2c+a+b)P_{n+1}^{(a,b)}(x; c) \\ & + 2(n+c+a)(n+c+b)(2n+2c+a+b+2)P_{n-1}^{(a,b)}(x; c) \\ & + (b^2 - a^2)(2n+2c+a+b+1)P_n^{(a,b)}(x; c) \\ & = (2n+2c+a+b+1)(2n+2c+a+b+2)(2n+2c+a+b)xP_n^{(a,b)}(x; c), \end{aligned} \tag{3.12}$$

$$P_{-1}^{(a,b)}(x; c) = 0, \quad P_0^{(a,b)}(x; c) = 1.$$

Setting $P_n^{(a,b)}(x; c) = U_n R_n^{(a,b)}(x; c)$ where

$$U_{-1} = 0, \quad U_0 = 1,$$

$$U_{n+1} = \sqrt{\frac{(n+c+a+1)(n+c+b+1)(2n+2c+a+b+1)}{(n+c+1)(n+c+a+b+1)(2n+2c+a+b+3)}} U_n,$$

we find

$$\begin{aligned} & \sqrt{\frac{4(n+c+1)(n+c+a+1)(n+c+b+1)(n+c+a+b+1)}{(2n+2c+a+b+2)^2(2n+2c+a+b+1)(2n+2c+a+b+3)}} R_{n+1}^{(a,b)}(x; c) \\ & + \sqrt{\frac{4(n+c)(n+c+a)(n+c+b)(n+c+a+b)}{(2n+2c+a+b)^2(2n+2c+a+b-1)(2n+2c+a+b+1)}} R_{n-1}^{(a,b)}(x; c) \\ & + \frac{(b^2 - a^2)}{(2n+2c+a+b+2)(2n+2c+a+b)} R_n^{(a,b)}(x; c) = x R_n^{(a,b)}(x; c), \end{aligned}$$

$$R_{-1}^{(a,b)}(x; c) = 0, \quad R_0^{(a,b)}(x; c) = 1. \tag{3.13}$$

The polynomials $R_n^{(a,b)}(x; c)$ and the associated Jacobi polynomials $P_n^{(a,b)}(x; c)$ have the same zeros $\lambda_n(c)$. In this case, we have

$$\alpha_{n+c}(a, b) = \sqrt{\frac{4(n+c+1)(n+c+a+1)(n+c+b+1)(n+c+a+b+1)}{(2n+2c+a+b+2)^2(2n+2c+a+b+1)(2n+2c+a+b+3)}},$$

$$\beta_{n+c}(a, b) = \frac{(b^2 - a^2)}{(2n+2c+a+b+2)(2n+2c+a+b)}, \quad n = 0, 1, 2, \dots, N-1,$$

and the relations (2.6), (2.7) give, respectively,

$$\sum_{n=0}^{N-1} \lambda_n(c) = \frac{N(b^2 - a^2)}{(2c+a+b)(2N+2c+a+b)}, \tag{3.14}$$

$$\begin{aligned}
\sum_{n=0}^{N-1} \lambda_n^2(c) &= \frac{(N-1)[2c(c+a+b+N) + N(a+b+1)]}{(2c+a+b+1)(2N+2c+a+b-1)} \\
&+ \frac{(b-a)^2[(3N-1)(b+a)^2(2c+a+b)(2N+2c+a+b)]}{2(2c+a+b+1)(2N+2c+a+b-1)(2c+a+b)^2(2N+2c+a+b)^2} \\
&- \frac{(b-a)^2[(N-1)(2c+a+b)^2(2N+2c+a+b)^2 + 2N^2(2N-1)(b+a)^2]}{2(2c+a+b+1)(2N+2c+a+b-1)(2c+a+b)^2(2N+2c+a+b)^2}.
\end{aligned} \tag{3.15}$$

Remark 3.1. For $c = 0$, in (3.3)–(3.6), (3.10), (3.11), (3.14), and (3.15), we get the known [5, 7] sum rules for the classical Laguerre, Hermite, and Jacobi polynomials, respectively, which were found using differential equations methods.

Remark 3.2. The sum rules (3.9), (3.10), (3.11) for the zeros of the associated Hermite polynomials have been found in [46], as an application of a program built in the Mathematica symbolic language.

4. Monotonicity properties of the zeros $\omega_n(\beta, \gamma, c)$

The operator (2.11) can be written

$$T = C^{-1/2}T_0C^{-1/2} + \beta C^{-1/2}P_0C^{-1/2}.$$

The eigenvalues $\omega_n(\beta, \gamma, c)$ of T , i.e., the zeros of the polynomial $Q_n(x; \beta, \gamma, c)$, are differentiable functions of β and the derivatives are given [20] by

$$\frac{\partial \omega_n(\beta, \gamma, c)}{\partial \beta} = (C^{-1/2}P_0C^{-1/2}y_n, y_n) = |(C^{-1/2}y_n, e_0)|^2 = |(y_n, C^{-1/2}e_0)|^2 \tag{4.1}$$

where $y_n, n = 0, 1, 2, \dots, N-1$, are the normalized eigenvectors of T which correspond to the eigenvalues $\omega_n(\beta, \gamma, c)$: $Ty_n = \omega_n(\beta, \gamma, c)y_n$.

The inequality

$$\frac{\partial \omega_n(\beta, \gamma, c)}{\partial \beta} \leq \frac{1}{\gamma} \tag{4.2}$$

follows immediately from (4.1) because $\|y_n\| = 1$ and $\|C^{-1/2}e_0\|^2 = \frac{1}{\gamma}$. We also observe that

$$\begin{aligned}
\frac{\partial}{\partial \beta} \left(\frac{\omega_n(\beta, \gamma, c)}{\beta} \right) &= \frac{\beta \frac{\partial \omega_n(\beta, \gamma, c)}{\partial \beta} - \omega_n(\beta, \gamma, c)}{\beta^2} = -\frac{1}{\beta^2} (C^{-1/2}T_0C^{-1/2}y_n, y_n) \\
&= -\frac{1}{\beta^2} (T_0x_n, x_n)
\end{aligned} \tag{4.3}$$

where $x_n = C^{-1/2}y_n$.

If $\lambda_0(c)$ is the smallest zero of $P_n(x; c)$, i.e., the first eigenvalue of T_0 , then the operator $T_0 + \lambda_0(c)$ is positive, i.e.,

$$((T_0 + \lambda_0(c))f, f) \geq 0$$

for every $f \in H_N$. Thus from (4.3), we find

$$\frac{\partial}{\partial \beta} \left(\frac{\omega_n(\beta, \gamma, c)}{\beta} \right) \leq \frac{\lambda_0(c)}{\beta^2} \|x_n\|^2 \leq \frac{\lambda_0(c)}{\beta^2} \|C^{-1/2}\|^2. \tag{4.4}$$

From the above, we obtain the following results:

(i) The function $\omega_n(\beta, \gamma, c)$ increases with β . This follows from (4.1).

(ii) The function $\omega_n(\beta, \gamma, c) - \frac{\beta}{\gamma}$ decreases with $\beta > 0$. This follows from (4.2) because

$$\frac{\partial}{\partial \beta}(\omega_n(\beta, \gamma, c) - \frac{\beta}{\gamma}) \leq 0. \tag{4.5}$$

(iii) If $\lambda_0(c) \leq 0$, then the function $\frac{\omega_n(\beta, \gamma, c)}{\beta}$ decreases with $\beta > 0$. This follows immediately from (4.4).

(iv) If $\lambda_0(c) > 0$, then the functions $\frac{\gamma\omega_n(\beta, \gamma, c) + \lambda_0(c)}{\beta}$ and $\frac{\omega_n(\beta, \gamma, c) + \lambda_0(c)}{\beta}$ decrease with $\beta > 0$ for $0 < \gamma \leq 1$ and for $\gamma > 1$, respectively. This follows from (4.4) because $\|C^{-1/2}\|^2 = \frac{1}{\gamma}$ for $0 < \gamma \leq 1$ and $\|C^{-1/2}\|^2 = 1$ for $\gamma > 1$.

Remark 4.1. From (4.5) we have for $\beta > 0$

$$\omega_n(\beta, \gamma, c) < \omega_n(0, \gamma, c) + \frac{\beta}{\gamma}. \tag{4.6}$$

This inequality gives an upper bound for all the zeros $\omega_n(\beta, \gamma, c)$, $n = 0, 1, 2, \dots, N-1$ of the scaled co-recursive associated polynomials, and the bound is sharp for small β because for $\beta = 0$, (4.6) becomes an equality.

Remark 4.2. From (4.1) we obtain

$$\frac{\partial \omega_n(\beta, \gamma, c)}{\partial \beta} = \frac{1}{\gamma} |(y_n, e_0)|^2$$

and

$$\sum_{n=0}^{N-1} \frac{\partial \omega_n(\beta, \gamma, c)}{\partial \beta} = \frac{1}{\gamma} \sum_{n=0}^{N-1} |(y_n, e_0)|^2 = \frac{1}{\gamma} \|e_0\|^2 = \frac{1}{\gamma}.$$

This means that

$$\frac{\partial}{\partial \beta} \left(\sum_{n=0}^{N-1} \omega_n(\beta, \gamma, c) - \frac{\beta}{\gamma} \right) = 0.$$

Thus we obtain the relation

$$\sum_{n=0}^{N-1} \omega_n(\beta, \gamma, c) - \frac{\beta}{\gamma} = \sum_{n=0}^{N-1} \omega_n(0, \gamma, c), \tag{4.7}$$

which follows also from the relation (2.16).

Acknowledgements. We thank the referees for their helpful remarks and suggestions and also for bringing references [9, 10, 35] to our attention.

References

1. W. R. Allaway, *The identification of a class of orthogonal polynomial sets*, PhD Thesis, University of Alberta, Canada, 1972.
2. R. Askey and J. Wimp, *Associated Laguerre and Hermite polynomials*, Proc. Roy. Soc. Edinburgh **96** (1984), 15–37.
3. S. Belmehdi and A. Ronveaux, *Polynomes associes des polynomes orthogonaux classiques. Construction via Reduce*, Actas del VI Simposium on Polinomios Orthogonales y Aplicaciones (1989), pp.72–83, Gijon, Spain.

4. ———, *The fourth-order differential equation satisfied by the associated orthogonal polynomials*, Rend. Math. Appl. **11** (1991), 313–326.
5. E. Buendia, J. S. Dehesa, and M. A. Sanchez-Buendia, *On the zeros of eigenfunctions of polynomial differential operators*, J. Math. Phys. **26** (1985), 2729–2736.
6. E. Buendia, J. S. Dehesa, and F. J. Galvez, *The distribution of zeros of the polynomial eigenfunctions of ordinary differential operators of arbitrary order*, Lect. Notes in Math. **1329** (1988), 222–235.
7. K. M. Case, *Sum rules for the zeros of polynomials I*, J. Math. Phys. **21** (1980), 702–708.
8. T. S. Chihara, *On co-recursive orthogonal polynomials*, Proc. Amer. Math. Soc. **8** (1957), 899–905.
9. J. S. Dehesa, *On the conditions for a Hamiltonian matrix to have an eigenvalue density with some prescribed characteristics*, J. Comput. Appl. Math. **2** (1976), 249–254.
10. ———, *The spectrum of Jacobi matrices in terms of its associated weight function*, J. Comput. Appl. Math. **4** (1978), 275–283.
11. ———, *The asymptotic behavior of zeros of orthogonal polynomials*, Letter Nuovo Cimento **24** (1979), 277–284.
12. J. S. Dehesa, E. Buendia, and M. A. Sanchez-Buendia, *On the polynomial solutions of ordinary differential equations of the fourth order*, J. Math. Phys. **26** (1985), 1547–1552.
13. E. A. van Doorn, *On a class of generalized orthogonal polynomials with real zeros*, in IMACS, Orthogonal Polynomials and Their Applications, pp.231-236, 1991.
14. ———, *Some new results for chain-sequence polynomials*, J. Comput. Appl. Math. **57** (1995), 309–317.
15. B. Germano and P. E. Ricci, *Representation formulas for the moments of the density of zeros of orthogonal polynomials*, Matematiche (Catania) **48** (1993), 77–86.
16. B. Germano, P. Natalini, and P. E. Ricci, *Computing the moments of the density of zeros for orthogonal polynomials*, Computers Math. Appl., Special issue on Concrete Analysis, **30** (1995), 69–81.
17. C. C. Grosjean, *Theory of recursive generation of systems of orthogonal polynomials: An illustrative example*, J. Comput. Appl. Math. **12, 13** (1985), 299–318.
18. ———, *The weight functions, generating functions and miscellaneous properties of the sequences of orthogonal polynomials of the second kind associated with Jacobi and the Gegenbauer polynomials*, J. Comput. Appl. Math. **16** (1986), 299–318.
19. W. Hahn, *Über Orthogonalpolynome mit drei parameteren*, Deutsche Math **5** (1940–41), 373–378.
20. E. K. Ifantis, *A theorem concerning differentiability of eigenvectors and eigenvalues with some applications*, Appl. Anal. **28** (1988), 257–283.
21. E. K. Ifantis and P. N. Panagopoulos, *On the zeros of a class of polynomials defined by a three term recurrence relation*, J. Math. Anal. Appl. **182** (1994), 361–370.
22. E. K. Ifantis and P. D. Siafarikas, *Differential inequalities on the greatest zero of Laguerre and ultraspherical polynomials*, Actas del VI Simposium on Polinomios Ortogonales y Aplicaciones, pp.187-197, Gijon, Spain, 1989.
23. ———, *On the zeros of a class of polynomials including the generalized Bessel polynomials*, in: Orthogonal Polynomials and Their Applications (Ed. J. S. Dehesa), J. Comput. Appl. Math. **49** (1993), 103–109.
24. ———, *Differential inequalities and monotonicity properties of the zeros of associated Laguerre and Hermite polynomials*, International Joint Symposium on Special Functions and Artificial Intelligence, Torino, Italy, 1993; Annals of Numerical Mathem. **2** (1995), 79–91.
25. ———, *Perturbation of the coefficients in the recurrence relation of a class of polynomials*, 4th International Symposium on Orthogonal Polynomials and Their Applications, Evian, France, 1992; J. Comput. Appl. Math. **57** (1995), 163–170.
26. ———, *An alternative proof of a theorem of Stieltjes and related results*, J. Comput. Appl. Math. **65** (1995), 165–172.
27. M. E. H. Ismail, D. R. Masson, J. Letessier, and G. Valent, *Birth and death processes and orthogonal polynomials*, in Orthogonal Polynomials: Theory and Practice (Ed. P. Nevai), pp.229–255, Kluwer, Dordrecht, 1990.
28. M. E. H. Ismail and M. E. Muldoon, *A discrete approach to monotonicity of zeros of orthogonal polynomials*, Trans. Amer. Math. Soc. **323**, (1991), 65–78.

29. J. Letessier, *Co-recursive associated Laguerre polynomials*, VII Simposium sobre Polinomios Ortogonales Aplicaciones, Granada, 1991; J. Comput. Appl. Math. **49** (1993), 127–136.
30. ———, *Some results on co-recursive associated Laguerre and Jacobi polynomials*, SIAM J. Math. Anal. **25** (1994), 528–548.
31. ———, *On co-recursive associated Jacobi polynomials*, 4th International Symposium on Orthogonal Polynomials and Their Applications, Evian, France, 1992; J. Comput. Appl. Math. **57**, (1995), 203–213.
32. S. Lewanowicz, *Results on the associated Jacobi and Gegenbauer polynomials*, in: Orthogonal Polynomials and Their Applications (Ed. J. S. Dehesa), J. Comput. Appl. Math., **49** (1993), 137–143.
33. F. Marcellan, J. S. Dehesa, and A. Ronveaux, *On orthogonal polynomials with perturbed recurrence relations*, J. Comput. Appl. Math. **30** (1990), 203–212.
34. F. Peherstorfer, *Finite perturbations of orthogonal polynomials*, J. Comput. Appl. Math. **44** (1992), 275–302.
35. I. V. V. Raghanacharyulu and A. R. Tekumalla, *Solution of the difference equations of generalized Lucas polynomials*, J. Math. Phys. **13** (1972), 321–324.
36. P. E. Ricci, *Sum rules for the zeros of polynomials and generalized Lucas polynomials*, J. Math. Phys. **34** (1993), 4884–4891.
37. A. Ronveaux, *Fourth-order differential equations for numerator polynomials*, J. Phys. A **21** (1988), 749–753.
38. ———, *Fourth-order differential equations and orthogonal polynomials of the Laguerre-Hahn class*, in: Orthogonal Polynomials and Their Applications (Eds. C. Brezinski, L. Gori, and A. Ronveaux), pp.379–385, IMACS Ann. Comput. Appl. Math. 9, Baltzer, Basel, 1991.
39. A. Ronveaux and F. Marcellan, *Co-recursive orthogonal polynomials and fourth-order differential equation*, J. Comput. Appl. Math. **25** (1989), 105–109.
40. A. Ronveaux, A. Zarzo, and E. Godoy, *Fourth-order differential equations satisfied by the generalized co-recursive of all classical orthogonal polynomials. A study of their distribution of zeros*, J. Comput. Appl. Math. **59** (1995), 295–328.
41. K. I. Sato, *On zeros of a system of polynomials and application to sojourn time distributions of birth-and-death processes*, Trans. Amer. Math. Soc. **309** (1988), 375–390.
42. H. A. Slim, *On co-recursive orthogonal polynomials and their application to potential scattering*, J. Math. Anal. Appl. **136** (1988), 1–19.
43. J. Wimp, *Explicit formulas for the associated Jacobi polynomials and some applications*, Canad. J. Math. **39** (1987), 983–1000.
44. A. Zarzo, *Estudio de las densidades discreta y asintotica de ceros de polinomios ortogonales*, Tesina de Licenciatura, Univ. Granada, 1990.
45. A. Zarzo, J. H. Dehesa, and A. Ronveaux, *Newton sum rules of zeros of semiclassical orthogonal polynomials*, J. Comput. Appl. Math. **33**, (1990), 85–96.
46. A. Zarzo, A. Ronveaux, and E. Godoy, *Fourth-order differential equation satisfied by the associated of any order of all classical orthogonal polynomials. A study of their distribution of zeros*, in: Orthogonal Polynomials and their Applications (Ed. J. S. Dehesa), J. Comput. Appl. Math. **49** (1993), 349–359.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF PATRAS, PATRAS, GREECE