

DARBOUX EQUATIONS IN EXTERIOR DOMAINS *

SAOUSSEN KALLEL-JALLOULI†

Abstract. We give sufficient conditions ensuring existence and regularity of a radial solution to the following equation

$$\begin{aligned} \det(\phi_{ij}) &= F(|x|, \phi, |\nabla\phi|), \text{ in } \Omega \\ \phi|_{\partial\Omega} &= c \end{aligned}$$

when Ω is an exterior domain.

1. Introduction. In this work, we consider the Dirichlet problem for real Monge-Ampère equations in exterior domains. More precisely, let $B \subset \mathbb{R}^n$ be an open ball, centered at the origin, that can be supposed, without loss of generality, to be the unit ball. Our purpose is to establish the existence of radial, convex solution $u \in C^2(\mathbb{R}^n \setminus B)$ of radially symmetric Monge-Ampère equation

$$\begin{cases} \det(\phi_{ij}) = F(|x|, \phi, |\nabla\phi|), & \text{in } \mathbb{R}^n \setminus \overline{B} \\ \phi|_{\partial B} = c \end{cases} \quad (1)$$

where F is a nonnegative continuous function. As usual, $|x|$ denotes the Euclidean length of $x = (x_1, \dots, x_n)$ and n is (all over this paper) the dimension of our Euclidean space. Additional hypothesis on F are described in §2.

When Ω is a strictly convex domain, this problem has received considerable study. Not many results are known about the solutions in unbounded domains. In the case when $F > 0$, F.Finster and O.C. Schnürer [2] proved the existence of smooth, strictly convex solution to (1) under some restrictions on F . We can also cite the work of T. Kusano and Ch.A. Swanson [3] related to radially symmetric two-dimensional elliptic Monge-Ampère equations.

Our attention will be directed toward the construction of radial solutions $u(x) = u(t)$ of (1), $t = |x|$. Direct computation (see [1]), shows that solving the equation (1) in C^2 is equivalent to solving the ordinary differential equation

$$\begin{cases} [(y')^n]' = nt^{n-1}F(t, y, y'), & \text{if } t > 1 \\ y(1) = c \end{cases} \quad (2)$$

Without loss of generality, we can take $c = 0$.

If we take as initial condition $y'(1) = 0$, we can easily transform (2) into the following integro-differential equation

$$y(r) = \int_1^r \left[\int_1^\rho nt^{n-1}F(t, y(t), y'(t)) dt \right]^{\frac{1}{n}} d\rho \quad (3)$$

*Received April 22, 2003; accepted for publication May 18, 2004.

†Faculté des Sciences, Campus Universitaire, 1060, Tunisie (Saoussen.Kallel@fst.rnu.tn).

EXAMPLE. Let $F(r) = (r-1)^{n-1-\varepsilon} r^{1-n}$, with $\varepsilon > 0$ small enough. Then, $u'(r) = \left[\frac{n}{n-\varepsilon}\right]^{\frac{1}{n}} (r-1)^{1-\frac{\varepsilon}{n}}$. In this case $F \in C^0$, but $u \notin C^2$.

This example shows that even when F depends only on r , it may not yield a C^2 solution, if F is allowed to vanish in the domain. This implies that we should place some restrictions on F .

Throughout this work, F satisfies some hypothesis be selected from the following list:

(H_1):

i) $F(t, y, z)$ is a nonincreasing function with respect to both y and z for each fixed (t, z) and (t, y) , respectively.

ii) $\int_1^{+\infty} t^{n-1} F(t, 0, 0) dt < +\infty$

(H_2): $F(t, y, z) \leq C_0 t^{-n-\alpha} |y|^\beta |z|^\theta$, with $\beta \geq 0$, $\alpha > \beta$, $\theta \geq 0$ and $C_0 \leq \frac{\alpha-\beta}{n}$ if $\beta + \theta = n$.

(H_3):

i) $F(t, y, z)$ is a nondecreasing function with respect to both y and z for each fixed (t, z) and (t, y) , respectively.

ii) There exists a constant $a > 0$ such that

$$\int_1^{+\infty} nt^{n-1} F(t, (t-1)a, a) dt \leq a^n$$

(H_4): $F(t, y, z) = (t-1)^l \tilde{F}(t, y, z)$, with $\tilde{F}(1, y, 0) \neq 0$, for $y \geq 0$, $l \geq n-1$, $\tilde{F} \in C^0$.

An example of a Monge-Ampère equation satisfying (H_3) is the Gauss curvature equation

$$\begin{cases} \det(u_{ij}) = p(|x|) u^\gamma (1 + |\nabla u|^2)^\delta, & x \in \mathbb{R}^n \setminus B \\ u|_{\partial B} = 1 \end{cases}$$

with $\gamma, \delta \geq 0$, $2\delta + \gamma < n$ and p is a non-négative function satisfying:

$$\int_1^{+\infty} t^{n+\gamma-1} p(t) dt < +\infty$$

In the following, \tilde{F} is used as introduced in (H_4). We shall prove

THEOREM A. If (H_4) and either (H_1), (H_2) or (H_3) holds, equation (1) has an infinitude of radial convex solutions $u \in C^2$ such that $\frac{u(x)}{|x|}$ has a positive finite limit at ∞ .

THEOREM B. If we suppose, in addition to the hypothesis of Theorem A, that

$$\tilde{F} \in C^k((\mathbb{R}^n \setminus B) \times \mathbb{R}^2) \quad (4)$$

and

$$\text{either } \frac{l+1}{n} \in \mathbb{N} \text{ or } \frac{l+1}{n} \geq k+1 \quad (5)$$

then the solutions given by theorem A are in C^{k+2}

2. Proof of theorem A. To prove the existence of a radially symmetric convex solution to the problem (1), we need to introduce the Frechet space C^1 of all continuously differentiable functions in $[1, +\infty[$, with the topology of uniform convergence of functions and their first derivatives on compact intervals. Consider now the closed convex subset \mathcal{K}_R of C^1

$$\mathcal{K}_R = \{y \in C^1 \mid y(1) = 0, 0 \leq y'(t) \leq R\} \quad (6)$$

and the operator $T : \mathcal{K}_R \rightarrow C^1$ defined by

$$T(y)(r) = \int_1^r \left[\int_1^\rho nt^{n-1}F(t, y(t), y'(t)) dt \right]^{\frac{1}{n}} d\rho, r \geq 1 \quad (7)$$

In order to prove that T has a fixed point $y \in \mathcal{K}_R$, we need to verify that T maps \mathcal{K}_R continuously into a relatively compact subset of \mathcal{K}_R .

If $y \in \mathcal{K}_R$, (7) implies that $T(y)(1) = 0$ and

$$0 \leq (Ty)'(r) = \left[\int_1^r nt^{n-1}F(t, y(t), y'(t)) dt \right]^{\frac{1}{n}}$$

We shall need to verify that we can find a constant $R > 0$ such that

$$\left[\int_1^{+\infty} ns^{n-1}F(s, y(s), y'(s)) ds \right]^{\frac{1}{n}} \leq R, \forall y \in \mathcal{K}_R \quad (8)$$

* If F satisfies (H_1) , we can write using $(H_1)(i)$,

$$(Ty)'(r) \leq \left[\int_1^r ns^{n-1}F(s, 0, 0) ds \right]^{\frac{1}{n}}$$

by $(H_1)(ii)$, it suffices then to take

$$R = \left[n \int_1^{+\infty} s^{n-1}F(s, 0, 0) ds \right]^{\frac{1}{n}}$$

and we get

$$(Ty)'(r) \leq R$$

* When F satisfies (H_2) , then, since

$$y(r) = \int_1^r y'(t) dt,$$

we get by (6),

$$|y(r)| \leq (r-1)R,$$

so,

$$(Ty)'(r) \leq \left[\int_1^r nC_0 s^{-\alpha-1} (s-1)^\beta R^{\beta+\theta} ds \right]^{\frac{1}{n}} \leq \left(\frac{n}{\alpha-\beta} C_0 \right)^{\frac{1}{n}} R^{\frac{\beta+\theta}{n}}$$

In order to get (8), it suffices to take R small enough when $(\beta+\theta) > n$, big enough when $(\beta+\theta) < n$. In the case when $\beta+\theta = n$ and $C_0 \leq \frac{\alpha-\beta}{n}$, any positive constant R lead to

$$(Ty)'(r) \leq R$$

* Finally, if F satisfies (H_3) , then, assumption (H_3) (i) shows that

$$(Ty)'(r) \leq \left[\int_1^r ns^{n-1} F(s, (s-1)R, R) ds \right]^{\frac{1}{n}}$$

it suffices then to take $R = a$ to ensure by (H_3) (ii) the inequality (8).

To establish the continuity of T , let (y_k) be a sequence in \mathcal{K}_R with $\lim_{k \rightarrow +\infty} y_k = y \in C^1$ in the C^1 -topology. By the dominated convergence theorem, we have then

$$\lim_{k \rightarrow +\infty} \int_1^r ns^{n-1} F(s, y_k(s), y'_k(s)) ds = \int_1^r ns^{n-1} F(s, y(s), y'(s)) ds$$

uniformly on $[1, +\infty[$, from which Ty_k and $(Ty_k)'$ converge uniformly to Ty and $(Ty)'$, respectively, on compact intervals in $[1, +\infty[$. this means that Ty_k converges to Ty in the C^1 -topology.

The relative compactness of $T(\mathcal{K}_R)$ is a consequence of Ascoli's Theorem; we need only verify the local uniform boundedness and local equicontinuity of the sets $T(\mathcal{K}_R)$ and $T(\mathcal{K}_R)' = \{(Ty)', y \in \mathcal{K}_R\}$.

Let us denote $G(t) = nF(t, u(t), u'(t))$ and $\tilde{G}(t) = n\tilde{F}(t, u(t), u'(t))$.

For every $y \in \mathcal{K}_R$, $1 \leq t_1 \leq t_2$, the inequality $a^{\frac{1}{n}} - b^{\frac{1}{n}} \leq (a-b)^{\frac{1}{n}}$, true for $a \geq b \geq 0$, implies

$$\begin{aligned} (Ty)'(t_2) - (Ty)'(t_1) &= \left(\int_1^{t_2} t^{n-1} G(t) dt \right)^{\frac{1}{n}} - \left(\int_1^{t_1} t^{n-1} G(t) dt \right)^{\frac{1}{n}} \\ &\leq \left(\int_{t_1}^{t_2} t^{n-1} G(t) dt \right)^{\frac{1}{n}} \end{aligned}$$

* If F satisfies (H_1) , then

$$G(t) \leq nF(t, 0, 0)$$

and

$$(Ty)'(t_2) - (Ty)'(t_1) \leq \left(\int_{t_1}^{t_2} nt^{n-1} F(t, 0, 0) dt \right)^{\frac{1}{n}} \rightarrow 0, \text{ as } t_1, t_2 \rightarrow \infty$$

* If F satisfies (H_2) , then,

$$\begin{aligned} (Ty)'(t_2) - (Ty)'(t_1) &\leq \left(\int_{t_1}^{t_2} nC_0 t^{n-1} t^{-n-\alpha} (t-1)^\beta R^{\beta+\theta} dt \right)^{\frac{1}{n}} \\ &\leq C_1 \left(\int_{t_1}^{t_2} t^{\beta-\alpha-1} dt \right)^{\frac{1}{n}} \rightarrow 0, \text{ as } t_1, t_2 \rightarrow \infty \end{aligned}$$

* Finally, when F satisfies (H_3) , then by (i), since $R = a$,

$$G(t) \leq nF(t, (t-1)a, a)$$

and

$$(Ty)'(t_2) - (Ty)'(t_1) \leq \left(\int_{t_1}^{t_2} nt^{n-1}F(t, (t-1)a, a) dt \right)^{\frac{1}{n}} \rightarrow 0, \text{ as } t_1, t_2 \rightarrow \infty$$

Then, in all these cases, for any compact interval I in $[1, +\infty[$ and arbitrary $\varepsilon > 0$, there is a corresponding $\delta > 0$, independent of t_1, t_2 and $y \in \mathcal{K}_R$, such that

$$|(Ty)'(t_2) - (Ty)'(t_1)| \leq \varepsilon$$

for all $t_1, t_2 \in I$ with $|t_1 - t_2| < \delta$.

The local equicontinuity of $T(\mathcal{K}_R)$ can be verified in the same way, and the local uniform boundedness is obvious.

Therefore the Schauder-Tychonoff fixed point theorem ([5]; lemma 1 and [6]; Theorem 4.5.1.) implies that T has a fixed point $u \in \mathcal{K}_R$, satisfying the integro-differential equation (3) for any R such that (8) holds. It remains to prove that $u' \in C^1$.

For $t > 1$, we have

$$u'(t) = \left[\int_1^t s^{n-1} (s-1)^l \tilde{G}(s) ds \right]^{\frac{1}{n}}$$

Since $\tilde{G}(1) \neq 0$, then $u' \in C^1]1, +\infty[$ and

$$\begin{aligned} u''(t) &= t^{n-1} (t-1)^l \tilde{G}(t) \left[\int_1^t s^{n-1} (s-1)^l \tilde{G}(s) ds \right]^{\frac{1}{n}-1} \\ &= t^{n-1} (t-1)^{\frac{l+1}{n}-1} \tilde{G}(t) \left[\int_0^1 [(t-1)s+1]^{n-1} s^l \tilde{G}((t-1)s+1) ds \right]^{\frac{1}{n}-1} \end{aligned}$$

which gives

$$\lim_{t \rightarrow 1^+} u''(t) = \begin{cases} 0, & \text{if } l > n-1 \\ \left[\frac{1}{l+1} \right]^{\frac{1}{n}-1} \tilde{G}(1)^{\frac{1}{n}} & \text{if } l = n-1 \end{cases}$$

Hence, $u \in C^2 [1, +\infty[$. It is not to be noted that u is a solution of (1) satisfying $u(1) = 0$ and $u'(1) = 0$.

Furthermore, the relation (3) and the inequality (8) imply that the limit

$$\lim_{t \rightarrow +\infty} \frac{u(t)}{t} = \lim_{t \rightarrow +\infty} u'(t) = \left[\int_1^{+\infty} ns^{n-1}F(s, u(s), u'(s)) ds \right]^{\frac{1}{n}}$$

is positive and finite, proving the asymptotic property in theorem A.

Since any non-negative constant b will serve as initial value $y'(1) = b$, there exists an infinitude of radial convex solutions to our problem.

3. Proof of theorem B. In this section, we study the regularity of the solution u given by theorem A. To prove the C^{k+2} regularity of u , let us proceed by induction on $k \in \mathbb{N}$. For $k = 0$, we have established in section 2, that $u \in C^2$. Suppose that

$$\tilde{F} \in C^{k-1} \Rightarrow u \in C^{k+1}$$

for some fixed $k \geq 1$. Assume now that $\tilde{F} \in C^k$. It follows in particular that $u \in C^{k+1}$. Hence, from the integral formula (7) and the hypothesis (H_4) , we get $u \in C^{k+2}]1, +\infty[$. It remains to check the regularity of u at the boundary $t = 1$.

The following preliminary result will be needed

LEMMA ([4] corollary 4.2). The k^{th} derivative of $g^{\frac{1}{n}}$, can be written as a sum of terms of the form

$$g^{\frac{1}{n}-\lambda} P_\lambda(g', g'', \dots, g^{(k+1-\lambda)})$$

where P_λ is a monomial of degree $\lambda \leq k$ and of weighted degree k .

Now, using the notation

$$H_y(t) = \int_0^1 [(t-1)s+1]^{n-1} s^l \widetilde{G}_y((t-1)s+1) ds, \quad (9)$$

we can write

$$u'(t) = (t-1)^{\frac{l+1}{n}} H_u^{\frac{1}{n}}(t)$$

where, by the induction hypothesis, $H_u \in C^k$. Then,

$$u^{(k+1)}(t) = \sum_{i=0}^k \binom{k}{i} \left[(t-1)^{\frac{l+1}{n}} \right]^{(i)} \left(H_u^{\frac{1}{n}} \right)^{(k-i)}(t)$$

furthermore, applying the above lemma, we get the following

$$\left(H_u^{\frac{1}{n}} \right)^{(l)} = \sum_{i=2}^l c_i H_u^{\frac{1}{n}-i} P_i(H'_u, \dots, H_u^{(l+1-i)}) + \frac{1}{n} H_u^{\frac{1}{n}-1} H_u^{(l)}, \quad \forall l \leq k$$

Since, $\forall j \leq k$,

$$(H_u)^{(j)}(t) = \sum_{i=0}^j c_{i,j} \int_0^1 \left([(t-1)s+1]^{n-1} \right)^{(j-i)} s^{l+i} \widetilde{G}_u^{(i)}((t-1)s+1) ds$$

it suffices then to prove that

$$f(t) = (t-1)^{\frac{l+1}{n}} h_k \in C^1([1, +\infty[)$$

where $h_k(t) = \int_0^1 [(t-1)s+1]^{n-1} s^{l+k} \widetilde{G}_u^{(k)}((t-1)s+1) ds$.

Differentiating f , yields

$$\forall t > 1, f'(t) = (t-1)^{\frac{l+1}{n}-1} \left[t^{n-1} \widetilde{G}_u^{(k)}(t) + c h_k(t) \right]$$

which implies

$$\lim_{t \rightarrow 1^+} f'(t) = \begin{cases} 0 & \text{if } l > n-1 \\ \widetilde{G}_u^{(k)}(1) \left[1 + c \int_0^1 s^{l+k} ds \right], & \text{if } l = n-1 \end{cases}$$

Consequently, $f \in C^1([1, +\infty[)$. Which completes the proof of theorem B.

Acknowledgement. The author would like to thank the referee for several helpful suggestions.

REFERENCES

- [1] PH. DELANOË, *Radially symmetric boundary value problems for real and complex elliptic Monge-Ampère equations*, Journal of Differential equations, 58 (1985), pp. 318–344.
- [2] F. FINSTER AND O.C. SCHNÜRER, *Hypersurfaces of prescribed Gauss curvature in exterior domains*, Calc. Var., 15 (2002), pp. 67–80.
- [3] T. KUSANO AND CH.A. SWANSON, *Existence theorems for elliptic Monge-Ampère equations in the plane*, Differential and Integral Equations, 3:3 (1990), pp. 487–493.
- [4] D. MONN, *Regularity of the complex Monge-Ampère equation for radially symmetric functions of the unit ball*, Math. Ann., 275 (1986), pp. 501–511.

- [5] A.J.B. POTTER, *An elementary version of the Leray-Schauder theorem*, J. Lond. Math. Soc., 2:5 (1972), pp. 414–416.
- [6] D.R. SMART, *Fixed point Theorems*, Cambridge Tracts in Mathematics, No. 66, Cambridge Univ. Press, London/New York, 1974.

