

Existence of hypersurfaces with prescribed mean curvature I – generic min-max*

XIN ZHOU[†] AND JONATHAN J. ZHU[‡]

We prove that, for a generic set of smooth prescription functions h on a closed ambient manifold, there always exists a nontrivial, smooth, closed hypersurface of prescribed mean curvature h . The solution is either an embedded minimal hypersurface with integer multiplicity, or a non-minimal almost embedded hypersurface of multiplicity one.

More precisely, we show that our previous min-max theory, developed for constant mean curvature hypersurfaces, can be extended to construct min-max prescribed mean curvature hypersurfaces for certain classes of prescription function, including a generic set of smooth functions, and all nonzero analytic functions. In particular we do not need to assume that h has a sign.

0. Introduction

Given a function $h : M \rightarrow \mathbb{R}$ on an ambient manifold M , a hypersurface $\Sigma \subset M$ has prescribed mean curvature h if its mean curvature satisfies

$$(0.1) \quad H_\Sigma = h|_\Sigma.$$

Prescribed mean curvature (PMC) hypersurfaces $\Sigma = \partial\Omega$ are critical points of the functional

$$(0.2) \quad \mathcal{A}^h = \text{Area} - \text{Vol}_h,$$

*The authors want to thank Prof. Richard Schoen and Prof. Bill Minicozzi for their interest, advice and encouragement on this work. The authors would like to thank the Institute for Advanced Study for their hospitality.

[†]The author is supported by NSF grant DMS-1811293 and an Alfred P. Sloan Research Fellowship.

[‡]The author is supported by NSF grant DMS-1607871. The author would also like to thank Boyu Zhang for helpful discussions, as well as Harvard University, where the majority of his contribution was carried out.

where $\text{Vol}_h(\Omega) = \int_{\Omega} h$ is the enclosed h -volume. PMC hypersurfaces are a canonical generalization of minimal and constant mean curvature (CMC) hypersurfaces, and have applications to physical phenomena such as capillary surfaces [22, §1.6]. Indeed, the local existence theory (or Dirichlet problem) for PMC hypersurfaces is quite well-understood, with several results extending naturally from the CMC to the PMC setting [35, 30, 31, 32, 60, 17]. The global theory or existence problem for closed PMC hypersurfaces, however, is to the authors' knowledge almost completely open for nonconstant prescription functions h .

In this article, we initiate a program to resolve the existence of closed PMC hypersurfaces, based on extending our min-max theory developed in [68] for CMC hypersurfaces. In particular, we prove that there exists a closed hypersurface of PMC h for a generic set of prescription functions h :

Theorem 0.1. *Let M^{n+1} be a smooth, closed Riemannian manifold of dimension $3 \leq n+1 \leq 7$. There is an open dense set $\mathcal{S} \subset C^{\infty}(M)$ of prescription functions h , for which there exists a nontrivial, smooth, closed, almost embedded hypersurface Σ^n of prescribed mean curvature h .*

In subsequent works, we plan to complete the global existence theory by approximating an arbitrary given smooth function $h \in C^{\infty}(M)$. Here a hypersurface is almost embedded if it locally decomposes into smooth sheets that touch but do not cross; in fact, our constructed PMC hypersurfaces Σ decompose into at most two sheets, and have touching set of codimension 1 - this is perhaps surprisingly the same regularity as we obtained in the CMC setting [68]. The dimension restriction comes from the regularity theory for stable minimal hypersurfaces [54] and is typical of variational methods for hypersurfaces.

After this article first appeared, the first author used our existence result for PMC hypersurfaces to prove the Multiplicity One Conjecture posed by Marques-Neves [48] for min-max minimal hypersurfaces. In his proof [67], it is crucial that our construction holds for a generic set of prescription functions h .

It was communicated to us by S.T. Yau that he conjectured the global existence problem for closed PMC surfaces in closed 3-manifolds in the 1980s; indeed he included the corresponding question for PMC surfaces in \mathbb{R}^3 in his 1982 problem list as [64, Problem 59]. The existence problem for closed PMC hypersurfaces may also be viewed as a higher dimensional *extended* Arnold-Novikov conjecture. The original question, which remains open, is the existence of closed embedded curves of prescribed constant geodesic curvature on a topological S^2 (see [26, 52, 53] for more backgrounds); Arnold

[6, 1996-17, 1996-18] later posed the natural extension to nonconstant prescribed curvature. Our previous theory [68] has already completely resolved the higher dimensional problem for constant (mean) curvature.

In Theorem 0.1, we use the set $\mathcal{S} = \mathcal{S}_{M,g}$ of Morse functions whose zero locus is a (possibly empty) hypersurface whose mean curvature vanishes to finite order. The mean curvature flow gives us a neat argument to show that this set is indeed generic (see Proposition 3.7 for the proof):

Proposition 0.2. *Let (M, g) be a smooth closed Riemannian manifold. Consider the set \mathcal{S} of smooth Morse functions h such that the zero set $\{h = 0\} =: \Sigma_0$ is a smooth closed hypersurface, and the mean curvature of Σ_0 vanishes to at most finite order. Then \mathcal{S} is open and dense in $C^\infty(M)$.*

In fact, our min-max theory is powerful enough to handle still more general prescription functions, namely those that satisfy any of the following conditions:

- (†) If $\Sigma^n \subset M^{n+1}$ is a smoothly embedded hypersurface and $h|_\Sigma$ vanishes to all orders at $p \in \Sigma$, then there exists $r > 0$ for which $\Sigma_0 = \{h = 0\} \cap B_r(p)$ is a connected, smoothly embedded hypersurface passing through p . Moreover, if the mean curvature of Σ_0 vanishes to infinite order at any point, then it vanishes identically.
- (‡) The zero set of $h : M^{n+1} \rightarrow \mathbb{R}$ is contained in a countable union of connected, smoothly embedded $(n - 1)$ -dimensional submanifolds.
- (RA) h is a real analytic function on a real analytic manifold M .

Any function in \mathcal{S} satisfies (†). A general function satisfying (†), however, could have nontrivial minimal hypersurfaces in its zero set. In this case each component of our constructed PMC hypersurface is either one of these (embedded) minimal components or a definitively non-minimal component, which is almost embedded with codimension 1 touching set. Note, however, that (‡) precludes the possibility of minimal PMC hypersurfaces.

The local existence theory of PMC hypersurfaces is a natural problem for geometric PDE, and has been fairly well-understood by parametric methods, such as [35, 30, 31, 32, 60, 17, 18], and also non-parametric methods [25, 23, 27, 5]. For the latter, however, graphical PMC hypersurfaces have typically been studied only for prescription functions independent of the vertical direction. Yau's conjecture for PMC surfaces in \mathbb{R}^3 inspired some important partial results such as [61, 24, 65]. Finally we take care to mention the results of [19], obtained through a homological (currents) approach.

In order to prove Theorem 0.1, we consider the variational properties of PMC hypersurfaces as critical points of the \mathcal{A}^h -functional. Because we are interested in the global problem of finding closed PMC hypersurfaces, it is far from clear direct minimization of the \mathcal{A}^h functional can produce a solution other than the empty domain $\mathcal{A}^h(\emptyset) = 0$ or the total manifold $\mathcal{A}^h(M) = -\int_M h$. Therefore, the min-max method becomes the natural way to find nontrivial critical points of \mathcal{A}^h .

The min-max theory for minimal submanifolds was initiated by Almgren [3], and has been a remarkably successful tool in the study of the area functional. Using methods of geometric measure theory, Almgren was able to prove the existence of a nontrivial weak solution as stationary integral varifolds [4] in any dimension and codimension. In codimension one, higher regularity was established by Pitts [51] (for $2 \leq n \leq 5$), and later by Schoen-Simon [37] (for $n \geq 6$). Colding-De Lellis [13] established the corresponding theory using smooth sweepouts based on ideas of Simon-Smith [58]. The preceding body of work completely resolved the $h \equiv 0$ case of Theorem 0.1.

Recently, Marques-Neves [45, 1, 47] used the Almgren-Pitts min-max theory to resolve a number of longstanding open problems in geometry, including their celebrated proof of the Willmore conjecture. Consequently, the min-max program has seen a series of developments in various directions, including (but not limited to) [44, 16, 29, 38, 46, 43, 15, 42, 37, 12, 48, 59]. In [68], we extended the min-max theory to study the \mathcal{A}^c functional ($h \equiv c$) for the construction of CMC hypersurfaces. The present work thus represents a natural continuation of our work, further extending the min-max theory to the PMC setting and the study of the \mathcal{A}^h functional.

0.1. Min-max procedure

We now give a heuristic overview of our min-max method. In the main proofs, for technical reasons we will work with discrete families as in Almgren-Pitts, but here we will describe the key ideas using continuous families to elucidate those ideas¹.

Let M, h be as in Theorem 0.1. The \mathcal{A}^h functional (0.2) is defined on open sets Ω with rectifiable boundary by $\mathcal{A}^h(\Omega) = \text{Area}(\partial\Omega) - \int_\Omega h$. Denote $I = [0, 1]$. Consider a continuous 1-parameter family of sets with rectifiable boundary

$$\{\Omega_x : x \in I\}, \text{ with } \Omega_0 = \emptyset \text{ and } \Omega_1 = M.$$

¹We note that a version of this min-max theory using continuous sweepouts was established (based on this article) by the first author in [67] after this article first appeared on arXiv.

Fix such a family $\{\Omega_x^0\}$, and consider its homotopy class $[\{\Omega_x^0\}] = \{\{\Omega_x\} \sim \{\Omega_x^0\}\}$. The h -min-max value (or h -width) is defined as

$$\mathbf{L}^h = \inf_{\{\Omega_x\} \sim \{\Omega_x^0\}} \max\{\mathcal{A}^h(\Omega_x) : x \in I\}.$$

A sequence $\{\{\Omega_x^i\} : i \in \mathbb{N}\}$ with $\max_{x \in I} \mathcal{A}^h(\Omega_x^i) \rightarrow \mathbf{L}^h$ is typically called a *minimizing sequence*, and any sequence $\{\Omega_{x_i}^i : x_i \in (0, 1), i \in \mathbb{N}\}$ with $\mathcal{A}^h(\Omega_{x_i}^i) \rightarrow \mathbf{L}^h$ is called a *min-max sequence*.

Our main result (for the precise statement see Theorem 4.8) then says that there is a nice minimizing sequence $\{\{\Omega_x^i\} : i \in \mathbb{N}\}$, and some min-max sequence $\{\Omega_{x_i}^i : x_i \in (0, 1), i \in \mathbb{N}\}$, such that:

Theorem 0.3. *Suppose that h satisfies (\dagger) , (\ddagger) or (RA) and that $\int_M h \geq 0$. Then the sequence $\partial\Omega_{x_i}^i$ converges as varifolds to a nontrivial, smooth, closed, almost embedded hypersurface Σ of prescribed mean curvature h . Each component of Σ is either a closed, embedded minimal hypersurface in the zero set of h ; or an almost embedded hypersurface on which the convergence is multiplicity 1.*

If h satisfies (\ddagger) , only the latter case may occur.

Here, and for the remainder of this article, we make the assumption that $\int_M h \geq 0$. This ensures that $\mathcal{A}_h(M) \leq 0$ and hence any positive width sweepout must have a nontrivial maximal slice; we can always guarantee this condition by changing the sign of h , and doing so does not affect the existence result, since a two-sided hypersurface with PMC h will have PMC $-h$ under its opposite orientation.

Our proof broadly follows the Almgren-Pitts scheme, but with several important difficulties. An important observation is that several of the innovations we developed in [68] may be used to handle the \mathcal{A}^h functional even for nonconstant h . However, in the present setting we must develop the background compactness and regularity theory for PMC hypersurfaces, and when h is allowed to have a significant zero set it becomes crucial to obtain good control of any touching phenomena.

To describe these challenges, we first review the Almgren-Pitts min-max method, which may be organized into five broad steps as follows:

- Construct a sweepout with positive width, and extract a minimizing sequence;
- Apply a ‘tightening’ map to construct a new sequence whose varifold limit satisfies a variational property and an ‘almost-minimizing’ property;

- Use these properties to construct ‘replacements’ on annuli which must be regular;
- Apply successive concentric annular replacements to the min-max limit and show that they coincide with each other, and hence extend to the center;
- Show that the min-max limit coincides with the replacement near the center.

Given a minimizing sequence, in the minimal ($h \equiv 0$) setting one constructs a new ‘tightened’ sequence, for which any min-max (varifold) limit must be stationary - that is, a weak solution in the sense of first variations. The \mathcal{A}^h functional, however, is not well-defined on varifolds, so it is not straightforward to formulate a notion of weak solution for its critical points. In the PMC setting, we are able to show that the min-max limit V (after tightening) has first variation bounded by $c = \sup |h|$. As in [68], it is an important and delicate observation that this relatively loose variational property still provides enough control to develop the remaining regularity theory.

We also show that the limit V is *h-almost minimizing*, which we formulate as property of being (a limit of) constrained almost-minimizers for the \mathcal{A}^h functional (see Definition 6.1). To construct (*h-*) *replacements* in a subset $U \subset M$, one then solves the corresponding series of constrained minimization problems. For the \mathcal{A}^h functional, each (local) \mathcal{A}^h -minimizer will be an open set Ω_i^* with stable, regular PMC boundary in U , and the *h*-replacement V^* is obtained as the varifold limit $\lim |\partial\Omega_i^*|$. At this point, it is clear that a good compactness theory for PMC hypersurfaces is essential for the regularity of V^* .

To establish the regularity of V , successive replacements V^* and V^{**} are applied on two overlapping concentric annuli A_1 and A_2 . The goal is to show that the replacements glue together smoothly on the overlapping region and may thus be extended all the way to the center, by taking further replacements. Here the main technical issue is to preserve the regularity and uniqueness across the gluing interface.

Finally, one would like to prove that the min-max limit V coincides with the extended replacement V^* near the center. In the minimal setting, one can appeal to the constancy theorem. Since in the PMC setting we also make an assertion about the multiplicity of V , we instead directly prove that V^* has at worst a removable singularity at the center, then use a moving sphere argument to show that the densities of V and V^* are the same in the annular region.

The primary concern in the PMC setting is the compactness and regularity theory for PMC hypersurfaces. Whilst it is not too difficult to show that smooth limits of PMC hypersurfaces remain almost embedded, for the moving sphere and gluing arguments above it is essential that we may avoid the bulk of any touching singularities of the replacement hypersurfaces by choosing the gluing interface to be transverse to the touching set. If the touching sets are small enough - $(n - 1)$ -rectifiable, for instance - then we may indeed achieve this by Sard's lemma.

Unlike the minimal or CMC settings, however, in the PMC setting we may not have a complete two-sided or even one-sided maximum principle. Moreover, whilst the difference of PMC hypersurfaces with opposite orientations will satisfy an elliptic PDE, there is an inhomogeneous term which does not have a sign for general h . The danger that arises is the possibility of infinite order but non-identical touching, for which the tamest example is two PMC sheets sticking together as a minimal hypersurface on a small subset. To overcome this, we use unique continuation for elliptic differential *inequalities* to prove the necessary compactness theorem: Under assumptions (\dagger) or (\ddagger) , any infinite order touching for the limit implies that all sheets are identical, minimal and contained in the zero set of h .

The possibility of minimal components also presents a technical issue for the gluing step, as again one would like to prevent non-minimal sheets gluing together into a higher multiplicity minimal sheet. We can rule this out by first obtaining a putative gluing to a smooth PMC hypersurface, then using unique continuation results coming from our work to control the touching set.

To complete the details of the min-max procedure, we adapt several ideas that we previously introduced to handle the CMC setting [68], demonstrating also the flexibility of those techniques. For instance, an essential observation is that the h -volume is still of lower order than the area term in the \mathcal{A}^h functional. This immediately allows us to show that \mathbf{L}^h is positive on any sweepout, as a consequence of the isoperimetric inequality for small volumes (see Theorems 2.3 and 4.9). Moreover, an important issue in both the CMC and PMC settings is that the total mass of the replacement V^* may differ from the total mass of the original varifold V ; however the mass defect is again controlled by the higher order term $\|h\|_\infty \text{Vol}(U)$, which converges to zero under any blowup process. Using this insight, we are able to prove that any blowup of the min-max limit V has the good replacement property of Colding-De Lellis, and is therefore regular (see Proposition 6.10); in particular the tangent cones of V are always planes.

Finally, to complete the gluing in the non-minimal setting, we must nevertheless work around a nontrivial touching set at the gluing interface. Again the compactness theory for PMC hypersurfaces is key, and an important step is to show that the second replacement V^{**} may be represented by a boundary in $A_1 \cup A_2$. This yields that V^* and V^{**} glue together - with matching orientations - as desired. Near the touching set, we then use the graphical decomposition into embedded sheets together with the gluing along regular part to properly match the sheets together.

0.2. Outline of the paper

Our basic notation and background material is described in Section 1. Then in Section 2 we recall some preliminary results including the regularity for \mathcal{A}^h -minimizers.

In Section 3, we describe curvature estimates and compactness for almost embedded PMC hypersurfaces, including unique continuation lemmas and estimates of the touching set for functions satisfying (\dagger) , (\ddagger) or (RA) . We also prove there the genericness of condition (\dagger) , that is, Proposition 0.2.

In Section 4 we formulate the precise min-max procedure, and prove the existence of nontrivial (positive width) sweepouts. Then in Section 5 we review the tightening process for varifolds of bounded first variation and its consequences for the \mathcal{A}^h functional. In Section 6 we further observe that the replacement theory for constant prescribed mean curvature extends to an h -replacement theory for our suitable functions h .

Finally, in Section 7 we complete the regularity of the min-max varifold.

1. Notation

In this section, we collect some notions. We refer to [57] and [51, §2.1] for further materials in geometric measure theory.

Let (M^{n+1}, g) denote a closed, oriented, smooth Riemannian manifold of dimension $3 \leq (n+1) \leq 7$. Assume that (M, g) is embedded in some \mathbb{R}^L , $L \in \mathbb{N}$. $B_r(p)$, $\tilde{B}_r(p)$ denote respectively the Euclidean ball of \mathbb{R}^L or the geodesic ball of (M, g) . We denote by \mathcal{H}^k the k -dimensional Hausdorff measure; $\mathbf{I}_k(M)$ the space of k -dimensional integral currents in \mathbb{R}^L with support in M ; $\mathcal{Z}_k(M)$ the space of integral currents $T \in \mathbf{I}_k(M)$ with $\partial T = 0$; $\mathcal{V}_k(M)$ the closure, in the weak topology, of the space of k -dimensional rectifiable varifolds in \mathbb{R}^L with support in M ; $G_k(M)$ the Grassmannian bundle of un-oriented k -planes over M ; \mathcal{F} and \mathbf{M} respectively the flat norm [57, §31] and mass norm [57, 26.4] on $\mathbf{I}_k(M)$; \mathbf{F} the varifold \mathbf{F} -metric on

$\mathcal{V}_k(M)$ and currents \mathbf{F} -metric on $\mathbf{I}_k(M)$, [51, 2.1(19)(20)]; $\mathcal{C}(M)$ or $\mathcal{C}(U)$ the space of sets $\Omega \subset M$ or $\Omega \subset U \subset M$ with finite perimeter (Caccioppoli sets), [57, §14], [28, §1.6]; and $\mathfrak{X}(M)$ or $\mathfrak{X}(U)$ the space of smooth tangent vector fields on M or supported in U .

We also utilize the following definitions:

- a) Given $T \in \mathbf{I}_k(M)$, $|T|$ and $\|T\|$ denote respectively the integral varifold and Radon measure in M associated with T ;
- b) Given $c > 0$, a varifold $V \in \mathcal{V}_k(M)$ is said to have *c-bounded first variation in an open subset $U \subset M$* , if

$$|\delta V(X)| \leq c \int_M |X| d\mu_V, \quad \text{for any } X \in \mathfrak{X}(U);$$

here the first variation of V along X is $\delta V(X) = \int_{G_k(M)} \text{div}_S X(x) dV(x, S)$, [57, §39];

- c) $U_r(V)$ denotes the ball in $\mathcal{V}_k(M)$ under \mathbf{F} -metric with center $V \in \mathcal{V}_k(M)$ and radius $r > 0$;
- d) Given $p \in \text{spt } \|V\|$, $\text{VarTan}(V, p)$ denotes the space of tangent varifolds of V at p , [57, 42.3];
- e) Given a smooth, immersed, closed, orientable hypersurface Σ in M , or a set $\Omega \in \mathcal{C}(M)$ with finite perimeter, $[[\Sigma]]$, $[[\Omega]]$ denote the corresponding integral currents with the natural orientation, and $[\Sigma]$ denotes the corresponding integer-multiplicity varifolds;
- f) $\partial\Omega$ denotes the (reduced)-boundary of $[[\Omega]]$ as an integral current, and $\nu_{\partial\Omega}$ denotes the outward pointing unit normal of $\partial\Omega$, [57, 14.2].

In this paper, we are interested in the following weighted area functional defined on $\mathcal{C}(M)$. Given $h : M \rightarrow \mathbb{R}$, define the \mathcal{A}^h -functional on $\mathcal{C}(M)$ as

$$(1.1) \quad \mathcal{A}^h(\Omega) = \mathcal{H}^n(\partial\Omega) - \int_{\Omega} h.$$

The *first variation formula* for \mathcal{A}^h along $X \in \mathfrak{X}(M)$ is (see [57, 16.2])

$$(1.2) \quad \delta \mathcal{A}^h|_{\Omega}(X) = \int_{\partial\Omega} \text{div}_{\partial\Omega} X d\mu_{\partial\Omega} - \int_{\partial\Omega} h \langle X, \nu \rangle d\mu_{\partial\Omega},$$

where $\nu = \nu_{\partial\Omega}$ is the outward unit normal on $\partial\Omega$.

When the boundary $\partial\Omega = \Sigma$ is a smooth immersed hypersurface, we have

$$\text{div}_{\Sigma} X = H \langle X, \nu \rangle,$$

where H is the mean curvature of Σ with respect to ν ; if Ω is a critical point of \mathcal{A}^h , then (1.2) directly implies that $\Sigma = \partial\Omega$ must have mean curvature $H = h|_\Sigma$ with respect to the outward unit normal ν . In this case, we can calculate the *second variation formula* for \mathcal{A}^h along normal vector fields $X \in \mathfrak{X}(M)$ such that $X = \varphi\nu$ along $\partial\Omega = \Sigma$ where $\varphi \in C^\infty(\Sigma)$, [9, Proposition 2.5],

$$(1.3) \quad \begin{aligned} \delta^2 \mathcal{A}^h|_\Omega(X, X) &= II_\Sigma(\varphi, \varphi) \\ &= \int_\Sigma (|\nabla\varphi|^2 - (Ric^M(\nu, \nu) + |A^\Sigma|^2 + \partial_\nu h) \varphi^2) d\mu_\Sigma. \end{aligned}$$

In the above formula, $\nabla\varphi$ is the gradient of φ on Σ ; Ric^M is the Ricci curvature of M ; A^Σ is the second fundamental form of Σ .

2. Preliminaries

In this section, we collect some preliminary results. We present a maximum principle for varifolds with bounded first variation, a regularity result for boundaries that minimize the \mathcal{A}^h -functional, and a result on isoperimetric profile for small volumes.

2.1. Maximum principle for varifolds with c -bounded first variation

We will need the following maximum principle which is essentially due to White [63, Theorem 5].

Proposition 2.1 (Maximum principle for varifolds with c -bounded first variation). *Suppose $V \in \mathcal{V}_n(M)$ has c -bounded first variation in a open subset $U \subset M$. Let $K \subset U$ be an open subset with compact closure in U , such that $\text{spt}(\|V\|) \subset K$, and*

- (i) ∂K is smoothly embedded in M ,
- (ii) the mean curvature of ∂K with respect to the outward pointing normal is greater than c .

Then $\text{spt}(\|V\|) \cap \partial K = \emptyset$.

2.2. Regularity for boundaries which minimize the \mathcal{A}^h functional

The following result about regularity of boundaries which minimize the \mathcal{A}^h functional can be found in [49].

Theorem 2.2. *Given $\Omega \in \mathcal{C}(M)$, $p \in \text{spt } \|\partial\Omega\|$, and some small $r > 0$, suppose that $\Omega \llcorner \tilde{B}_s(p)$ minimizes the \mathcal{A}^h -functional: that is, for any other $\Lambda \in \mathcal{C}(M)$ with $\text{spt } \|\Lambda - \Omega\| \subset \tilde{B}_s(p)$, we have $\mathcal{A}^h(\Lambda) \geq \mathcal{A}^h(\Omega)$. Then except for a set of Hausdorff dimension at most $n - 7$, $\partial\Omega \llcorner \tilde{B}_s(p)$ is a smooth and embedded hypersurface, and is real analytic if the ambient metric on M is real analytic.*

Proof. The regularity will follow from the arguments in [49, Section 3] - in particular [49, Corollary 3.7, 3.8] - so long as we can verify condition [49, 3.1(2)]. That is, setting $A = \mathcal{H}^n(\partial\Omega \cap \tilde{B}_r(p))$ and $A' = \mathcal{H}^n(\partial\Lambda \cap \tilde{B}_r(p))$, it suffices to prove that, for small enough r ,

$$(2.1) \quad A' - A \geq -CrA,$$

for all $\Lambda \in \mathcal{C}(M)$ as in the statement of the theorem. Indeed, for such Λ , since $\Omega \llcorner \tilde{B}_r(p)$ minimizes the \mathcal{A}^h -functional in $\tilde{B}_r(p)$ we have that

$$(2.2) \quad A' - A \geq - \left| \int_{\Lambda} h - \int_{\Omega} h \right| \geq -c\mathcal{H}^{n+1}(\Lambda \Delta \Omega),$$

where $c = \sup |h|$ and $\Lambda \Delta \Omega$ is the symmetric difference.

In fact this essentially replaces condition [49, 3.1(1)], and we finish the proof in the same way, by estimating $\mathcal{H}^{n+1}(\Lambda \Delta \Omega) \leq 2\omega_{n+1}r^{n+1}$, where ω_n is the volume of the unit n -ball and, by the isoperimetric inequality, $\mathcal{H}^{n+1}(\Lambda \Delta \Omega) \leq C\mathcal{H}^n(\partial(\Lambda \Delta \Omega))^{\frac{n+1}{n}} \leq C(A' + A)^{\frac{n+1}{n}}$. Combining these estimates with (2.2) yields that

$$A' - A \geq -Cr(A' + A),$$

which is clearly equivalent to (2.1) for small r . This completes the proof. \square

2.3. Isoperimetric profiles for small volume

We will use the following consequence of the fact that the isoperimetric profile is asymptotically Euclidean for small volumes [11] (see also [50, Theorem 3]). Note that the result indeed holds for any $\Omega \in \mathcal{C}(M)$ by using the regularity theory for isoperimetric domains (c.f. Theorem 2.2).

Theorem 2.3. *There exists constants $C_0 > 0$ and $V_0 > 0$ depending only on (M, g) such that*

$$\text{Area}(\partial\Omega) \geq C_0 \text{Vol}(\Omega)^{\frac{n}{n+1}}, \text{ whenever } \Omega \in \mathcal{C}(M) \text{ and } \text{Vol}(\Omega) \leq V_0.$$

3. Stable PMC hypersurfaces

In this section, we establish curvature estimates and the compactness theory for stable hypersurfaces of prescribed mean curvature $h : M \rightarrow \mathbb{R}$. In particular, when h satisfies any of the assumptions (\dagger) , (\ddagger) or (RA) , we obtain good control on the touching set that arises upon taking the limit of embedded stable PMC hypersurfaces.

3.1. Stability and curvature estimates

Definition 3.1. Let Σ be a smooth, immersed, two-sided hypersurface with unit normal vector ν , and $U \subset M$ an open subset. We say that Σ is a *stable h -hypersurface* in U if

- the mean curvature H of $\Sigma \cap U$ with respect to ν equals to $h|_{\Sigma}$; and
- $II_{\Sigma}(\varphi, \varphi) \geq 0$ for all $\varphi \in C^{\infty}(\Sigma)$ with $\text{spt } \varphi \subset \Sigma \cap U$, where II_{Σ} is as in (1.3).

Definition 3.2. Let Σ_i , $i = 1, 2$, be connected embedded hypersurfaces in a connected open subset $U \subset M$, with $\partial\Sigma_i \cap U = \emptyset$ and unit normals ν_i . We say that Σ_2 *lies on one side of* Σ_1 if Σ_1 divides U into two connected components $U_1 \cup U_2 = U \setminus \Sigma_1$, where ν_1 points into U_1 , and either:

- $\Sigma_2 \subset \text{Clos}(U_1)$, which we write as $\Sigma_1 \leq \Sigma_2$ or that Σ_2 lies on the positive side of Σ_1 ; or
- $\Sigma_2 \subset \text{Clos}(U_2)$, which we write as $\Sigma_1 \geq \Sigma_2$ or that Σ_2 lies on the negative side of Σ_1 .

Definition 3.3 (Almost embedding). Let $U \subset M^{n+1}$ be an open subset, and Σ^n be a smooth n -dimensional manifold. A smooth immersion $\phi : \Sigma \rightarrow U$ is said to be an *almost embedding* if at any point $p \in \phi(\Sigma)$ where Σ fails to be embedded, there is a small neighborhood $W \subset U$ of p , such that

- $\Sigma \cap \phi^{-1}(W)$ is a disjoint union of connected components $\cup_{i=1}^l \Sigma_i$;
- $\phi(\Sigma_i)$ is an embedding for each $i = 1, \dots, l$;
- for each i , any other component $\phi(\Sigma_j)$, $j \neq i$, lies on one side of $\phi(\Sigma_i)$ in W .

We will simply denote $\phi(\Sigma)$ by Σ and denote $\phi(\Sigma_i)$ by Σ_i . The subset of points in Σ where Σ fails to be embedded will be called the *touching set*, and denoted by $\mathcal{S}(\Sigma)$. We will call $\Sigma \setminus \mathcal{S}(\Sigma)$ the regular set, and denote it by $\mathcal{R}(\Sigma)$.

Remark 3.4. From the definition, the collection of components $\{\Sigma_i\}$ meet tangentially along $\mathcal{S}(\Sigma)$.

Definition 3.5 (Almost embedded h -boundaries).

1. An almost embedded hypersurface $\Sigma \subset U$ is said to be a *boundary* if there is an open subset $\Omega \in \mathcal{C}(U)$, such that Σ is equal to the boundary $\partial\Omega$ (in U) in the sense of currents;
2. The *outer unit normal* ν_Σ of Σ is the choice of the unit normal of Σ which points outside of Ω along the regular part $\mathcal{R}(\Sigma)$;
3. Σ is called a *stable h -boundary* if Σ is a boundary as well as a stable immersed h -hypersurface.

We have the following variant of the famous Schoen-Simon-Yau (for $2 \leq n \leq 5$) [55] and Schoen-Simon ($n = 6$) [54] curvature estimates.

Theorem 3.6 (Curvature estimates for stable h -hypersurfaces). *Let $2 \leq n \leq 6$, and $U \subset M$ be an open subset. If $\Sigma \subset U$ is a smooth, immersed (almost embedded when $n = 6$), two-sided, stable h -hypersurface in U with $\partial\Sigma \cap U = \emptyset$, and $\text{Area}(\Sigma) \leq C$, then there exists C_1 depending only on $n, M, \|h\|_{C^3(M)}, C$, such that*

$$|A^\Sigma|^2(x) \leq \frac{C_1}{\text{dist}_M^2(x, \partial U)} \quad \text{for all } x \in \Sigma.$$

Moreover if $\Sigma_k \subset U$ is a sequence of smooth, immersed (almost embedded when $n = 6$), two-sided, stable h -hypersurfaces in U with $\partial\Sigma_k \cap U = \emptyset$ and $\sup_k \text{Area}(\Sigma_k) < \infty$, then up to a subsequence, Σ_k converges locally smoothly (possibly with multiplicity) to some stable h -hypersurface Σ_∞ in U .

Proof. The compactness statement follows in the standard way from the curvature estimates. The curvature estimates follow from standard blowup arguments together with the Bernstein Theorem [55, Theorem 2] and [54, Theorem 3], the key being that the blowup will be a stable minimal hypersurface, and when $n = 6$, the blowup of a sequence of almost embedded h -hypersurfaces will be embedded by the classical maximum principle for embedded minimal hypersurfaces (c.f. [14]). \square

3.2. Some hypotheses on the zero set

We consider smooth prescription functions $h : M^{n+1} \rightarrow \mathbb{R}$ satisfying the following property:

- (†) If $\Sigma^n \subset M^{n+1}$ is a smoothly embedded hypersurface and $h|_\Sigma$ vanishes to all orders at $p \in \Sigma$, then there exists $r > 0$ for which $\Sigma_0 = \{h = 0\} \cap B_r(p)$ is a connected, smoothly embedded hypersurface passing through p . Moreover, if the mean curvature of Σ_0 vanishes to infinite order at any point, then it vanishes identically.

We can show that functions satisfying (†) are generic:

Proposition 3.7. *Let (M, g) be a smooth closed Riemannian manifold. Consider the set \mathcal{S} of smooth Morse functions h such that the zero set $\{h = 0\} =: \Sigma_0$ is a smooth closed hypersurface, and the mean curvature of Σ_0 vanishes to at most finite order.*

Then \mathcal{S} is open and dense in $C^\infty(M)$. Moreover each function in \mathcal{S} satisfies (†).

Proof. Recall that the set of Morse functions is open and dense in $C^\infty(M)$. In fact, since nondegenerate critical points are isolated, the singular set $\{h = \nabla h = 0\}$ of any Morse function h has only finitely many points. It follows that the set \mathcal{S}_0 of Morse functions h with empty singular set $\{h = \nabla h = 0\} = \emptyset$ is also open and dense. By the implicit function theorem, having empty singular set is equivalent to the condition that $\Sigma_0 = \{h = 0\}$ is a smooth, embedded, closed hypersurface.

First we address the openness of \mathcal{S} . Take $h \in \mathcal{S}$. Near Σ_0 , the gradient is bounded below, $|\nabla h| \geq \delta > 0$, so any small perturbation of h will have zero set given by a smooth hypersurface close to Σ_0 . Moreover, since the mean curvature of the level sets of h are given by $H = \operatorname{div} \left(\frac{\nabla h}{|\nabla h|} \right)$, any bound for the order of vanishing of H will be preserved under smooth perturbation.

To show that \mathcal{S} is dense, it suffices to show, given $h \in \mathcal{S}_0$, that we can construct a smooth perturbation of h whose zero set does not have mean curvature vanishing to infinite order. First, it is clear that we may perturb h so that no component of Σ_0 has identically vanishing mean curvature, so we assume this without loss of generality. The idea is then to run the mean curvature flow $\Sigma_{0,t}$ starting from the zero set Σ_0 , and construct a smooth deformation h^t of h which has the flowing hypersurface $\Sigma_{0,t}$ as its zero set.

One way to perform this procedure is to extend the mean curvature vector of $\Sigma_{0,t}$, for a short time interval $[0, t_0]$, to a smooth time-dependent vector field X_t supported in a fixed neighbourhood of Σ_0 . We may then take h^t to be the pullback of h under the flow of X_t . Alternatively, we can get somewhat more control by running mean curvature flow on all nearby level sets, and interpolating as follows:

Denote $\mathcal{N}_\epsilon = \{|h| < \epsilon\}$. Since $|\nabla h|$ is nonzero on Σ_0 , for sufficiently small $|s| < \epsilon$ the level sets $\Sigma_s = \{h = s\}$ are smooth, closed, embedded,

hypersurfaces. Fix a smooth function $0 \leq \phi \leq 1$ that is equal to 1 outside $\mathcal{N}_{\epsilon/3}$, and equal to 0 inside $\mathcal{N}_{\epsilon/4}$.

We claim that for a short time interval $[0, t_0]$, the level set flow starting from h yields a smooth family of smooth functions \tilde{h}^t , whose level sets $\{\tilde{h}^t = s\}$ coincide on $\mathcal{N}_{\epsilon/2}$ with the classical mean curvature flows $\Sigma_{s,t}$ starting from Σ_s . For small enough ϵ and t_0 , we also have $\{|\tilde{h}^t| < \epsilon/5\} \subset \mathcal{N}_{\epsilon/4}$; in particular h and \tilde{h}^t have the same sign at each point of $\mathcal{N}_{\epsilon/3} \setminus \mathcal{N}_{\epsilon/4}$. We defer the proof to Section 3.3. With the functions \tilde{h}^t in hand we may define $h^t = \phi h + (1 - \phi)\tilde{h}^t$.

By either construction, we have a smooth family of smooth Morse functions h^t , with zero set $\{h^t = 0\} = \Sigma_{0,t}$ as desired. We claim that at positive times the mean curvature of a hypersurface flowing by mean curvature cannot vanish to infinite order, unless it was a minimal hypersurface to begin with. Again the proof is deferred to Section 3.3. But we already ensured that no component of Σ_0 was minimal, so by the claim, the mean curvature of $\Sigma_{0,t}$ can only vanish to at most finite order as desired.

Finally, if $h \in \mathcal{S}$ and p, Σ are as in (\dagger) , then obviously $p \in \Sigma_0$. In fact, we can also show that Σ is tangent to Σ_0 at p : Otherwise, the vanishing of $h|_{\Sigma}$ to all orders would force p to be a degenerate critical point of h on M . \square

We can also handle smooth prescription functions with small zero set. In particular we can consider smooth functions $h : M^{n+1} \rightarrow \mathbb{R}$ satisfying the property:

- (\ddagger) The zero set $\{h = 0\}$ is contained in a countable union of connected, smoothly embedded $(n - 1)$ -dimensional submanifolds.

3.3. Some results on mean curvature flow

Here we prove two results on mean curvature flow that were needed to show that the set \mathcal{S} was a generic set. First we state a smooth short-time existence result for level set flows when everything is uniformly smooth.

Proposition 3.8. *Suppose that $h : M \rightarrow \mathbb{R}$ is a smooth function and that $0 < \delta \leq |\nabla h| \leq \delta^{-1}$ on $\mathcal{N}_{\epsilon} = \{|h| < \epsilon\}$. Further suppose that the level sets $\Sigma_s = \{h = s\}$ are smooth, closed, embedded hypersurfaces, which then have uniformly bounded curvature. Then there exists $t_0 > 0$ and a smooth family of smooth functions $\tilde{h}^t : \mathcal{N}_{\epsilon/2} \rightarrow \mathbb{R}$, $t \in [0, t_0]$, such that for each fixed $|s| \leq \epsilon/2$, the level sets $\{\tilde{h}^t = s\}$ are given by $\Sigma_{s,t} \cap \mathcal{N}_{\epsilon/2}$, where $\{\Sigma_{s,t} : t \in [0, t_0]\}$ is the classical mean curvature flow of Σ_s .*

Proof. First, it follows from Evans-Spruck [20] that there is a unique weak solution \tilde{h}^t of the level set flow with initial data h . By short-time existence and continuous dependence on the smooth initial data for the mean curvature flow, there exists a short time interval $t_0 > 0$ such that for any $|s| \leq \epsilon$, the mean curvature flow $\{\Sigma_{s,t}\}$ starting from Σ_s exists and is smooth for $t \in [0, t_0]$.

Therefore by [20] again, the level sets $\{\tilde{h}^t = s\}$ must coincide with the classical flows $\Sigma_{s,t}$. In particular, for $t \in [0, t_0]$, the mean curvature flow defines a smooth family of smooth, surjective maps $F^t : \mathcal{N}_\epsilon \rightarrow \mathcal{N}_\epsilon^t$, where $\mathcal{N}_\epsilon^t := \{|\tilde{h}^t| < \epsilon\}$. We claim that the $\{F^t\}_{t \in [0, t_0]}$ are in fact a smooth family of smooth *diffeomorphisms*.

By the avoidance principle for mean curvature flow, each F^t is certainly injective. In fact, by [20] once more, the distance between distinct level sets is bounded by $d(\Sigma_{s_1,t}, \Sigma_{s_2,t}) \geq d(\Sigma_{s_1}, \Sigma_{s_2})$. This implies that dF^t is uniformly nonsingular and the inverse function theorem then implies the claim.

By taking t_0 smaller if needed we can ensure that $\mathcal{N}_\epsilon^t \supset \mathcal{N}_{\epsilon/2}$. Then \tilde{h}^t is well-defined on $\mathcal{N}_{\epsilon/2}$ and satisfies $\tilde{h}^t = h \circ (F^t)^{-1}$, which completes the proof. \square

Second, we establish that at positive times, the mean curvature on a hypersurface flowing by mean curvature cannot vanish to infinite order unless it is minimal.

Proposition 3.9. *Let $\{\Sigma_t^n\}_{0 \leq t \leq t_0}$ be a smooth mean curvature flow of closed connected hypersurfaces in M^{n+1} . If H_{Σ_t} vanishes to all orders at $p \in \Sigma_t$ for some $t > 0$, then Σ_0 must have been a minimal hypersurface.*

Proof. For hypersurfaces under mean curvature flow, the mean curvature satisfies the parabolic PDE $(\partial_t - \Delta_g)H = (|A|^2 + \text{Ric}(\nu, \nu))H$, where Δ_g is the hypersurface Laplacian at time t . Since the flow is smooth on $[0, t_0]$, we have $|A|^2 + |\text{Ric}(\nu, \nu)| \leq C < \infty$. The conclusion follows from the spacelike strong unique continuation principle for parabolic PDE (see for instance [62, Corollary 4.2.7] or [21, 2]) and backwards uniqueness for mean curvature flow [39, 41]. \square

3.4. Estimates for the touching set

Let $h : M^{n+1} \rightarrow \mathbb{R}$ be a smooth function. Suppose that Σ_1^n, Σ_2^n are connected smoothly embedded hypersurfaces of prescribed mean curvature h in a connected open subset $U \subset M$, which lie to one side of one another. We would like to show that for certain classes of prescription functions h , it follows that either $\Sigma_1 = \Sigma_2$ or the touching set $\Sigma_1 \cap \Sigma_2$ is $(n-1)$ -rectifiable.

This section is mainly devoted to proving the following theorem (see also Remark 3.18 for the real analytic case):

Theorem 3.10. *Suppose that $h : M^{n+1} \rightarrow \mathbb{R}$ satisfies either property (\dagger) or (\ddagger) .*

Let Σ_1, Σ_2 be connected smoothly embedded hypersurfaces of prescribed mean curvature h in a connected open subset $U \subset M$, which lie to one side of one another. Then either $\Sigma_1 = \Sigma_2$, or the touching set $\Sigma_1 \cap \Sigma_2$ is contained in a countable union of connected, smoothly embedded $(n-1)$ -dimensional submanifolds.

Proof. By a covering argument, it suffices to prove that the assertion holds on a neighbourhood of each touching point p . So let $p \in \Sigma_1 \cap \Sigma_2$, then we may take a chart identifying $(\tilde{B}_r(p), g) \simeq (B_r^{n+1}(0), g')$ for which $T_p \Sigma_1 = T_p \Sigma_2$ corresponds to the plane $P = \{x_{n+1} = 0\}$, $\nu_1(p)$ corresponds to e_{n+1} and the hypersurfaces Σ_i may be written as graphs $x_{n+1} = u_i(x)$ of smooth functions u_i over the plane P .

The functions u_i then satisfy the quasilinear elliptic PDE

$$(3.1) \quad \begin{aligned} Hu_1 &= h(x, u_1(x)), \\ Hu_2 &= \begin{cases} h(x, u_2(x)), & \text{if } \nu_2(p) = \nu_1(p) \\ -h(x, u_2(x)), & \text{if } \nu_2(p) = -\nu_1(p). \end{cases} \end{aligned}$$

Here $H = Q + H_0$ is the mean curvature operator, and the zero order term H_0 is the mean curvature of P in $(B_r(0), g')$.

We now divide into cases:

3.4.1. Same orientation. If $\nu_1(p) = \nu_2(p)$, then the u_i satisfy the same quasilinear elliptic PDE $Hu = h(x, u(x))$. Since h is smooth, $u_1 - u_2$ has a sign, and $u_1(0) = u_2(0)$, the strong maximum principle of Serrin [56] implies that $u_1 \equiv u_2$. The standard connectedness argument for unique continuation then yields:

Lemma 3.11. *Let $h : M^{n+1} \rightarrow \mathbb{R}$ be a smooth function. Suppose that Σ_1, Σ_2 are connected smoothly embedded hypersurfaces of prescribed mean curvature h in a connected open subset $U \subset M$, which lie to one side of one another. Suppose that $p \in \Sigma_1 \cap \Sigma_2$ and $\nu_1(p) = \nu_2(p)$. Then $\Sigma_1 \equiv \Sigma_2$.*

3.4.2. Minimal sheets. It will be useful to record what occurs if one sheet, say Σ_2 , is in fact a minimal hypersurface, so that $h|_{\Sigma_2} = 0$. In this case, irrespective of orientation, u_2 again satisfies $Hu_2 = \pm 0 = h(x, u_2(x))$. Applying the strong maximum principle of Serrin [56] again we have:

Lemma 3.12. *Let $h : M^{n+1} \rightarrow \mathbb{R}$ be a smooth function. Suppose that Σ_1, Σ_2 are connected smoothly embedded hypersurfaces of prescribed mean curvature h in a connected open subset $U \subset M$, which lie to one side of one another. Suppose that $p \in \Sigma_1 \cap \Sigma_2$ and Σ_2 is minimal. Then $\Sigma_1 \equiv \Sigma_2$.*

3.4.3. Opposite orientation. In the opposite orientation case, we do not need the assumption that Σ_1, Σ_2 lie to one side of one another but just assume they are tangent. In this setting, we define the touching set to be $\{p \in \Sigma_1 \cap \Sigma_2 | T_p \Sigma_1 = T_p \Sigma_2\}$ and we will estimate it directly.

If $\nu_1(p) = -\nu_2(p)$, then the difference $v := u_1 - u_2$ satisfies an inhomogeneous linear elliptic PDE of the form

$$(3.2) \quad Lv = h(x, u_1(x)) + h(x, u_2(x)).$$

If h does not vanish at p , then $Lv(p) \neq 0$, so the Hessian of v at p has rank at least 1. The implicit function theorem then implies that, on a possibly smaller neighbourhood of p , the touching set $\{v = Dv = 0\}$ is contained in an $(n-1)$ -dimensional submanifold; (for more details one may consult [68, Lemma 2.8]). Thus we have shown:

Lemma 3.13. *Let $h : M^{n+1} \rightarrow \mathbb{R}$ be a smooth function. Suppose that Σ_1, Σ_2 are connected smoothly embedded hypersurfaces of prescribed mean curvature h in a connected open subset $U \subset M$. Suppose that $p \in \Sigma_1 \cap \Sigma_2$ and $\nu_1(p) = -\nu_2(p)$. Then $\Sigma_1 \cap \Sigma_2 \setminus \{h = 0\}$ is contained in a countable union of connected, smoothly embedded $(n-1)$ -dimensional submanifolds.*

It remains to estimate the stationary touching set $\Sigma_1 \cap \Sigma_2 \cap \{h = 0\}$, so in the remainder of this section we assume $h(p) = 0$. Of course, if h satisfies (\ddagger) , that is, if $\{h = 0\}$ is already contained in a countable union of connected, smoothly embedded $(n-1)$ -dimensional submanifolds, then we are already done.

First suppose that Σ_1, Σ_2 have a finite-order touching at p , that is, if $v = u_1 - u_2$ vanishes to finite order at $x = 0$. Then by the work of Hardt-Simon [33] the touching set $\{v = Dv = 0\}$ is in fact, again on a possibly smaller neighbourhood of p , contained in a countable union of $(n-2)$ -dimensional submanifolds.

It is for the remaining case of infinite-order touching that we require property (\dagger) : Suppose that $v = u_1 - u_2$ vanishes to infinite order at 0. Then by differentiating (3.2) we see that $h(x, u_i(x))$ vanishes to infinite order at $x = 0$, that is, $h|_{\Sigma_i}$ must vanish to infinite order at p .

Therefore by property (\dagger) , $\{h = 0\}$ is given by a smooth hypersurface Σ_0 near p . If it intersects Σ_i transversally at p , then $\Sigma_1 \cap \Sigma_2 \cap \{h = 0\}$ is

clearly contained in an $(n-1)$ -dimensional submanifold. So we may assume that Σ_0 is tangent to Σ_1 and Σ_2 at p .

Choose a ball $\tilde{B}_r(p)$ and a chart such that $\Sigma_0 \cap \tilde{B}_r(p)$ is identified with the set $P = \{x_{n+1} = 0\} \subset B_r^{n+1}(0)$. The smooth functions u_i then describe the height of Σ_i above Σ_0 . In particular, inside $\tilde{B}_r(p)$ we have

$$\{u_1 = u_2\} \cap \Sigma_0 \subset \{u_1 = u_2 = h = 0\},$$

and the u_i satisfy (3.1) with zero order term H_{Σ_0} .

If either u_i vanishes to finite order at p , then by a result of Bär [8] (see also [10]) the zero set of u_i is locally contained in the union of countably many $(n-1)$ -dimensional submanifolds.

Therefore the only remaining case is when both u_i vanish to infinite order at p . In this case, at p the Σ_i agree with Σ_0 to infinite order, and in particular the mean curvature H_{Σ_0} vanishes to infinite order (since $H_{\Sigma_i} = h|_{\Sigma_i}$ vanish to infinite order at p). By property (\dagger) , this means Σ_0 is minimal and in particular $H_{\Sigma_0} \equiv 0$ on a possibly smaller ball.

To proceed it is useful to record the following lemma which handles the case where either graph separately vanishes to infinite order:

Lemma 3.14. *Let $h : (B_r^{n+1}, g) \rightarrow \mathbb{R}$ be a smooth function satisfying $\Sigma_0 := \{x_{n+1} = 0\} \subset \{h = 0\}$, and that Σ_0 is minimal with respect to g . Suppose that Σ is a hypersurface in B_r of prescribed mean curvature h and that Σ is given by the graph of u over Σ_0 . If u vanishes to infinite order at $x = 0$, then there exists $\delta > 0$ such that $u \equiv 0$ on $|x| < \delta$; that is, Σ coincides with Σ_0 on B_δ .*

Proof. Since u vanishes to infinite order at $x = 0$, the mean curvature operator $Qu = H_\Sigma - H_{\Sigma_0}$ will be a perturbation of the Laplacian near 0. On the other hand, by supposition we have $H_\Sigma - H_{\Sigma_0} = h|_\Sigma$. So using that h is Lipschitz, for sufficiently small $\delta > 0$ we will have $|L_0 u| \leq C|u| + \epsilon|Du|$ for $|x| < \delta$, where $L_0 = a^{ij}D_{ij}$ is a uniformly elliptic operator with smooth coefficients $|a^{ij} - \delta^{ij}| < \epsilon \ll 1$. Strong unique continuation for elliptic differential inequalities (see for instance Aronszajn [7]) then implies that $u \equiv 0$ for $|x| < \delta$. \square

We pause to record the following corollary:

Corollary 3.15 (Unique minimal continuation). *Suppose that $h : M^{n+1} \rightarrow \mathbb{R}$ satisfies property (\dagger) . Let Σ be a connected smoothly embedded hypersurface of prescribed mean curvature h in a connected open subset $U \subset M$. If h vanishes on an open subset of Σ then $h|_\Sigma \equiv 0$, that is, Σ is a minimal hypersurface in U .*

(The corresponding result for (\ddagger) is trivial as in this case h cannot vanish on an open subset of Σ .)

Since each u_i vanishes to infinite order at $x = 0$, applying the lemma for each i we have $\Sigma_i = \Sigma_0$ on a neighborhood of p . That is, near p , the hypersurfaces Σ_0 , Σ_1 and Σ_2 all coincide and are in particular minimal. Again the usual connectedness argument for unique continuation then shows that the Σ_i must all be the same minimal hypersurface in U .

This concludes the proof of Theorem 3.10. \square

For convenience we state the following obvious corollary of Theorem 3.10 for almost embedded hypersurfaces:

Proposition 3.16 (Touching sets for almost embedded h -hypersurfaces). *If the metric on U^{n+1} is smooth, and $h : U^{n+1} \rightarrow \mathbb{R}$ satisfies either property (\dagger) or (\ddagger) then for any almost embedded hypersurface $\Sigma^n \subset U$ of prescribed mean curvature h , the touching set $\mathcal{S}(\Sigma)$ is contained in a countable union of connected, embedded $(n-1)$ -dimensional submanifolds.*

In particular, the regular set $\mathcal{R}(\Sigma)$ is open and dense in Σ .

Note that in the same orientation case $p \in \Sigma_1 \cap \Sigma_2$ with $\nu_1(p) = \nu_2(p)$, the difference $v := u_1 - u_2$ satisfies

$$(3.3) \quad Lv = h(x, u_1(x)) - h(x, u_2(x)),$$

so in particular $|Lv| \leq C|u_1 - u_2| = C|v|$. Thus, even without assuming that Σ_1, Σ_2 lie to one side of one another (that is, without assuming a sign for v), our work above together with strong unique continuation for differential inequalities [7] gives the following unique continuation for PMC:

Corollary 3.17 (Unique continuation for almost embedded h -hypersurfaces). *Suppose that $h : M^{n+1} \rightarrow \mathbb{R}$ satisfies either property (\dagger) or (\ddagger) . Let Σ_1^n, Σ_2^n be two connected almost embedded hypersurfaces of prescribed mean curvature h in a connected open subset U . If Σ_1 and Σ_2 coincide on a nonempty open neighbourhood U' , then $\Sigma_1 = \Sigma_2$.*

Remark 3.18. Both Proposition 3.16 and Corollary 3.17 also hold under condition (RA).

Indeed, when the metric on M and the function h are real analytic, we have the stronger statement that the touching set is a finite union of real analytic subvarieties $\bigcup_{k=0}^{n-1} S^k$ of respective dimension k . This follows from [40, Theorem 5.2.3], since in this setting the operator L will have analytic coefficients, and hence the difference v is also real analytic.

3.5. Compactness of stable PMC hypersurfaces

We are now in a position to prove our main compactness theorem.

Theorem 3.19 (Compactness for almost embedded stable h -hypersurfaces).

Let $2 \leq n \leq 6$. Suppose $\Sigma_k \subset U$ is a sequence of smooth, almost embedded, two-sided, stable $h^{(k)}$ -hypersurfaces in U , with $\sup_k \text{Area}(\Sigma_k) < \infty$. Further suppose that the functions $h^{(k)}$ converge smoothly to a smooth function h satisfying any property (\dagger) , (\ddagger) or (RA) .

Then, up to a subsequence, $\{\Sigma_k\}$ converges locally smoothly (with multiplicity) to some almost embedded stable h -hypersurface Σ_∞ in U .

If additionally $\{\Sigma_k\}$ are all boundaries, then each connected component Σ_∞^i of Σ_∞ is either:

- 1. minimal and smoothly embedded, or*
- 2. not minimal, with density 1 along its regular part and 2 along the touching set $\mathcal{S}(\Sigma_\infty)$;*

moreover, unless Σ_∞^i is minimal with multiplicity $l \geq 2$, we have that Σ_∞^∞ is locally a boundary - that is, for any point $p \in \Sigma_\infty^i$ there is a neighborhood \tilde{B}_p of p and an $\Omega \in \mathcal{C}(M)$ so that

$$\Sigma_\infty \cap \tilde{B}_p = \Sigma_\infty^i \cap \tilde{B}_p = \partial\Omega \llcorner \tilde{B}_p.$$

On the other hand, if $h^{(k)} \rightarrow h = 0$, then up to a subsequence, $\{\Sigma_k\}$ converges locally smoothly (with multiplicity) to some smoothly embedded stable minimal hypersurface Σ_∞ in U .

Proof of Theorem 3.19. The locally smooth convergence results follow straightforwardly from Theorem 3.6 and the almost embedded assumption. One can also rule out the case when sheets of the same orientation come together but do not coincide, by using the maximum principle Lemma 3.11 and the classical maximum principle for embedded minimal hypersurfaces (c.f. [14]) respectively.

Now we consider the case where the Σ_k are all boundaries. Namely, denote $\Sigma_k = \partial\Omega_k$ for some $\Omega_k \in \mathcal{C}(U)$. By standard compactness [57, Theorem 6.3], a subsequence of the $\partial\Omega_k$ converges weakly as currents to some $\partial\Omega_\infty$ with $\Omega_\infty \in \mathcal{C}(U)$.

Take an arbitrary point $p \in \Sigma_\infty$. Suppose p has density $l \geq 1$. Since we know the limit is almost embedded, we have an ordered graphical decomposition of Σ_∞ in a neighborhood $\tilde{B}_p \subset U$ of p given by $\bigcup_{i=1}^l \Sigma_\infty^i$, where each sheet has outward unit normal ν_∞^i and all sheets touch at p . By the locally

smooth convergence of Σ_k to Σ_∞ , for k large enough $\Sigma_k \cap \tilde{B}_p$ also has an ordered graphical decomposition $\bigcup_{i=1}^{l_k} \Sigma_k^i$, where $\Sigma_k^i \rightarrow \Sigma_\infty^i$.

We claim that if there are two distinct sheets $\Sigma_\infty^i, \Sigma_\infty^j$ with the same orientation $\nu_\infty^i(p) = \nu_\infty^j(p)$, then there must be another sheet Σ_∞^m in between, $i < m < j$, with the opposite orientation; and hence, $\Sigma_\infty \cap \tilde{B}_p$ is a minimal hypersurface with multiplicity l .

Indeed, suppose for the sake of contradiction that Σ_∞ has two sheets with the same orientation and no sheet in between. Then the same is true for Σ_k for large enough k , and we may assume without loss of generality that $i = 1, j = 2$. Then Σ_k^1, Σ_k^2 have unit normals ν_k^1, ν_k^2 pointing in the same direction. But then $\tilde{B}_p \setminus (\Sigma_k^1 \cup \Sigma_k^2)$ has three connected components U_0, U_1, U_2 such that, counting orientation, $(\partial U_0) \llcorner \tilde{B}_p = \Sigma_k^1$, $(\partial U_1) \llcorner \tilde{B}_p = \Sigma_k^2 - \Sigma_k^1$, and $(\partial U_2) \llcorner \tilde{B}_p = -\Sigma_k^2$.

On the other hand, for each i the Constancy Theorem [57, Theorem 26.27] applied to $\Omega_k \llcorner U_i$ implies that $\Omega_k \llcorner U_i$ is identical to either \emptyset or U_i . That is, $\Omega_k \llcorner \tilde{B}_p = \sum_{i=0}^2 a_i U_i$, where each $a_i = 0, 1$. It is then easy to see that any choice of the a_i will contradict the fact that, counting orientation, $\partial(\Omega_k \llcorner \tilde{B}_p) \llcorner \tilde{B}_p = \Sigma_k \cap \tilde{B}_p = \Sigma_k^1 + \Sigma_k^2$.

Thus if there were the two sheets $\Sigma_\infty^i, \Sigma_\infty^j$ with the same orientation, then there must be another sheet Σ_∞^m in between with the opposite orientation. But since all sheets touch at p , by Lemma 3.11 the sheets $\Sigma_\infty^i, \Sigma_\infty^j$ must coincide, which forces Σ_∞^m to coincide as well.

But then $\Sigma_\infty^i = \Sigma_\infty^m$ has prescribed mean curvature h with respect to both orientations, so h must vanish and it must be a minimal hypersurface. Lemma 3.12 then implies that any other sheet touching at p must coincide with Σ_∞^i as claimed.

Of course, if $l > 2$ then there must be two sheets converging with the same orientation. Thus we have shown that either: p has density $l = 1$ and is a regular point; p has density $l = 2$ and is a touching point between two non-minimal sheets of opposite orientation; or p is a regular point on a minimal hypersurface of density $l \geq 2$.

In particular if Σ_∞^i is not minimal then it has density 1 on its regular part and density 2 on its touching set. Having density 1 on the regular part is enough to conclude that $\Sigma_\infty \cap \tilde{B}_p = \Sigma_\infty^i \cap \tilde{B}_p = \partial \Omega_\infty \llcorner \tilde{B}_p$ as currents. \square

4. The h -min-max construction

In this section, we present the setup of the min-max construction mainly following Pitts [51]. We also prove the existence of a non-trivial sweepout with positive \mathcal{A}^h -min-max value.

4.1. Homotopy sequences

We will introduce the min-max construction using the scheme developed by Almgren and Pitts [3, 4, 51]; see also [68, Section 3].

Definition 4.1. (cell complex).

1. Denote $I = [0, 1]$, $I_0 = \partial I = I \setminus (0, 1)$;
2. For $j \in \mathbb{N}$, $I(1, j)$ is the cell complex of I , whose 1-cells are all intervals of form $[\frac{i}{3^j}, \frac{i+1}{3^j}]$, and 0-cells are all points $[\frac{i}{3^j}]$;
3. For $p = 0, 1$, $\alpha \in I(1, j)$ is a p -cell if $\dim(\alpha) = p$. A 0-cell is also called a vertex;
4. $I(1, j)_p$ denotes the set of all p -cells in $I(1, j)$, and $I_0(1, j)_0$ denotes the set $\{[0], [1]\}$;
5. Given a 1-cell $\alpha \in I(1, j)_1$, and $k \in \mathbb{N}$, $\alpha(k)$ denotes the 1-dimensional sub-complex of $I(1, j+k)$ formed by all cells contained in α . For $q = 0, 1$, $\alpha(k)_q$ and $\alpha_0(k)_q$ denote respectively the set of all q -cells of $I(1, j+k)$ contained in α , or in the boundary of α ;
6. The boundary homeomorphism $\partial : I(1, j) \rightarrow I(1, j)$ is given by $\partial[a, b] = [b] - [a]$ if $[a, b] \in I(1, j)_1$, and $\partial[a] = 0$ if $[a] \in I(1, j)_0$;
7. The distance function $\mathbf{d} : I(1, j)_0 \times I(1, j)_0 \rightarrow \mathbb{N}$ is defined as $\mathbf{d}(x, y) = 3^j|x - y|$;
8. The map $\mathbf{n}(i, j) : I(1, i)_0 \rightarrow I(1, j)_0$ is defined as: $\mathbf{n}(i, j)(x) \in I(1, j)_0$ is the unique element of $I(1, j)_0$, such that $\mathbf{d}(x, \mathbf{n}(i, j)(x)) = \inf \{ \mathbf{d}(x, y) : y \in I(1, j)_0 \}$.

For a map to the space of Caccioppoli sets: $\phi : I(1, j)_0 \rightarrow \mathcal{C}(M)$, the *fineness* of ϕ (with respect to \mathbf{M}) is defined as:

$$(4.1) \quad \mathbf{f}(\phi) = \sup \left\{ \frac{\mathbf{M}(\partial\phi(x) - \partial\phi(y))}{\mathbf{d}(x, y)} : x, y \in I(1, j)_0, x \neq y \right\}.$$

Similarly we can define the fineness of ϕ with respect to the \mathcal{F} -norm and \mathbf{F} -metric. We use $\phi : I(1, j)_0 \rightarrow (\mathcal{C}(M), \{0\})$ to denote a map such that $\phi(I(1, j)_0) \subset \mathcal{C}(M)$ and $\partial\phi|_{I_0(1, j)_0} = 0$, i.e. $\phi([0]), \phi([1]) = \emptyset$ or M .

Definition 4.2. Given $\delta > 0$ and $\phi_i : I(1, k_i)_0 \rightarrow (\mathcal{C}(M), \{0\})$, $i = 0, 1$, we say ϕ_1 is 1-homotopic to ϕ_2 in $(\mathcal{C}(M), \{0\})$ with fineness δ , if $\exists k_3 \in \mathbb{N}$, $k_3 \geq \max\{k_1, k_2\}$, and

$$\psi : I(1, k_3)_0 \times I(1, k_3)_0 \rightarrow \mathcal{C}(M),$$

such that

- $\mathbf{f}(\psi) \leq \delta$;
- $\psi([i], x) = \phi_i(\mathbf{n}(k_3, k_i)(x))$, $i = 0, 1$;
- $\partial\psi(I(1, k_3)_0 \times I_0(1, k_3)_0) = 0$.

Definition 4.3. A $(1, \mathbf{M})$ -homotopy sequence of mappings into $(\mathcal{C}(M), \{0\})$ is a sequence of mappings $\{\phi_i\}_{i \in \mathbb{N}}$,

$$\phi_i : I(1, k_i)_0 \rightarrow (\mathcal{C}(M), \{0\}),$$

such that ϕ_i is 1-homotopic to ϕ_{i+1} in $(\mathcal{C}(M), \{0\})$ with fineness δ_i , and

- $\lim_{i \rightarrow \infty} \delta_i = 0$;
- $\sup_i \{\mathbf{M}(\partial\phi_i(x)) : x \in I(1, k_i)_0\} < +\infty$.

Remark 4.4. Note that the second condition implies that $\sup_i \{\mathcal{A}^h(\phi_i(x)) : x \in I(1, k_i)_0\} < +\infty$.

Definition 4.5. Given two $(1, \mathbf{M})$ -homotopy sequences of mappings $S_1 = \{\phi_i^1\}_{i \in \mathbb{N}}$ and $S_2 = \{\phi_i^2\}_{i \in \mathbb{N}}$ into $(\mathcal{C}(M), \{0\})$, S_1 is homotopic to S_2 if $\exists \{\delta_i\}_{i \in \mathbb{N}}$, such that

- ϕ_i^1 is 1-homotopic to ϕ_i^2 in $(\mathcal{C}(M), \{0\})$ with fineness δ_i ;
- $\lim_{i \rightarrow \infty} \delta_i = 0$.

It is easy to see that the relation “is homotopic to” is an equivalence relation on the space of $(1, \mathbf{M})$ -homotopy sequences of mappings into $(\mathcal{C}(M), \{0\})$. An equivalence class is a $(1, \mathbf{M})$ -homotopy class of mappings into $(\mathcal{C}(M), \{0\})$. Denote the set of all equivalence classes by $\pi_1^\#(\mathcal{C}(M, \mathbf{M}), \{0\})$.

4.2. Min-max construction

Definition 4.6 (Min-max definition). Given $\Pi \in \pi_1^\#(\mathcal{C}(M, \mathbf{M}), \{0\})$, define $\mathbf{L}^h : \Pi \rightarrow \mathbb{R}^+$ to be the function given by:

$$\mathbf{L}^h(S) = \mathbf{L}^h(\{\phi_i\}_{i \in \mathbb{N}}) = \limsup_{i \rightarrow \infty} \max \{\mathcal{A}^h(\phi_i(x)) : x \text{ lies in the domain of } \phi_i\}.$$

The \mathcal{A}^h -min-max value of Π is defined as

$$(4.2) \quad \mathbf{L}^h(\Pi) = \inf \{\mathbf{L}^h(S) : S \in \Pi\}.$$

A sequence $S = \{\phi_i\} \in \Pi$ is called a *critical sequence* if $\mathbf{L}^h(S) = \mathbf{L}^h(\Pi)$.

Given a critical sequence S , then $K(S) = \{V = \lim_{j \rightarrow \infty} |\partial \phi_{i_j}(x_j)| : x_j \text{ lies in the domain of } \phi_{i_j}\}$ is a compact subset of $\mathcal{V}_n(M^{n+1})$. The *critical set* of S is the subset $C(S) \subset K(S)$ defined by

$$(4.3) \quad C(S) = \{V = \lim_{j \rightarrow \infty} |\partial \phi_{i_j}(x_j)| : \text{with } \lim_{j \rightarrow \infty} \mathcal{A}^h(\phi_{i_j}(x_j)) = \mathbf{L}^h(S)\};$$

we call any such sequence $\{\partial \phi_{i_j}(x_j)\}$ as in (4.3) a *min-max sequence*.

Note that by [51, 4.1(4)], we immediately have:

Lemma 4.7. *Given any $\Pi \in \pi_1^\#(\mathcal{C}(M, \mathbf{M}), \{0\})$, there exists a critical sequence $S \in \Pi$.*

The main theorem of this paper is as follows:

Theorem 4.8. *Let $2 \leq n \leq 6$. Consider a smooth closed Riemannian manifold M^{n+1} and a smooth function h , which satisfies $\int_M h \geq 0$ as well as any property (\dagger) , (\ddagger) or (RA) . There exists $\Pi \in \pi_1^\#(\mathcal{C}(M, \mathbf{M}), \{0\})$ and a critical sequence $S \in \Pi$ such that:*

- $\mathbf{L}^h(\Pi) = \mathbf{L}^h(S) > 0$;
- *There exists an element V of $C(S)$ induced by a nontrivial, smooth, closed, almost embedded, hypersurface $\Sigma^n \subset M$ of prescribed mean curvature h . Moreover V has multiplicity one, except possibly on components of Σ on which h vanishes.*

Proof of Theorem 4.8. This follows from combining Theorem 4.9, Theorem 6.4 and Theorem 7.1. \square

4.3. Existence of nontrivial sweepouts

Theorem 4.9. *There exists $\Pi \in \pi_1^\#(\mathcal{C}(M, \mathbf{M}), \{0\})$, such that for any function $h : M \rightarrow \mathbb{R}$ with $\sup |h| = c < \infty$ and $\int_M h \geq 0$, we have $\mathbf{L}^h(\Pi) > 0$. Moreover, for any critical sequence $S \in \Pi$, the critical set $C(S)$ is non-empty.*

Remark 4.10. We first describe a heuristic argument using smooth sweepouts which will help to reveal the key idea. Let $C_0 > 0$ and $V_0 > 0$ to be the constants in Theorem 2.3, and fix $0 < V \leq V_0$ such that $V^{\frac{-1}{n+1}} > 2c/C_0$. Note that V only depends on c, C_0, V_0 . Then for any Ω with $\text{Vol}(\Omega) = V$, we have

$$(4.4) \quad \mathcal{A}^h(\Omega) \geq C_0 V^{\frac{n}{n+1}} - cV > cV > 0.$$

Now consider any smooth 1-parameter family $\{\Omega_x : x \in [0, 1]\}$ satisfying $\Omega_0 = \emptyset$ and $\Omega_1 = M$. Since $\{\Omega_x\}$ sweeps out M , there must exist some $x_0 \in (0, 1)$ such that $\text{Vol}(\Omega_{x_0}) = V$, whence $\max_{x \in [0, 1]} \mathcal{A}^h(\Omega_x) \geq cV > 0$. Since this holds for any sweepout we then have $\mathbf{L}^h(\Pi) \geq cV > 0$.

Proof of Theorem 4.9. Note that $|\int_{\Omega} h| \leq c \text{Vol}(\Omega)$. The proof for the existence of $\Pi \in \pi_1^{\sharp}(\mathcal{C}(M, \mathbf{M}), \{0\})$ with $\mathbf{L}^h(\Pi) > 0$ proceeds as in the CMC setting [68, Theorem 3.9].

The remaining assertion that the critical sets are nontrivial follows from the following lemma.

Lemma 4.11. *Given $\mathbf{L} > 0$, then*

$$\inf\{\mathbf{M}(\partial\Omega) : \Omega \in \mathcal{C}(M) \text{ and } \mathcal{A}^h(\Omega) \geq \mathbf{L}\} > 0.$$

Proof: Suppose this is not true, then there exists a sequence $\{\Omega_i\} \subset \mathcal{C}(M)$, with

$$(4.5) \quad \mathcal{A}^h(\Omega_i) = \mathbf{M}(\partial\Omega_i) - \int_{\Omega_i} h \geq \mathbf{L} > 0,$$

but

$$\lim_{i \rightarrow \infty} \mathbf{M}(\partial\Omega_i) = 0.$$

Up to a subsequence, we can assume that $\lim_{i \rightarrow \infty} \Omega_i = \Omega_{\infty}$ weakly in the sense of Caccioppoli sets for some $\Omega_{\infty} \in \mathcal{C}(M)$. In particular, the characteristic functions $\chi_{\Omega_i} \rightarrow \chi_{\Omega_{\infty}}$ in $L^1(M)$. By the lower semi-continuity, we have

$$\mathbf{M}(\partial\Omega_{\infty}) \leq \lim_{i \rightarrow \infty} \mathbf{M}(\partial\Omega_i) = 0.$$

By the Constancy Theorem, we must have $\Omega_{\infty} = \emptyset$ or $\Omega_{\infty} = M$.

Since we assumed that $\int_M h \geq 0$, either case then yields a contradiction upon taking $i \rightarrow \infty$ in (4.5). This completes the proof of the lemma and hence the proof of Theorem 4.9. \square

5. Tightening

In this section, we recall the tightening process, which we constructed to study the \mathcal{A}^c functional (defined like \mathcal{A}^h but with the constant function c) as part of our min-max theory for constant mean curvature surfaces [68, §4]. The same process can be applied to the \mathcal{A}^h functional, and we will

prove that after applying the tightening map to a critical sequence, every element in the critical set has uniformly bounded first variation. Note that the tightening process is adapted from those in [13, §4] and [51, §4.3].

5.1. Review of constructions in [68, §4]

We first recall several key ingredients obtained in [68, §4] for the tightening process. In this section, we always let $c \equiv \sup_M |h|$. Given $L > 0$, consider the set of varifolds in $\mathcal{V}_n(M)$ with $2L$ -bounded mass: $A^L = \{V \in \mathcal{V}_n(M) : \|V\|(M) \leq 2L\}$. Denote

$$A_\infty^c = \{V \in A^L : |\delta V(X)| \leq c \int |X| d\mu_V, \text{ for any } X \in \mathfrak{X}(M)\}.$$

Given $V \in A^L$, we denote

$$\gamma = \gamma(V) = \mathbf{F}(V, A_\infty^c).$$

Given $X \in \mathfrak{X}(M)$, we use $\Phi_X : \mathbb{R}^+ \times M \rightarrow M$ to denote the one parameter group of diffeomorphisms generated by X .

We have constructed in [68, §4]:

- (i) a map $X : A^L \rightarrow \mathfrak{X}(M)$, which is continuous with respect to the C^1 topology on $\mathfrak{X}(M)$;
- (ii) two continuous functions $g : \mathbb{R}^+ \rightarrow \mathbb{R}^+$ and $\rho : \mathbb{R}^+ \rightarrow \mathbb{R}^+$, such that $\rho(0) = 0$ and

$$\delta W(X(V)) + c \int |X(V)| d\mu_W \leq -g(\gamma(V)),$$

whenever $W \in A^L, \mathbf{F}(W, V) \leq \rho(\gamma(V))$;

- (iii) a continuous time function $T : [0, \infty) \rightarrow [0, \infty)$, such that

- $\lim_{t \rightarrow 0} T(t) = 0$, and $T(t) > 0$ if $t \neq 0$;
- For any $V \in A^L$, denote $\Phi_V = \Phi_{X(V)}$, and $V_t = (\Phi_V(t))_\# V$, then $\mathbf{F}(V_t, V) < \rho(\gamma)$ for all $0 \leq t \leq T(\gamma)$.

Our goal is to show that given $\Omega \in \mathcal{C}(M)$ with $|\partial\Omega| \in A^L$, we can deform Ω by $\Phi_{|\partial\Omega|}(t)$ to get a 1-parameter family $\Omega_t = \Phi_{|\partial\Omega|}(t)(\Omega) \in \mathcal{C}(M)$, so that the \mathcal{A}^h functional of Ω_t for some $t > 0$ can be deformed down by a fixed amount depending only on $\gamma(|\partial\Omega|) = \mathbf{F}(|\partial\Omega|, A_\infty^c)$.

In particular, given $V \in A^L \setminus A_\infty^c$, denote

$$\Psi_V(t, \cdot) = \Phi_V(T(\gamma)t, \cdot), \quad \text{for } t \in [0, 1],$$

and $L : \mathbb{R}^+ \rightarrow \mathbb{R}^+$, with $L(\gamma) = T(\gamma)g(\gamma)$; then $L(0) = 0$ and $L(\gamma) > 0$ if $\gamma > 0$. We can deform V through the family $\left\{ V_t = (\Psi_V(t))_\# V : t \in [0, 1] \right\} \subset U_{\rho(\gamma)}(V)$, so that

1. The map $(t, V) \rightarrow V_t$ is continuous under the \mathbf{F} -metric;
2. When $V = |\partial\Omega|$, $\Omega \in \mathcal{C}(M)$, $\gamma = \mathbf{F}(|\partial\Omega|, A_\infty^c) > 0$, we have by (1.2)

$$(5.1) \quad \begin{aligned} \mathcal{A}^h(\Omega_1) - \mathcal{A}^h(\Omega) &\leq \int_0^{T(\gamma)} [\delta \mathcal{A}^h|_{\Omega_t}](X(|\partial\Omega|)) dt \\ &\leq -T(\gamma)g(\gamma) = -L(\gamma) < 0. \end{aligned}$$

(Note that $\delta \mathcal{A}_\Omega^h(X) \leq \delta |\partial\Omega|(X) + c \int |X| d\mu_{\partial\Omega}$ by (1.2).)

Finally note that the flow $\Psi_V(t, \cdot)$ is generated by the vector field

$$(5.2) \quad \tilde{X}(V) = T(\gamma)X(V).$$

5.2. Deforming sweepouts by the tightening map

We now apply our tightening map to the critical sequence provided by Lemma 4.7. As in our min-max theory for the CMC setting, the conclusion is that varifolds in the critical set have c -bounded first variation, where now $c = \sup |h|$. Indeed, the proof proceeds essentially unchanged, with the \mathcal{A}^c functional replaced by \mathcal{A}^h .

Proposition 5.1 (Tightening). *Let $\Pi \in \pi_1^\sharp(\mathcal{C}(M, \mathbf{M}), \{0\})$, $\int h \geq 0$, and assume $\mathbf{L}^h(\Pi) > 0$. For any critical sequence S^* for Π , there exists another critical sequence S for Π such that $C(S) \subset C(S^*)$ and each $V \in C(S)$ has c -bounded first variation, $c = \sup |h|$.*

Proof. Take $S^* = \{\phi_i^*\}$, where $\phi_i^* : I(1, k_i)_0 \rightarrow (\mathcal{C}(M), \{0\})$, and ϕ_i^* is 1-homotopic to ϕ_{i+1}^* in $(\mathcal{C}(M), \{0\})$ with fineness $\delta_i \searrow 0$. Let $\Xi_i : I(1, k_i)_0 \times [0, 1] \rightarrow \mathcal{C}(M)$ be defined as

$$\Xi_i(x, t) = \Psi_{|\partial\phi_i^*(x)|}(t)(\phi_i^*(x)),$$

where $\Psi(\cdot)$ is the tightening map defined above.

Denote $\phi_i^t(\cdot) = \Xi_i(\cdot, t)$. First, we first recall the following fact, the proof of which follows by a standard argument using (5.1); (see [68, §4.4 Claim 1]):

Claim 1: if $\lim_{i \rightarrow \infty} \mathcal{A}^h(\phi_i^1(x_i)) = \mathbf{L}^h(\Pi)$, then (up to relabeling) any converging subsequences of $\{\partial\phi_i^1(x_i)\}$ converges (as varifolds) to a varifold in $C(S^)$ of c -bounded first variation.*

Proof of Claim 1: By (5.1),

$$(5.3) \quad \mathcal{A}^h(\phi_i^1(x_i)) - \mathcal{A}^h(\phi_i^*(x_i)) = -L(\gamma_i),$$

where $\gamma_i = \mathbf{F}(|\partial\phi_i^*(x_i)|, A_\infty^c)$. Therefore,

$$\mathbf{L}^h(\Pi) = \lim \mathcal{A}^h(\phi_i^1(x_i)) = \lim \mathcal{A}^h(\phi_i^*(x_i)) - L(\lim \gamma_i) \leq \mathbf{L}^h(\Pi) - L(\lim \gamma_i),$$

so actually we must have $\lim \gamma_i = 0$ and this implies that $\lim |\partial\phi_i^*(x_i)| \in A_\infty^c$. Moreover, by our construction of the tightening map (see property (iii) in §5.1), each $|\partial\phi_i^1(x_i)|$ had to be $\rho(\gamma_i)$ -close to $|\partial\phi_i^*(x_i)|$ under the \mathbf{F} -metric, therefore

$$\lim |\partial\phi_i^1(x_i)| = \lim |\partial\phi_i^*(x_i)| \in A_\infty^c \cap C(S^*),$$

and this finishes the proof of the claim.

Heuristically, one would like to set $\phi_i = \phi_i^1$ as the desired sequence, but since the isotopies $\Psi_{|\partial\phi_i^*(x)|}$ depend on x , the fineness of $\{\phi_i^1\}$ could be large even if $\mathbf{f}(\phi_i^*)$ is small. Thus we need to interpolate ϕ_i^1 to get the desired ϕ_i , but we need to make sure the values of ϕ_i after interpolation are \mathbf{F} -close to those of ϕ_i^1 . The idea is to extend ϕ_i^1 to a *piecewise* continuous (with respect to the \mathbf{F} -metric) map on I and then apply the discretization result in [66, Theorem 5.1].

Similar difficulties appeared in the same way in [45, §15]. Instead, we use another interpolation method in Claim 2 as developed in [68]. For completeness, we provide detailed proof in Appendix B².

Claim 2: there exist integers $l_i > k_i$ and maps $\phi_i : I(1, l_i)_0 \rightarrow (C(M), \{0\})$ for each i , such that $S = \{\phi_i\}$ is homotopic to S^ , and*

- (a) $\phi_i^1 = \phi_i \circ \mathbf{n}(l_i, k_i)$ on $I(1, k_i)_0$;
- (b) $\mathbf{f}(\phi_i) \rightarrow 0$, as $i \rightarrow \infty$;

²We note that an alternative interpolation argument analogous to [45, §14] was found by the first author in [67, §1.3] after this article first appeared on arXiv.

- (c) $\mathcal{A}^h(\phi_i(x)) - \max\{\mathcal{A}^h(\phi_i^1(y)) : \alpha \in I(1, k_i)_1, x, y \in \alpha\} \rightarrow 0$, uniformly in $x \in I(1, l_i)_0$ as $i \rightarrow \infty$.
- (d) $\max\{\mathbf{F}(\partial\phi_i(x), \partial\phi_i^1(y)) : \alpha \in I(1, k_i)_1, x, y \in \alpha\} \rightarrow 0$, as $i \rightarrow \infty$.

In particular, S is a valid sequence in Π , and we now check that it satisfies the requirements of the proposition. First, property (c) and the fact that S^* is a critical sequence directly imply that S is also a critical sequence. It remains to show that every element in $C(S)$ must lie in $C(S^*)$ and have c -bounded first variation. Given $V \in C(S)$, one can find a subsequence (without relabeling) $\{\phi_i(\bar{x}_i) : \bar{x}_i \in I(1, l_i)_0\} \subset \mathcal{C}(M)$, such that $V = \lim |\partial\phi_i(\bar{x}_i)|$ as varifolds, and

$$\lim \mathcal{A}^h(\phi_i(\bar{x}_i)) = \mathbf{L}^h(\Pi).$$

We will need to first consider $\phi_i(x_i) = \phi_i^1(x_i)$, where x_i is the nearest point to \bar{x}_i in $I(1, k_i)_0$. By (c) and (d), we have $\lim \mathcal{A}^h(\phi_i^1(x_i)) = \mathbf{L}^h(\Pi)$ and also $\lim |\partial\phi_i(\bar{x}_i)| = \lim |\partial\phi_i^1(x_i)|$ as varifolds. Then by Claim 1, we conclude that $V \in A_\infty^c \cap C(S^*)$. This completes the proof. \square

6. h -almost minimizing

In this section, we introduce the notion of h -almost minimizing varifolds, and show the existence of such a varifold from min-max construction. We also construct h -replacements for any h -almost minimizing varifold. Using the properties of these replacements, we show that all blowups of h -almost minimizing varifolds are regular. As an easy consequence, the tangent cones of such varifolds are always planar.

Definition 6.1 (h -almost minimizing varifolds). Let $\underline{\nu}$ be the \mathcal{F} or \mathbf{M} -norm, or the \mathbf{F} -metric. For any given $\epsilon, \delta > 0$ and an open subset $U \subset M$, we define $\mathcal{A}^h(U; \epsilon, \delta; \underline{\nu})$ to be the set of all $\Omega \in \mathcal{C}(M)$ such that if $\Omega = \Omega_0, \Omega_1, \Omega_2, \dots, \Omega_m \in \mathcal{C}(M)$ is a sequence with:

- (i) $\text{spt}(\Omega_i - \Omega) \subset U$;
- (ii) $\underline{\nu}(\partial\Omega_{i+1}, \partial\Omega_i) \leq \delta$;
- (iii) $\mathcal{A}^h(\Omega_i) \leq \mathcal{A}^h(\Omega) + \delta$, for $i = 1, \dots, m$,

then $\mathcal{A}^h(\Omega_m) \geq \mathcal{A}^h(\Omega) - \epsilon$.

We say that a varifold $V \in \mathcal{V}_n(M)$ is h -almost minimizing in U if there exist sequences $\epsilon_i \rightarrow 0$, $\delta_i \rightarrow 0$, and $\Omega_i \in \mathcal{A}^h(U; \epsilon_i, \delta_i; \mathcal{F})$, such that $\mathbf{F}(|\partial\Omega_i|, V) \leq \epsilon_i$.

A simple consequence of the definition is that h -almost minimizing implies bounded first variation:

Lemma 6.2. *Let $V \in \mathcal{V}_n(M)$ be h -almost minimizing in U , then V has c -bounded first variation in U , where $c = \sup |h|$.*

Following similar arguments to [51, 4.10], we can show that every sequence which has been pulled tight using Proposition 5.1 has a critical limit which is h -almost minimizing on small annuli:

Definition 6.3. A varifold $V \in \mathcal{V}_n(M)$ is said to be h -almost minimizing in small annuli if for each $p \in M$, there exists $r_{am}(p) > 0$ such that V is h -almost minimizing in $A_{s,r}(p) \cap M$ for all $0 < s < r \leq r_{am}(p)$, where $A_{s,r}(p) = B_r(p) \setminus B_s(p)$.

Theorem 6.4 (Existence of h -almost minimizing varifold).

Let $\Pi \in \pi_1^\sharp(\mathcal{C}(M, \mathbf{M}), \{0\})$, $\int h \geq 0$, $c = \sup |h|$ and assume that $\mathbf{L}^h(\Pi) > 0$. There exists a nontrivial $V \in \mathcal{V}_n(M)$, such that

- (i) $V \in C(S)$ for some critical sequence S of Π ;
- (ii) V has c -bounded first variation;
- (iii) V is h -almost minimizing in small annuli.

In fact, one may show that for any critical sequence S of Π , there exists a varifold $V \in C(S)$, so that for any small annulus A , there exists a min-max sequence $\{\Omega_i\}$ such that Ω_i are eventually in $\mathcal{A}^h(A; \epsilon_i, \delta_i; \mathbf{M})$ for $\epsilon_i, \delta_i \rightarrow 0$, and have the nontrivial varifold limit $|\partial\Omega_i| \rightarrow V$. The main idea is that if there is no such sequence, the procedures described in [51, 4.10] allows one to deform S homotopically to a new sequence \tilde{S} with $\mathbf{L}^h(\tilde{S}) < \mathbf{L}^h(S)$, which contradicts the criticality of S .

We omit the proof of Theorem 6.4 except to point out the following equivalence result among several almost minimizing concepts using the three different topologies. In particular, it implies that we can work with the \mathbf{M} -norm at the expense of shrinking the open subset $U \subset M$.

Proposition 6.5. *Given $V \in \mathcal{V}_n(M)$, the following statements satisfy (a) \implies (b) \implies (c) \implies (d):*

- (a) V is h -almost minimizing in U ;
- (b) For any $\epsilon > 0$, there exists $\delta > 0$ and $\Omega \in \mathcal{A}^h(U; \epsilon, \delta; \mathbf{F})$ such that $\mathbf{F}(V, |\partial\Omega|) < \epsilon$;
- (c) For any $\epsilon > 0$, there exists $\delta > 0$ and $\Omega \in \mathcal{A}^h(U; \epsilon, \delta; \mathbf{M})$ such that $\mathbf{F}(V, |\partial\Omega|) < \epsilon$;
- (d) V is h -almost minimizing in W for any relatively open subset $W \subset\subset U$.

Remark 6.6. The proof of Proposition 6.5 for the area functional was originally due to Pitts [51, Theorem 3.9]. In our context, we work with boundaries instead of general integral currents. Furthermore, in Definition 6.1(iii), we use the \mathcal{A}^h functional instead of the mass \mathbf{M} . The main point is to derive (c) \implies (d), which follows from a standard interpolation process; see Lemma A.1.

Now we formulate and solve a natural constrained minimization problem.

Lemma 6.7 (A constrained minimization problem). *Given $\epsilon, \delta > 0$, $U \subset M$ and any $\Omega \in \mathcal{A}^h(U; \epsilon, \delta; \mathcal{F})$, fix a compact subset $K \subset U$. Let \mathcal{C}_Ω be the set of all $\Lambda \in \mathcal{C}(M)$ such that there exists a sequence $\Omega = \Omega_0, \Omega_1, \dots, \Omega_m = \Lambda$ in $\mathcal{C}(M)$ satisfying:*

- (a) $\text{spt}(\Omega_i - \Omega) \subset K$;
- (b) $\mathcal{F}(\partial\Omega_i - \partial\Omega_{i+1}) \leq \delta$;
- (c) $\mathcal{A}^h(\Omega_i) \leq \mathcal{A}^h(\Omega) + \delta$, for $i = 1, \dots, m$.

Then there exists $\Omega^ \in \mathcal{C}(M)$ such that:*

- (i) $\Omega^* \in \mathcal{C}_\Omega$, and

$$\mathcal{A}^h(\Omega^*) = \inf\{\mathcal{A}^h(\Lambda) : \Lambda \in \mathcal{C}_\Omega\},$$

- (ii) Ω^* is locally \mathcal{A}^h -minimizing in $\text{int}(K)$,
- (iii) $\Omega^* \in \mathcal{A}^h(U; \epsilon, \delta; \mathcal{F})$.

Proof. Let us first describe the construction of Ω^* . Take any minimizing sequence $\{\Lambda_j\} \subset \mathcal{C}_\Omega$, i.e.

$$\lim_{j \rightarrow \infty} \mathcal{A}^h(\Lambda_j) = \inf\{\mathcal{A}^h(\Lambda) : \Lambda \in \mathcal{C}_\Omega\}.$$

Notice that $\text{spt}(\Lambda_j - \Omega) \subset K$ and $\mathcal{A}^h(\Lambda_j) \leq \mathcal{A}^h(\Omega) + \delta$ for all j . By standard compactness [57, Theorem 6.3], after passing to a subsequence, Λ_j converges weakly to some $\Omega^* \in \mathcal{C}(M)$ and $\text{spt}(\Omega^* - \Omega) \subset K$. Since Λ_j converges weakly to Ω^* , we have that $\mathcal{H}^n(\partial\Omega^*) \leq \lim_{j \rightarrow \infty} \mathcal{H}^n(\partial\Lambda_j)$ and $\int_{\Omega^*} h = \lim_{j \rightarrow \infty} \int_{\Lambda_j} h$. Therefore,

$$(6.1) \quad \mathcal{A}^h(\Omega^*) \leq \inf\{\mathcal{A}^h(\Lambda) : \Lambda \in \mathcal{C}_\Omega\}.$$

Now by taking the advantage of the discrete deformation sequence, properties (i-iii) follow in the same way as those in [68, Lemma 5.7], and we leave the details to readers. \square

Given an h -almost minimizing varifold V , we can construct our h -replacements by applying Lemma 6.7 to the approximating sequence as in Definition 6.1 and taking the limit:

Proposition 6.8 (Existence and properties of replacements). *Let $V \in \mathcal{V}_n(M)$ be h -almost minimizing in an open set $U \subset M$ and $K \subset U$ be a compact subset, then there exists $V^* \in \mathcal{V}_n(M)$, called an h -replacement of V in K such that, with $c = \sup |h|$,*

- (i) $V \llcorner (M \setminus K) = V^* \llcorner (M \setminus K)$;
- (ii) $-c \operatorname{Vol}(K) \leq \|V\|(M) - \|V^*\|(M) \leq c \operatorname{Vol}(K)$;
- (iii) V^* is h -almost minimizing in U ;
- (iv) $V^* = \lim_{i \rightarrow \infty} |\partial \Omega_i^*|$ as varifolds for some $\Omega_i^* \in \mathcal{C}(M)$ such that $\Omega_i^* \in \mathcal{A}^h(U; \epsilon_i, \delta_i; \mathcal{F})$ with $\epsilon_i, \delta_i \rightarrow 0$; furthermore Ω_i^* locally minimizes \mathcal{A}^h in $\operatorname{int}(K)$;
- (v) if V has c -bounded first variation in M , then so does V^* .

Lemma 6.9 (Regularity of h -replacement). *Let $2 \leq n \leq 6$. Suppose that h satisfies (\dagger) , (\ddagger) or (RA) . Under the same hypotheses as Proposition 6.8, if $\Sigma = \operatorname{spt} \|V^*\| \cap \operatorname{int}(K)$, then Σ is a smooth, almost embedded, stable h -hypersurface. Furthermore, $V^* \llcorner \operatorname{int}(K) = \sum_{i=1}^L m_i \Sigma_i$, where each component Σ_i satisfies either:*

1. Σ_i is not minimal, and $m_i = 1$ so the density of V^* is 1 along $\mathcal{R}(\Sigma_i)$ and 2 along $\mathcal{S}(\Sigma_i)$; or
2. Σ_i is minimal and smoothly embedded, but m is some natural number.

Finally, if $p \in \Sigma_i$ and $m_i = 1$ then Σ is locally a boundary in a neighborhood of p .

Proof. By the regularity for local minimizers of the \mathcal{A}^h functional (Theorem 2.2), we know that each $\partial \Omega_i^*$ is a smoothly embedded h -boundary in $\operatorname{int}(K)$ by Proposition 6.8(iv). Moreover, $\partial \Omega_i^*$ is stable in $\operatorname{int}(K)$ in the sense of Definition 3.1. If this were not true, one can deform Ω_i^* by ambient isotopies supported in $\operatorname{int}(K)$ such that the \mathcal{A}^h values are strictly decreasing; it is then easy to see this contradicts Lemma 6.7(i). The lemma then follows from the compactness Theorem 3.19. \square

By iterating Proposition 6.8, we see that blowups of an h -almost minimizing varifolds have the good replacement property of [13, 16]. This allows us to characterize certain blowups of the the h -min-max varifold. In particular the tangent cones are planar; see also Section 7. The detailed proofs follow similar to [68, Lemma 5.10, Proposition 5.11], and we omit them here.

Given $p \in M, r > 0$, let $\eta_{p,r} : \mathbb{R}^L \rightarrow \mathbb{R}^L$ be the dilation defined by $\eta_{p,r}(x) = \frac{x-p}{r}$.

Proposition 6.10. *Let $2 \leq n \leq 6$, and $V \in \mathcal{V}_n(M)$ be an h -almost minimizing varifold in U . Given a sequence $p_i \in U$ with $p_i \rightarrow p \in U$ and, a sequence $r_i > 0$ with $r_i \rightarrow 0$, let $\bar{V} = \lim(\eta_{p_i,r_i})_{\#} V$ be the varifold limit. Then \bar{V} is an integer multiple of some complete embedded minimal hypersurface Σ in $T_p M$, and moreover, Σ is proper.*

Proposition 6.11 (Tangent cones are planes). *Let $2 \leq n \leq 6$. Suppose $V \in \mathcal{V}_n(M)$ has c -bounded first variation in M and is h -almost minimizing in small annuli. Then V is integer rectifiable. Moreover, for any $C \in \text{VarTan}(V, p)$ with $p \in \text{spt } \|V\|$,*

$$(6.2) \quad C = \Theta^n(\|V\|, p) |S| \text{ for some } n\text{-plane } S \subset T_p M$$

where $\Theta^n(\|V\|, p) \in \mathbb{N}$.

7. Regularity for h -min-max varifold

In this section, we prove the regularity of our min-max varifolds. In particular we prove that every varifold which has c -bounded variation and is h -almost minimizing in small annuli is a smooth, closed, almost embedded, hypersurface of h -prescribed mean curvature whose non-minimal components have multiplicity one.

Theorem 7.1 (Main regularity). *Let $2 \leq n \leq 6$, and (M^{n+1}, g) be an $(n+1)$ -dimensional smooth, closed Riemannian manifold. Further let $h : M \rightarrow \mathbb{R}$ be a smooth function satisfying (\dagger) , (\ddagger) or (RA) , and set $c = \sup |h|$.*

If $V \in \mathcal{V}_n(M)$ is a varifold which

1. *has c -bounded first variation in M , and*
2. *is h -almost minimizing in small annuli,*

then V is induced by Σ , where Σ is a closed, almost embedded h -hypersurface (possibly disconnected). Moreover, each component $\Sigma^{(i)}$ of Σ satisfies either:

- (i) *$\Sigma^{(i)}$ is not minimal, the density of V is exactly 1 at the regular set $\mathcal{R}(\Sigma^{(i)})$ and 2 at the touching set $\mathcal{S}(\Sigma^{(i)})$; or*
- (ii) *$\Sigma^{(i)}$ is a smoothly embedded minimal hypersurface (in particular $h \equiv 0$ on $\Sigma^{(i)}$), and V has integer density m_i on $\Sigma^{(i)}$.*

Proof. The conclusion is purely local, so we only need to prove the regularity of V near an arbitrary point $p \in \text{spt } \|V\|$. Fix a $p \in \text{spt } \|V\|$, then there

exists $0 < r_0 < r_{am}(p)$ such that for any $r < r_0$, the mean curvature H of $\partial B_r(p) \cap M$ in M is greater than c . Here $r_{am}(p)$ is as in Definition 6.3.

In particular, if $r < r_0$ and $W \in \mathcal{V}_n(M)$ has c -bounded first variation in M and $W \neq 0$ in $B_r(p)$, then by a standard contradiction argument applying the maximum principle (Proposition 2.1) to the varifold $W \llcorner G_n(B_r(p))$, we have

$$(7.1) \quad \emptyset \neq \text{spt } \|W\| \cap \partial B_r(p) = \text{Clos}(\text{spt } \|W\| \setminus \text{Clos}(B_r(p))) \cap \partial B_r(p).$$

Note that in the second equality we need a localized version of Proposition 2.1 which holds true by the remark after [63, Theorem 2].

We will show that $V \llcorner B_{r_0}(p)$ is either an embedded minimal hypersurface (on which $h \equiv 0$) with integer multiplicity, or an almost embedded hypersurface of prescribed mean curvature h with density equal to 2 along its touching set.

The argument consists of five steps:

Step 1: *Constructing successive h -replacements V^* and V^{**} on two overlapping concentric annuli.*

Step 2: *Gluing the h -replacements smoothly (as immersed hypersurfaces) on the overlap.*

Step 3: *Extending the h -replacements to the point p to get a h -‘replacement’ \tilde{V} on the punctured ball.*

Step 4: *Showing that the singularity of \tilde{V} at p is removable, so that \tilde{V} is regular.*

Step 5: *V coincides with the almost embedded hypersurface \tilde{V} on a small neighborhood of p .*

We now proceed to the proof.

Step 1

We first describe the construction of h -replacements on overlapping annuli; a key property will be that the replacements are also boundaries in the chosen annulus (see Claim 1), at least near points of multiplicity one.

Fix $0 < s < t < r_0$. By the choice of r_0 , we can apply Proposition 6.8 to V to obtain an h -replacement V^* in $K = \text{Clos}(A_{s,t}(p) \cap M)$. By (7.1) and Lemma 6.9, the restriction

$$\Sigma_1 = \text{spt } \|V^*\| \llcorner (A_{s,t}(p) \cap M)$$

is a nontrivial, smooth, almost embedded, stable h -hypersurface with some unit normal ν_1 ; when the multiplicity is 1, Σ_1 is locally a boundary so we can choose ν_1 to be the outer normal.

By Proposition 3.16 (see also Remark 3.18), the touching set $\mathcal{S}(\Sigma_1)$ is contained in a countable union of $(n-1)$ -dimensional connected submanifolds $\bigcup S_1^{(k)}$. Since a countable union of sets of measure zero still has measure zero, it follows from Sard's theorem that we may choose $s_2 \in (s, t)$ such that $\partial(B_{s_2}(p) \cap M)$ intersects Σ_1 and all the $S_1^{(k)}$ transversally.

Given any $s_1 \in (0, s)$, by Proposition 6.8(iii), V^* is still h -almost-minimizing in small annuli, and we can apply Proposition 6.8 again to get an h -replacement V^{**} of V^* in $K = \text{Clos}(A_{s_1, s_2}(p) \cap M)$. By (7.1) and Lemma 6.9 again, the restriction

$$\Sigma_2 = \text{spt} \|V^{**}\| \llcorner (A_{s_1, s_2}(p) \cap M)$$

is also a nontrivial, smooth, almost embedded, stable h -hypersurface with some unit normal ν_2 , which points outward at multiplicity 1 points in $A_{s_1, s_2}(p)$. Note that by Proposition 6.8(v), both V^* and V^{**} have c -bounded first variation.

We first observe that the second h -replacement can be chosen to be locally given by a boundary near ‘multiplicity one’ points in $\partial B_{s_2}(p)$, and in particular near non-minimal components of Σ_2 .

*Claim 1: there exists a set $\Omega^{**} \in \mathcal{C}(M)$ satisfying the following: Suppose $q \in \text{spt}(V^{**}) \cap \partial B_{s_2}(p)$, and Σ_1, Σ_2 have multiplicity 1 in a neighborhood of q . Then there exists $\epsilon > 0$ so that*

- a) Σ_1 and Σ_2 are the boundaries of Ω^{**} in $A_{s_2, t}(p) \cap B_\epsilon(q)$ and $A_{s_1, s_2}(p) \cap B_\epsilon(q)$ respectively;
- b) ν_1, ν_2 coincide with the outer unit normal of Ω^{**} in $A_{s_2, t}(p) \cap B_\epsilon(q)$ and $A_{s_1, s_2}(p) \cap B_\epsilon(q)$ respectively;
- c) if $\|V^{**}\|(\partial B_{s_2}(p)) = 0$, then V^{**} is identical to $|\partial \Omega^{**}|$ in $B_\epsilon(q) \cap M$.

The claim follows by constructing the ϵ_i -replacements $\Omega_i^{**} \rightarrow V^{**}$ from corresponding replacements $\Omega_i^* \rightarrow V^*$ (in particular, with consistent orientations). We refer the reader to [68, Theorem 6.1, Claim 1] for the details.

Step 2

We now show that Σ_1 and Σ_2 glue smoothly (as immersed hypersurfaces) across $\partial(B_{s_2}(p) \cap M)$. Indeed, define the intersection set

$$(7.2) \quad \Gamma = \Sigma_1 \cap \partial(B_{s_2}(p) \cap M), \quad \mathcal{S}(\Gamma) = \Gamma \cap \mathcal{S}(\Sigma_1).$$

Then by transversality, Γ is an almost embedded hypersurface in $\partial(B_{s_2}(p) \cap M)$, and $\mathcal{S}(\Gamma)$ is its touching set. Notice that

$$(7.3) \quad \mathcal{S}(\Gamma) \text{ is closed, and } \mathcal{R}(\Gamma) = \Gamma \setminus \mathcal{S}(\Gamma) \text{ is open in } \Gamma.$$

It follows from the maximum principle that

$$\text{Clos}(\Sigma_2) \cap \partial(B_{s_2}(p) \cap M) \subset \Gamma.$$

Indeed, (7.1) implies that any $y \in \text{Clos}(\Sigma_2) \cap \partial(B_{s_2}(p) \cap M)$ is also a limit point of $\text{spt} \|V^{**}\|$ from the outer side of $\partial B_{s_2}(p)$, on which V^{**} coincides with Σ_1 . In fact, we also have

Claim 2: $\text{Clos}(\Sigma_2) \cap \partial(B_{s_2}(p) \cap M) = \Gamma$, and then Σ_1 glues together continuously with Σ_2 .

Proof of Claim 2: By Proposition 6.8(i), we have

$$(7.4) \quad V^* = V^{**} = \Sigma_1, \quad \text{in } A_{s_2, t}(p) \cap M.$$

Given any $x \in \Gamma$, using (7.4), Proposition 6.11 and the fact that Σ_1 meets $\partial(B_{s_2}(p) \cap M)$ transversally, we have

$$(7.5) \quad \text{VarTan}(V^{**}, x) \text{ is an integer multiple of } T_x \Sigma_1.$$

This implies that x is a limit point of $\text{spt} \|V^{**}\|$ from inside of $\partial B_{s_2}(p)$, and thus completes the proof of the claim. \square

Furthermore, we will show that Σ_1 glues with Σ_2 in C^1 , i.e. the tangent spaces of Σ_1 and Σ_2 agree along Γ , with matching normals.

First we have the following.

Claim 3: Fix $x \in \Gamma$, and denote the plane $P_x = T_x \Sigma_1$ (with multiplicity 1). Then for any sequence of $x_i \rightarrow x$ with $x_i \in \Gamma$ and $r_i \rightarrow 0$, up to a subsequence we have

$$\begin{aligned} & \lim_{i \rightarrow \infty} (\eta_{x_i, r_i})_{\sharp} V^{**} \\ &= \begin{cases} \Theta^n(\|V^*\|, x) P_x, & \text{if } x \in \mathcal{R}(\Gamma), \\ P_x + \tau_v P_x, & \text{if } x \in \mathcal{S}(\Gamma), \liminf_{i \rightarrow \infty} \text{dist}_{\mathbb{R}^L}(x_i, \mathcal{S}(\Gamma))/r_i = \infty, \\ 2P_x, & \text{if } x \in \mathcal{S}(\Gamma), \liminf_{i \rightarrow \infty} \text{dist}_{\mathbb{R}^L}(x_i, \mathcal{S}(\Gamma))/r_i < \infty, \end{cases} \end{aligned}$$

where τ_w denotes translation by a vector w , and $v \in (P_x)^\perp$ is a vector in $T_x M$ orthogonal to P_x (v may be ∞ , in which case $\tau_v P$ is understood to be the empty set).

The proof of Claim 3 follows from the half space theorem for minimal hypersurfaces [36, Theorem 3] and the classical maximum principle for minimal hypersurfaces, after using that V^* and V^{**} coincide on $A_{s_2, t}(p)$ to determine the blowup on a halfspace. We refer the reader to [68, Theorem 6.1, Claim 3(A)(B)] for the details.

Since $\{(\eta_{x_i, r_i})_\# V^{**} : i \in \mathbb{N}\}$ have uniformly bounded first variation, a standard argument using the monotonicity formula implies that, in the Hausdorff topology, either

$$(7.6) \quad \text{spt } \|(\eta_{x_i, r_i})_\# V^{**}\| \rightarrow \begin{cases} T_x \Sigma_1, \text{ or} \\ T_x \Sigma_1 + \tau_v T_x \Sigma_1, & v \in (T_x \Sigma_1)^\perp. \end{cases}$$

To show that Σ_1 and Σ_2 glue together along Γ in C^1 near q , we will need:

Claim 4: For each $x \in \Gamma$, we have

$$(7.7) \quad \lim_{z \rightarrow x, z \in \Sigma_2} [T_z \Sigma_2] = [T_x \Sigma_1],$$

where $[T_z \Sigma_2]$ and $[T_x \Sigma_1]$ denote the un-oriented tangent planes of Σ_2 and Σ_1 (without counting multiplicity) respectively; and the convergence is uniform in x on compact subsets of Γ .

Moreover, if $x \in \mathcal{R}(\Gamma)$ and x lies in a multiplicity 1 component of Σ_1 , then in fact $\nu_2(z) \rightarrow \nu_1(x)$.

Proof of Claim 4: The uniformity follows from the fact that $[T \Sigma_1]$ is continuous on Γ , so we only need to establish the convergence to $[T \Sigma_1]$.

So consider a sequence $z_i \in \Sigma_2$ converging to some $x \in \Gamma$. Take $x_i \in \Gamma$ to be the nearest point projection (in \mathbb{R}^L) of z_i to Γ and $r_i = |z_i - x_i|$. Note that $x_i \rightarrow x \in \Gamma$ and $r_i \rightarrow 0$, so we are in the situation of Claim 3. Note that $\Sigma_2 \cap B_{r_i/2}(z_i)$ is an almost embedded, stable h -hypersurface in M , so by Theorem 3.19 a subsequence of the blow-ups $\eta_{x_i, r_i}(\Sigma_2 \cap B_{r_i/2}(z_i))$ converges smoothly to a smooth, embedded, stable, minimal hypersurface Σ_∞ contained in a half-space of $T_x M$.

On the other hand, Claim 3 and (7.6) imply that $\eta_{x_i, r_i}(\Sigma_2 \cap B_{r_i/2}(z_i))$ converges in the Hausdorff topology to a domain in $T_x \Sigma_1$. Therefore, we have

$\Sigma_\infty \subset T_x \Sigma_1$. The smooth convergence then implies that $\nu_2(z_i)$ converges to one of the unit normals $\pm \nu_1(x)$ of $T_x \Sigma_1$.

Now suppose that $x \in \mathcal{R}(\Gamma)$ and x lies in a multiplicity 1 component of Σ_1 . Then the multiplicity of Σ_2 is also 1 near x , by the upper semi-continuity of density function for varifolds with bounded first variation [57, 17.8]. Then by (7.5), [57, Theorem 3.2(2)] and Claim 1(c), we have

$$(7.8) \quad \begin{aligned} & \|V^{**}\|(\partial B_{s_2}(p)) = 0, \\ & \text{and hence } V^{**} = |\partial \Omega^{**}| \text{ in } B_\epsilon(x) \cap M \text{ for some } \epsilon > 0. \end{aligned}$$

This implies that the limit of $\nu_2(z_i)$ must be $\nu_1(x)$ by Claim 1, so *Claim 4* is proved. \square

We first consider the gluing near a regular point $q \in \mathcal{R}(\Gamma)$. By Claim 4, Σ_1 and Σ_2 glue together along Γ as a C^1 hypersurface with matching unit normals and prescribed mean curvature h (in the weak sense). Note that if either Σ_i was a minimal hypersurface near q then we may have had to choose the opposite normal to ν_i to ensure the C^1 gluing. (This is allowed since in this situation h vanishes on Σ_i , so it still has prescribed mean curvature h with respect to the opposite normal.)

The higher regularity then follows from a standard elliptic PDE argument. More precisely, Σ_1 and Σ_2 can be written as graphs of some functions u_1, u_2 over $T_q \Sigma_1$ respectively. Since Σ_1 and Σ_2 both have prescribed mean curvature h with respect to unit normals pointing to the same side of $T_q \Sigma_1$, they satisfy the same mean curvature type elliptic PDE with inhomogeneous term given by $h|_{\Sigma_1 \cup \Sigma_2}$, which has the same regularity as the glued hypersurface $\Sigma_1 \cup \Sigma_2$. The higher regularity follows from the elliptic regularity of this PDE.

Thus we have proven that Σ_2 glues smoothly with $\Sigma_1 \cap A_{s_2,t}(p)$ along $\mathcal{R}(\Gamma)$. Moreover, by the unique continuation for elliptic PDE (for instance Corollary 3.17) we know that Σ_2 is identical to Σ_1 in a neighborhood of $\mathcal{R}(\Gamma)$ in $A_{s_2,t}(p) \cap M$. We now show that the smooth gluing extends to the touching set $\mathcal{S}(\Gamma)$.

Now consider an arbitrary fixed singular point $q \in \mathcal{S}(\Gamma)$. By Lemma 6.9, in some small neighborhood $U \subset M$ of q , $\Sigma_1 \cap U$ is the union of two connected, embedded h -hypersurfaces $\Sigma_{1,1} \cup \Sigma_{1,2}$ with unit normals $\nu_{1,1}$ and $\nu_{1,2}$, such that $\Sigma_{1,2}$ lies on one side of $\Sigma_{1,1}$ and they touch tangentially at $\mathcal{S}(\Sigma_1) \cap U = \Sigma_{1,1} \cap \Sigma_{1,2}$. By Lemma 3.11, $\nu_{1,1} = -\nu_{1,2}$ along the touching set $\mathcal{S}(\Sigma_1) \cap U$. Denote $\Gamma \cap \Sigma_{1,1} = \Gamma_1$ and $\Gamma \cap \Sigma_{1,2} = \Gamma_2$, then as embedded

submanifolds of $\partial(B_{s_2}(p) \cap U)$, Γ_2 lies on one-side of Γ_1 and they touch tangentially along $\mathcal{S}(\Gamma) \cap U$.

By Claim 4, using the regularity of Σ_2 in Lemma 6.9, near q the hypersurface Σ_2 can be written as a set of graphs $\{\Sigma_{2,i} : i = 1 \cdots l\}$ over the half space $[T_q(\Sigma_1 \cap B_{s_2}(p))]$. Now since Σ_2 glues smoothly with Σ_1 along $\mathcal{R}(\Gamma)$, and since $\mathcal{R}(\Gamma)$ is an open and dense subset of Γ , we know that the set $\{\Sigma_{2,i} : i = 1 \cdots l\}$ consists of exactly two elements: one of them, denoted by $\Sigma_{2,1}$, glues smoothly with $\Sigma_{1,1}$ along $\Gamma_1 \setminus \mathcal{S}(\Gamma)$; the other one, denoted by $\Sigma_{2,2}$, glues smoothly with $\Sigma_{1,2}$ along $\Gamma_2 \setminus \mathcal{S}(\Gamma)$. This, together with (7.7), implies that the pairs $(\Sigma_{1,1}, \Sigma_{2,1})$ and $(\Sigma_{1,2}, \Sigma_{2,2})$ glue together in C^1 near q respectively (with matching orientations). Again higher regularity follows from the elliptic PDE argument as for the regular part. This completes the smooth gluing near the touching set.

Step 3

We now wish to extend the replacements, via unique continuation, all the way to the center p .

Henceforth we denote V^{**} by $V_{s_1}^{**}$ and Σ_2 by Σ_{s_1} to indicate the dependence on s_1 . By varying $s_1 \in (0, s)$, we obtain a family of nontrivial, smooth, almost embedded, stable h -hypersurfaces $\{\Sigma_{s_1} \subset A_{s_1, s_2}(p) \cap M\}$. Since unique continuation holds for almost embedded prescribed mean curvature hypersurfaces (Corollary 3.17) by Step 2 we have $\Sigma_{s_1} = \Sigma_1$ in $A_{s, s_2}(p)$, and moreover, for any $s'_1 < s_1 < s$, we have $\Sigma_{s'_1} = \Sigma_{s_1}$ in $A_{s_1, s_2}(p)$. Hence

$$\Sigma := \bigcup_{0 < s_1 < s} \Sigma_{s_1}$$

is a nontrivial, smooth, almost embedded, stable h -hypersurface in $(B_{s_2}(p) \setminus \{p\}) \cap M$.

Let $\Sigma^{(i)}$, $i = 1, \cdots, l$ be the connected components of Σ . We also have

$$\text{spt} \|V_{s_1}^{**}\| = \Sigma, \text{ in } A_{s_1, s_2}(p), V_{s_1}^{**} = V^* \text{ in } A_{s, s_2}(p),$$

$$\text{and for any } s'_1 < s_1 < s, V_{s'_1}^{**} = V_{s_1}^{**} \text{ in } A_{s_1, s_2}(p).$$

Indeed, to show the matching between V^* , $V_{s_1}^{**}$, and $V_{s'_1}^{**}$, we only need to show the matching of densities. Even though there is no satisfied Constancy Theorem as the minimal case, the matching follows from (7.5) applied to their supports along $\partial B_s(p)$.

By Proposition 6.8, $V_{s_1}^{**}$ has c -bounded first variation and uniformly bounded mass for all $0 < s_1 < s$. Therefore as $s_1 \rightarrow 0$, the family $V_{s_1}^{**}$ will converge to a varifold $\tilde{V} \in \mathcal{V}_n(M)$ with c -bounded first variation, i.e.

$$(7.9) \quad \tilde{V} = \lim_{s_1 \rightarrow 0} V_{s_1}^{**}, \quad \text{such that}$$

$$\tilde{V} = \begin{cases} \sum_{i=1}^l m_i \Sigma^{(i)} & \text{in } (B_{s_2}(p) \setminus \{p\}) \cap M \\ V^* & \text{in } M \setminus B_s(p) \end{cases}, \quad \text{and } \|\tilde{V}\|(\{p\}) = 0,$$

where $m_i > 1$ only if $\Sigma^{(i)}$ is minimal. Since $p \in \text{spt } \|V_{s_1}^{**}\|$, by the upper semi-continuity of density function for varifolds with bounded first variation [57, 17.8], we know that $p \in \text{spt } \|\tilde{V}\|$.

Step 4

We now determine the regularity of \tilde{V} at p .

First, \tilde{V} has c -bounded first variation. Second, $\text{spt } \|\tilde{V}\|$, when restricted to any small annulus $A_{r,2r}(p) \cap M$, already coincides with a smooth, almost embedded, stable h -hypersurface Σ . Using these two ingredients, we can use a standard blowup argument (see for instance Proposition 6.11 or [68, Proposition 5.11]) to show that every tangent varifold of \tilde{V} at p is an integer multiple of some n -plane, i.e. for any $C \in \text{VarTan}(V, p)$,

$$C = \Theta^n(\|\tilde{V}\|, p) |S|, \quad \text{for some } n\text{-plane } S \subset T_p M \text{ where } \Theta^n(\|\tilde{V}\|, p) \in \mathbb{N}.$$

Now the removability of the singularity of \tilde{V} at p (as an almost embedded hypersurface) follows similarly to [51, Theorem 7.12]. We include the details for completeness. We can assume that

$$\Theta^n(\|\tilde{V}\|, p) = m$$

for some $m \in \mathbb{N}$. Since Σ is stable in a punctured ball of p , by Theorem 3.19, for any sequence $r_j \rightarrow 0$,

$$\eta_{p,r_j}(\Sigma) \rightarrow m \cdot S$$

(up to subsequences) locally smoothly in $\mathbb{R}^L \setminus \{0\}$ for some n -plane $S \subset T_p M$. However, S may depend on the sequence r_j . By the convergence and the regularity of Σ , there exists $\sigma_0 > 0$ small enough, such that for any $0 < \sigma \leq \sigma_0$, Σ has an ordered (in the sense of Definition 3.2) graphical

decomposition in $A_{\sigma/2,\sigma}(p)$:

$$(7.10) \quad \tilde{V} \llcorner A_{\sigma/2,\sigma}(p) = \sum_{i=1}^l m_i |\Sigma_i(\sigma)|, \quad \sum_{i=1}^l m_i = m$$

Here $\Sigma_i(\sigma)$ are distinct graphs over $A_{\sigma/2,\sigma}(p) \cap S$ for some n -plane $S \subset T_p M$, and $m_i > 1$ only if $\Sigma_i(\sigma)$ is minimal.

Since (7.10) holds for all σ , by continuity of Σ we can continue each $\Sigma_i(\sigma_0)$ to $(B_{\sigma_0}(p) \setminus \{p\}) \cap M$, and we denote this continuation by Σ_i . Since each piece Σ_i has prescribed mean curvature h , by a standard extension argument (c.f. the proof in [34, Theorem 4.1]), each Σ_i can be extended as a varifold with c -bounded first variation in $B_{\sigma_0}(p) \cap M$ (recall $c = \sup |h|$). Given $C_i \in \text{VarTan}(\Sigma_i, p)$, to see that C_i has multiplicity one, first notice that

$$(7.11) \quad \Theta^n(\|C_i\|, p) \geq 1,$$

since each Σ_i is h -stable and thus its re-scalings converge with multiplicity to a smooth, embedded, stable, minimal hypersurface by Theorem 3.19. If equality does not hold for some i in (7.11), this will derive a contradiction since

$$\tilde{V} \llcorner B_{\sigma_0}(p) = \sum_{i=1}^l m_i |\Sigma_i|, \quad \sum_{i=1}^l m_i = m$$

Therefore, each Σ_i has c -bounded first variation in $B_{\sigma_0}(p) \cap M$ and $\Theta^n(\|\Sigma_i\|, p) = 1$; by the Allard regularity theorem [57, Theorem 24.2] and elliptic regularity, Σ_i extends as a smooth, embedded h -hypersurface across p . Moreover, each minimal Σ_i extends as a smooth minimal hypersurface across p . Finally, by the maximum principle (Lemma 3.11, Lemma 3.12) either some sheet Σ_i is indeed minimal so all sheets must coincide and $\tilde{V} \llcorner B_{\sigma_0}(p) = m[\Sigma_i]$, or none are minimal and $m = 1$ or $m = 2$. This shows that \tilde{V} extends as an almost embedded h -hypersurface across p with the desired regularity.

Step 5

Finally, to complete the proof we show that V coincides with \tilde{V} on a small ball about p .

We will need the following simple corollary of the first variation formula; (see [68, Lemma 6.2]).

Lemma 7.2. *For small enough r the set*

$$\mathrm{Tr}_p^V = \left\{ y \in \mathrm{spt} \|V\| \cap (B_r(p) \setminus \{p\}) : \begin{array}{l} \mathrm{VarTan}(V, y) \text{ consists of an} \\ \text{integer multiple of an } n\text{-plane} \\ \text{transverse to } \partial(B_{\mathrm{dist}(y,p)}(p) \cap M) \end{array} \right\}$$

is a dense subset of $\mathrm{spt} \|V\| \cap B_r(p)$.

Claim 5: *For small enough r , $\mathrm{spt} \|V\| = \Sigma$ in the punctured ball $(B_r(p) \setminus \{p\}) \cap M$.*

Proof of Claim 5: We first prove that $\mathrm{Tr}_p^V \subset \Sigma$, which combined with Lemma 7.2 will imply that $\mathrm{spt} \|V\| \cap (B_r(p) \setminus \{p\}) \subset \Sigma$. Fix $y \in \mathrm{Tr}_p^V \cap (B_r(p) \setminus \{p\})$, and let $\rho = \mathrm{dist}(y, p)$. Consider V_ρ^{**} . By transversality we have $y \in \mathrm{Clos}(\mathrm{spt} \|V\| \cap B_\rho(p))$. On the other hand, $V_\rho^{**} = V^* = V$ inside $B_\rho(p)$, so by (7.1) we have

$$\begin{aligned} & \mathrm{Clos}(\mathrm{spt} \|V\| \cap B_\rho(p)) \cap \partial B_\rho(p) \\ &= \mathrm{Clos}(\mathrm{spt} \|V_\rho^{**}\| \cap B_\rho(p)) \cap \partial B_\rho(p) \\ &\subset \mathrm{Clos}(\mathrm{spt} \|V_\rho^{**}\| \setminus \mathrm{Clos}(B_\rho(p))) \cap \partial B_\rho(p). \end{aligned}$$

Since $\mathrm{spt} \|V_\rho^{**}\| = \Sigma$ on $A_{\rho, s_2}(p)$, we therefore have $y \in \Sigma$.

Next we show the reverse inclusion $\Sigma \subset \mathrm{spt} \|V\|$. Since Σ extends across p as an almost embedded hypersurface, we know that $T_y \Sigma$ is transverse to $\partial(B_{\mathrm{dist}(y,p)}(p) \cap M)$ for all $y \in \Sigma \cap B_r(p)$ for small enough r . Let ρ and V_ρ^{**} be as above, then $y \in \mathrm{spt} \|V_\rho^{**}\|$. By Proposition 6.11, $\mathrm{VarTan}(V_\rho^{**}, y) = \{\Theta^n(\|V_\rho^{**}\|, y) | T_y \Sigma\}$. By the transversality, we then have $y \in \mathrm{Clos}(\mathrm{spt} \|V_\rho^{**}\| \cap B_\rho(p))$, so since $V_\rho^{**} = V$ inside $B_\rho(p)$ we conclude that $y \in \mathrm{Clos}(\mathrm{spt} \|V\| \cap B_\rho(p)) \subset \mathrm{spt} \|V\|$ as desired. \square

Note that we do not have a suitable Constancy Theorem (c.f. [57, 41.1]) for varifolds with bounded first variation. In order to show that V coincides with Σ near p , our strategy is to show that $V = \tilde{V}$ as varifolds in a neighborhood of p . By the transversality argument as above, we only need to show that the densities of V and \tilde{V} are identical along $\Sigma \cap (B_r(p) \setminus \{p\})$, where r is chosen as in Claim 5.

Claim 6: $\Theta^n(\|V\|, \cdot) = \Theta^n(\|\tilde{V}\|, \cdot)$ on $\Sigma \cap B_r(p) \setminus \{p\}$.

Proof of Claim 6: Let $y \in \Sigma$ and $\rho = \mathrm{dist}(y, p) < r$ be as above. Then since $V_\rho^{**} = V$ inside $B_\rho(p)$, by transversality and Proposition 6.11 we have

$\text{VarTan}(V, y) = \text{VarTan}(V_\rho^{**}, y)$. But $V_\rho^{**} = \tilde{V}$ on $A_{\rho, s_2}(p)$, so we must have $\text{VarTan}(V_\rho^{**}, y) = \{\Theta^n(\|\tilde{V}\|, y) | T_y \Sigma | \}$. Thus $\Theta^n(\|V\|, y) = \Theta^n(\|\tilde{V}\|, y)$. \square

Combining Claims 5 and 6 yields that $V = \tilde{V}$ on $B_r(p) \cap M$. This finishes the proof of Step 5, and hence also completes the proof of the main Theorem 7.1. \square

Finally we have the following result which is implied by the above proof.

Proposition 7.3. *Let V be as in Theorem 7.1. Assume that $V = \lim_{i \rightarrow \infty} |\partial \Omega_i|$, where $\{\Omega_i\}$ are the approximating sets of the h -almost-minimizing varifold V , given by Definition 6.1. If there are no minimal sheets in $\Sigma = \text{spt } V$, then Σ is a boundary of some $\Omega \in \mathcal{C}(M)$ where its mean curvature with respect to the unit outer normal is h , and*

$$\mathcal{A}^h(\Omega) = \lim_{i \rightarrow \infty} \mathcal{A}^h(\Omega_i).$$

In particular, if $V \in C(S)$ for some critical sequence $S \in \Pi$ as given by Theorem 6.4 and the remarks following it, we have

$$\mathcal{A}^h(\Omega) = \mathbf{L}^h(\Pi).$$

Proof. We may assume without loss of generality that Σ is connected. It suffices to prove that $\partial \Omega_i$ sub-converges to Σ weakly as currents. Take Ω as the weak limit of Ω_i as Caccioppoli sets (up to a subsequence), then $\text{spt}(\partial \Omega) \subset \Sigma$, and $|\partial \Omega| = 0$ or Σ as varifolds by the Constancy Theorem (by taking Σ as the ambient space). Fix an arbitrary regular (non-touching) point p in Σ . As in Step 1 of the proof of Theorem 7.1, we can take a first replacement V^* near p . In fact, we showed that *a posteriori* V^* coincides with V .

The construction of V^* , however, came with the constrained minimizers Ω_i^* by Proposition 6.8. The Constancy Theorem still implies that Ω_i^* converges weakly to Ω , but now according to the proof of Lemma 6.9, the $\partial \Omega_i^*$ are all smoothly embedded h -hypersurfaces, converging smoothly to Σ in $\text{int}(K)$. Therefore $\partial \Omega_i^*$ must converge to Σ as currents in $\text{int}(K)$. This shows that $\partial \Omega = \Sigma$ inside $\text{int}(K)$ as currents, and hence they must coincide everywhere, which concludes the proof. \square

Appendix A. An interpolation lemma

The following interpolation lemma was proved in in [68, Appendix A] when h is a constant function. This type of result was essentially due to Pitts

[51, Lemma 3.8], but the modification to find the interpolation sequence using boundaries of Caccioppoli sets was completed by the first author [66, Proposition 5.3]. The extension to non-constant functions h proceeds similar to [68, Appendix A], the key being to control $|\int_{\Omega} h| \leq c \text{Vol}(\Omega)$, so we omit the details here.

Lemma A.1. *Suppose $L > 0$, $\eta > 0$, W is a compact subset of U , and $\Omega \in \mathcal{C}(M)$. Then there exists $\delta = \delta(L, \eta, U, W, \Omega) > 0$, such that for any $\Omega_1, \Omega_2 \in \mathcal{C}(M)$ satisfying*

- (a) $\text{spt}(\Omega_i - \Omega) \subset W$, $i = 1, 2$,
- (b) $\mathbf{M}(\partial\Omega_i) \leq L$, $i = 1, 2$,
- (c) $\mathcal{F}(\partial\Omega_1 - \partial\Omega_2) \leq \delta$,

there exist a sequence $\Omega_1 = \Lambda_0, \Lambda_1, \dots, \Lambda_m = \Omega_2 \in \mathcal{C}(M)$ such that for each $j = 0, \dots, m-1$,

- (i) $\text{spt}(\Lambda_j - \Omega) \subset U$;
- (ii) $\mathcal{A}^h(\Lambda_j) \leq \max\{\mathcal{A}^h(\Omega_1), \mathcal{A}^h(\Omega_2)\} + \eta$;
- (iii) $\mathbf{M}(\partial\Lambda_j - \partial\Lambda_{j+1}) \leq \eta$.

Appendix B. Interpolation process

Proof of Claim 2 in Proposition 5.1. Here we describe the construction of $\{\phi_i\}$ by interpolating $\{\phi_i^1\}$.

Fix $i \in \mathbb{N}$ and consider a 1-cell $\alpha \in I(1, k_i)$. We only need to show how to interpolate ϕ_i^1 when restricted to α_0 . For notational simplicity we write $\alpha = [0, 1]$. For $x \in \alpha$ let $\tilde{X}(x)$ be the linear interpolation between $\tilde{X}_i(0) = \tilde{X}(|\partial\phi_i^*(0)|)$ and $\tilde{X}_i(1) = \tilde{X}(|\partial\phi_i^*(1)|)$. The continuity of the map $V \rightarrow \tilde{X}(V)$ implies that $\|\tilde{X}_i(x) - \tilde{X}_i(0)\|_{C^1(M)} \rightarrow 0$ uniformly as $i \rightarrow \infty$. Define $\bar{Q}_i(x)$ to be the push-forward of $\phi_i^*(0)$ by the flow of $\tilde{X}_i(x)$ up to time 1; this gives a map $\bar{Q}_i : \alpha \rightarrow \mathcal{C}(M)$. Note that $\partial\bar{Q}_i : \alpha \rightarrow \mathcal{Z}_n(M)$ is continuous under the \mathbf{F} -metric.

Since $\bar{Q}_i(x)$ and $\phi_i^1(0)$ are the push-forwards of the same initial set $\phi_i^*(0)$ under the flows of $\tilde{X}_i(x)$ and $\tilde{X}_i(0)$ respectively, we have

$$(B.1) \quad \begin{aligned} \mathbf{F}(\partial\bar{Q}_i(x), \partial\phi_i^1(0)) &\rightarrow 0 \\ \mathbf{M}(\bar{Q}_i(x) - \phi_i^1(0)) &\rightarrow 0 \end{aligned} \quad , \text{ uniformly in } x, \alpha \text{ as } i \rightarrow \infty.$$

As $\bar{Q}_i(1)$ and $\phi_i^1(1)$ are the respective push-forwards of $\phi_i^*(0)$ and $\phi_i^*(1)$ under the same flow of $\tilde{X}_i(1)$, we have

$$(B.2) \quad \mathbf{M}(\partial\bar{Q}_i(1) - \partial\phi_i^1(1)) \rightarrow 0, \text{ uniformly in } \alpha \text{ as } i \rightarrow \infty.$$

Now we can apply the interpolation result [66, Theorem 5.1] (see also [45, Theorem 13.1]) to \bar{Q}_i , which gives that for any $\eta > 0$, there exist $l_\eta > 0$ and $Q_i : \alpha(l_\eta)_0 \rightarrow \mathcal{C}(M)$, such that

(i) given $x \in \alpha(l_\eta)_0$,

$$\mathbf{M}(\partial Q_i(x)) \leq \mathbf{M}(\partial \bar{Q}_i(x)) + \eta/2,$$

and

$$\mathbf{M}(Q_i(x) - \bar{Q}_i(x)) \leq \eta/(2c),$$

where $c = \sup |h|$, and hence

$$\mathcal{A}^h(Q_i(x)) \leq \mathcal{A}^h(\bar{Q}_i(x)) + \eta;$$

(ii) $\mathbf{f}(Q_i) \leq \eta$;

(iii) $\sup\{\mathcal{F}(\partial Q_i(x) - \partial \bar{Q}_i(x)) : x \in \alpha(l_\eta)_0\} < \eta$.

When $\eta \rightarrow 0$, by (i, iii) and [51, 2.1(20)] (see also [45, Lemma 4.1]), we have

$$\limsup_{\eta \rightarrow 0} \{\mathbf{F}(\partial Q_i(x), \partial \bar{Q}_i(x)) : x \in \alpha(l_\eta)_0\} = 0.$$

Take a sequence $\eta_i \rightarrow 0$, and denote $l_i = k_i + l_{\eta_i} + 1$, then we construct $\phi_i : I(1, k_i + l_{\eta_i} + 1) \rightarrow \mathcal{C}(M)$ by defining ϕ_i on each $\alpha(l_{\eta_i} + 1)_0$ by

$$\phi_i(x) = \begin{cases} Q_i(3x) & \text{for } x \in [0, 1/3] \cap \alpha(l_{\eta_i} + 1)_0 \\ \phi_i^1(1) & \text{otherwise.} \end{cases}$$

The desired properties (a, b, c, d) of ϕ_i follow straightforwardly from (B.1)–(B.2) and the properties of Q_j . Since \bar{Q}_i is obtained from a continuous deformation from ϕ_i^* , a further interpolation argument shows that S is homotopic to S^* , and hence we finish the proof of Claim 2. \square

References

- [1] Ian Agol, Fernando C. Marques, and André Neves. Min-max theory and the energy of links. *J. Amer. Math. Soc.*, 29(2):561–578, 2016.
- [2] Giovanni Alessandrini and Sergio Vessella. Remark on the strong unique continuation property for parabolic operators. *Proc. Amer. Math. Soc.*, 132(2):499–501, 2004.
- [3] Frederick Justin Almgren, Jr. The homotopy groups of the integral cycle groups. *Topology*, 1:257–299, 1962.

- [4] Frederick Justin Almgren, Jr. The theory of varifolds. *Mimeographed notes, Princeton*, 1965.
- [5] P. Amster and M. C. Mariani. The prescribed mean curvature equation for nonparametric surfaces. *Nonlinear Anal.*, 52(4):1069–1077, 2003.
- [6] Vladimir I. Arnold. *Arnold's problems*. Springer-Verlag, Berlin; PHASIS, Moscow, 2004. Translated and revised edition of the 2000 Russian original, With a preface by V. Philippov, A. Yakivchik and M. Peters.
- [7] N. Aronszajn. A unique continuation theorem for solutions of elliptic partial differential equations or inequalities of second order. *J. Math. Pures Appl. (9)*, 36:235–249, 1957.
- [8] Christian Bär. Zero sets of solutions to semilinear elliptic systems of first order. *Invent. Math.*, 138(1):183–202, 1999.
- [9] J. Lucas Barbosa, Manfredo do Carmo, and Jost Eschenburg. Stability of hypersurfaces of constant mean curvature in Riemannian manifolds. *Math. Z.*, 197(1):123–138, 1988.
- [10] Thomas Beck, Spencer Becker-Kahn, and Boris Hanin. Nodal sets of smooth functions with finite vanishing order and p -sweepouts. *Calc. Var. Partial Differential Equations*, 57(5):Art. 140, 13, 2018.
- [11] Pierre Bérard and Daniel Meyer. Inégalités isopérimétriques et applications. *Ann. Sci. École Norm. Sup. (4)*, 15(3):513–541, 1982.
- [12] Otis Chodosh and Christos Mantoulidis. Minimal surfaces and the allencahn equation on 3-manifolds: index, multiplicity, and curvature estimates. *to appear in Ann. Math.*, *arXiv:1803.02716*, 2018.
- [13] Tobias H. Colding and Camillo De Lellis. The min-max construction of minimal surfaces. In *Surveys in differential geometry, Vol. VIII (Boston, MA, 2002)*, volume 8 of *Surv. Differ. Geom.*, pages 75–107. Int. Press, Somerville, MA, 2003.
- [14] Tobias Holck Colding and William P. Minicozzi, II. *A course in minimal surfaces*, volume 121 of *Graduate Studies in Mathematics*. American Mathematical Society, Providence, RI, 2011.
- [15] Camillo De Lellis and Jusuf Ramic. Min-max theory for minimal hypersurfaces with boundary. *Ann. Inst. Fourier (Grenoble)*, 68(5):1909–1986, 2018.
- [16] Camillo De Lellis and Dominik Tasnady. The existence of embedded minimal hypersurfaces. *J. Differential Geom.*, 95(3):355–388, 2013.

- [17] Frank Duzaar. On the existence of surfaces with prescribed mean curvature and boundary in higher dimensions. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, 10(2):191–214, 1993.
- [18] Frank Duzaar and Joseph F. Grotowski. Existence and regularity for higher-dimensional H -systems. *Duke Math. J.*, 101(3):459–485, 2000.
- [19] Frank Duzaar and Klaus Steffen. Existence of hypersurfaces with prescribed mean curvature in Riemannian manifolds. *Indiana Univ. Math. J.*, 45(4):1045–1093, 1996.
- [20] L. C. Evans and J. Spruck. Motion of level sets by mean curvature. I. *J. Differential Geom.*, 33(3):635–681, 1991.
- [21] F. J. Fernandez. Unique continuation for parabolic operators. II. *Comm. Partial Differential Equations*, 28(9-10):1597–1604, 2003.
- [22] Robert Finn. *Equilibrium capillary surfaces*, volume 284 of *Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences]*. Springer-Verlag, New York, 1986.
- [23] Claus Gerhardt. Existence, regularity, and boundary behavior of generalized surfaces of prescribed mean curvature. *Math. Z.*, 139:173–198, 1974.
- [24] Claus Gerhardt. Closed hypersurfaces of prescribed mean curvature in locally conformally flat Riemannian manifolds. *J. Differential Geom.*, 48(3):587–613, 1998.
- [25] Mariano Giaquinta. On the Dirichlet problem for surfaces of prescribed mean curvature. *Manuscripta Math.*, 12:73–86, 1974.
- [26] Viktor L. Ginzburg. On closed trajectories of a charge in a magnetic field. An application of symplectic geometry. In *Contact and symplectic geometry (Cambridge, 1994)*, volume 8 of *Publ. Newton Inst.*, pages 131–148. Cambridge Univ. Press, Cambridge, 1996.
- [27] Enrico Giusti. On the equation of surfaces of prescribed mean curvature. Existence and uniqueness without boundary conditions. *Invent. Math.*, 46(2):111–137, 1978.
- [28] Enrico Giusti. *Minimal surfaces and functions of bounded variation*, volume 80 of *Monographs in Mathematics*. Birkhäuser Verlag, Basel, 1984.
- [29] Marco A. M. Guaraco. Min-max for phase transitions and the existence

- of embedded minimal hypersurfaces. *J. Differential Geom.*, 108(1):91–133, 2018.
- [30] Robert Gulliver and Joel Spruck. The Plateau problem for surfaces of prescribed mean curvature in a cylinder. *Invent. Math.*, 13:169–178, 1971.
 - [31] Robert Gulliver and Joel Spruck. Existence theorems for parametric surfaces of prescribed mean curvature. *Indiana Univ. Math. J.*, 22:445–472, 1972/73.
 - [32] Robert D. Gulliver, II. The Plateau problem for surfaces of prescribed mean curvature in a Riemannian manifold. *J. Differential Geometry*, 8:317–330, 1973.
 - [33] Robert Hardt and Leon Simon. Nodal sets for solutions of elliptic equations. *J. Differential Geom.*, 30(2):505–522, 1989.
 - [34] Reese Harvey and Blaine Lawson. Extending minimal varieties. *Invent. Math.*, 28:209–226, 1975.
 - [35] Stefan Hildebrandt. Randwertprobleme für Flächen mit vorgeschriebener mittlerer Krümmung und Anwendungen auf die Kapillaritätstheorie. I. Fest vorgebener Rand. *Math. Z.*, 112:205–213, 1969.
 - [36] David Hoffman and William H. Meeks, III. The strong halfspace theorem for minimal surfaces. *Invent. Math.*, 101(2):373–377, 1990.
 - [37] Kei Irie, Fernando C. Marques, and André Neves. Density of minimal hypersurfaces for generic metrics. *Ann. of Math. (2)*, 187(3):963–972, 2018.
 - [38] Daniel Ketover and Xin Zhou. Entropy of closed surfaces and min-max theory. *J. Differential Geom.*, 110(1):31–71, 2018.
 - [39] Brett Kotschwar. A short proof of backward uniqueness for some geometric evolution equations. *Internat. J. Math.*, 27(12):1650102, 17, 2016.
 - [40] Steven G. Krantz and Harold R. Parks. *A primer of real analytic functions*, volume 4 of *Basler Lehrbücher [Basel Textbooks]*. Birkhäuser Verlag, Basel, 1992.
 - [41] Man-Chun Lee and John Man-shun Ma. Uniqueness theorem for non-compact mean curvature flow with possibly unbounded curvatures. *arXiv:1709.00253*, 2017.

- [42] Martin Li and Xin Zhou. Min-max theory for free boundary minimal hypersurfaces I-regularity theory. *to appear in J. Differential Geom.*, *arXiv:1611.02612*, 2016.
- [43] Yevgeny Liokumovich, Fernando C. Marques, and André Neves. Weyl law for the volume spectrum. *Ann. of Math. (2)*, 187(3):933–961, 2018.
- [44] Fernando C. Marques and André Neves. Rigidity of min-max minimal spheres in three-manifolds. *Duke Math. J.*, 161(14):2725–2752, 2012.
- [45] Fernando C. Marques and André Neves. Min-max theory and the Willmore conjecture. *Ann. of Math. (2)*, 179(2):683–782, 2014.
- [46] Fernando C. Marques and André Neves. Morse index and multiplicity of min-max minimal hypersurfaces. *Camb. J. Math.*, 4(4):463–511, 2016.
- [47] Fernando C. Marques and André Neves. Existence of infinitely many minimal hypersurfaces in positive Ricci curvature. *Invent. Math.*, 209(2):577–616, 2017.
- [48] Fernando C. Marques and André Neves. Morse index of multiplicity one min-max hypersurfaces. *arXiv:1803.04273v1*, 2018.
- [49] Frank Morgan. Regularity of isoperimetric hypersurfaces in Riemannian manifolds. *Trans. Amer. Math. Soc.*, 355(12):5041–5052, 2003.
- [50] Stefano Nardulli. The isoperimetric profile of a smooth riemannian manifold for small volumes. *Annals of global analysis and geometry*, Vol. 36, No. 2 (2009), p., 36(2):111–132, 2009.
- [51] Jon T. Pitts. *Existence and regularity of minimal surfaces on Riemannian manifolds*, volume 27 of *Mathematical Notes*. Princeton University Press, Princeton, N.J.; University of Tokyo Press, Tokyo, 1981.
- [52] Harold Rosenberg and Matthias Schneider. Embedded constant-curvature curves on convex surfaces. *Pacific J. Math.*, 253(1):213–218, 2011.
- [53] Matthias Schneider. Closed magnetic geodesics on S^2 . *J. Differential Geom.*, 87(2):343–388, 2011.
- [54] Richard Schoen and Leon Simon. Regularity of stable minimal hypersurfaces. *Comm. Pure Appl. Math.*, 34(6):741–797, 1981.
- [55] Richard Schoen, Leon Simon, and Shing-Tung Yau. Curvature estimates for minimal hypersurfaces. *Acta Math.*, 134(3-4):275–288, 1975.
- [56] James Serrin. On the strong maximum principle for quasilinear second order differential inequalities. *J. Functional Analysis*, 5:184–193, 1970.

- [57] Leon Simon. *Lectures on geometric measure theory*, volume 3 of *Proceedings of the Centre for Mathematical Analysis, Australian National University*. Australian National University, Centre for Mathematical Analysis, Canberra, 1983.
- [58] Francis R. Smith. *On the existence of embedded minimal 2-spheres in the 3-sphere, endowed with an arbitrary Riemannian metric*. PhD thesis, Phd thesis, Supervisor: Leon Simon, University of Melbourne, 1982.
- [59] Antoine Song. Existence of infinitely many minimal hypersurfaces in closed manifolds. *arXiv:1806.08816v1*, 2018.
- [60] Klaus Steffen. On the existence of surfaces with prescribed mean curvature and boundary. *Math. Z.*, 146(2):113–135, 1976.
- [61] Andrejs E. Treibergs and S. Walter Wei. Embedded hyperspheres with prescribed mean curvature. *J. Differential Geom.*, 18(3):513–521, 1983.
- [62] Sergio Vessella. Unique continuation properties and quantitative estimates of unique continuation for parabolic equations. In *Handbook of differential equations: evolutionary equations. Vol. V*, Handb. Differ. Equ., pages 421–500. Elsevier/North-Holland, Amsterdam, 2009.
- [63] Brian White. The maximum principle for minimal varieties of arbitrary codimension. *Comm. Anal. Geom.*, 18(3):421–432, 2010.
- [64] Shing Tung Yau. Problem section. In *Seminar on Differential Geometry*, volume 102 of *Ann. of Math. Stud.*, pages 669–706. Princeton Univ. Press, Princeton, N.J., 1982.
- [65] Shing Tung Yau. A remark on the existence of sphere with prescribed mean curvature. *Asian J. Math.*, 1(2):293–294, 1997.
- [66] Xin Zhou. Min-max hypersurface in manifold of positive Ricci curvature. *J. Differential Geom.*, 105(2):291–343, 2017.
- [67] Xin Zhou. On the multiplicity one conjecture in min-max theory. *arXiv:1901.01173*, 2019.
- [68] Xin Zhou and Jonathan J. Zhu. Min-max theory for constant mean curvature hypersurfaces. *Invent. Math. (2019) 218:441–490*, 2019.

XIN ZHOU

DEPARTMENT OF MATHEMATICS

UNIVERSITY OF CALIFORNIA SANTA BARBARA

SANTA BARBARA, CA 93106

USA

E-mail address: zhou@math.ucsb.edu

JONATHAN ZHU
DEPARTMENT OF MATHEMATICS
PRINCETON UNIVERSITY
PRINCETON, NJ 08544
USA
E-mail address: jjzhu@math.princeton.edu

RECEIVED MARCH 22, 2019