

## ON $H$ -SPACES AND A CONGRUENCE OF CATALAN NUMBERS

TAMAR FRIEDMANN AND JOHN HARPER

(communicated by Donald M. Davis)

### Abstract

For  $p$  an odd prime and  $F$  the cyclic group of order  $p$ , we show that the number of conjugacy classes of embeddings of  $F$  in  $SU(p)$  such that no element of  $F$  has 1 as an eigenvalue is  $(1 + C_{p-1})/p$ , where  $C_{p-1}$  is a Catalan number. We prove that the only coset space  $SU(p)/F$  that admits a  $p$ -local  $H$ -structure is the classical Lie group  $PSU(p)$ . We also show that  $SU(4)/\mathbb{Z}_3$ , where  $\mathbb{Z}_3$  is embedded off the center of  $SU(4)$ , is a novel example of an  $H$ -space, even globally. We apply our results to the study of homotopy classes of maps from  $BF$  to  $BSU(n)$ .

## 1. Introduction

In [FS] conjugacy classes of elements of finite order dividing  $m$  in  $SU(n)$  are identified and counted. That result poses the question whether any of these elements can be used to produce new finite  $H$ -spaces with non-trivial fundamental group.

Let  $F$  denote a subgroup of  $SU(n)$  that is cyclic of finite order dividing  $m$ . Since  $SU(n)$  is the union of conjugacy classes of a maximal torus, we may assume that  $F$  is contained in a maximal torus of  $SU(n)$ . In order for a coset space  $SU(n)/F$  to admit the structure of an  $H$ -space, there are restrictions on the values of  $n$  and  $m$ . The basic topological restriction comes from W. Browder's work on differential Hopf-algebras [Br]. In particular, if  $p$  is a prime that divides  $m$ , then from Theorem 4.7 of [Br] it follows that the rational cohomology of  $SU(n)$  must have a generator in some dimension of the form  $2p^f - 1$ . Recall that the generators for  $SU(n)$  have odd dimensions  $2k - 1$  with  $k$  between 2 and  $n$ . Thus, the minimal case that satisfies the restriction is  $f = 1$  and  $m = n = p$ , and we consider this case, as well as the case  $f = 1$ ,  $m = 3$ ,  $n = 4$ .

When  $m = n = p$ , we show that it is only when  $F$  is the center of  $SU(p)$  that  $SU(p)/F$  is an  $H$ -space (Theorem 2.1), yet we find a new example of an  $H$ -space when  $m = 3$  and  $n = 4$  (Theorem 5.1). As far as we know, previous examples of non-classical finite (possibly  $p$ -local)  $H$ -spaces with a non-trivial fundamental group have used the center of  $SU(n)$  as an ingredient [H, KK]. But the center of  $SU(4)$  is  $\mathbb{Z}_4$ , so our example is off-center.

Borel's structure theorem for  $H$ -spaces, which plays a central role in the proof of Theorem 2.1, requires that when  $m = p$ , for  $SU(p)/F$  to be an  $H$ -space the embedding

---

Received September 8, 2016, revised November 10, 2016; published on July 19, 2017.

2010 Mathematics Subject Classification: 11A07, 11B50, 55P45, 05E15, 05A15.

Key words and phrases:  $H$ -space, Catalan number, homotopy class of maps.

Article available at <http://dx.doi.org/10.4310/HHA.2017.v19.n2.a2>

Copyright © 2017, Tamar Friedmann and John Harper. Permission to copy for private use granted.

$t$  of  $F$  in  $SU(p)$  is such that no eigenvalue of  $t(g)$ ,  $g \in F$  is equal to 1. Generalizing the eigenvalue condition to any  $SU(n)/\mathbb{Z}_m$  leads us to count, a-la [FS], the number of conjugacy classes of elements of finite order dividing  $m$  in  $SU(n)$  with no eigenvalue equal to 1, for any pair of positive integers  $m$  and  $n$  (Theorem 3.2). We call these *special* conjugacy classes.

The combinatorics for the class of cases where  $n = m = p$  for  $p > 3$  yields many examples of embeddings of  $F$  not conjugate to the center of  $SU(p)$ , for which – as we show –  $SU(p)/F$  is not an  $H$ -space. The combinatorics also contains a pleasant surprise; the number of special conjugacy classes of  $\mathbb{Z}_p$  subgroups in  $SU(p)$  is given by  $(1 + C_{p-1})/p$  where  $C_n$  is the  $n$ -th Catalan number (Theorem 3.5; a related observation involving a pair of distinct primes  $p$  and  $q$  appears in Theorem 3.4). As far as we know, the first observation that this expression yields an integer appears in OEIS #A098796, submitted by Chapoton; our result may be viewed as a proof of this fact. The related fact that  $p \nmid C_{p-1}$  is well-known [AK].

The combinatorial count in Theorem 3.2 is of additional interest based on a theorem proved independently by Dwyer and Wilkerson [DW] and by Zabrodsky [Z] and Notbohm [N]. The theorem asserts (among other things) that up to homotopy, essential maps of  $BF$  to  $BSU(n)$  are in one-to-one correspondence with conjugacy classes of non-constant homomorphisms of  $F$  to  $SU(n)$ , where  $F$  is a cyclic group of prime order  $p$ . Hence Theorem 3.2 and its corollary count the number of essential maps of  $BF$  to  $BSU(n)$  that do not factor through  $BSU(n-1)$ . We draw attention to the fact that other results in [FS] may be interpreted in a similar manner.

As of this writing, the general question of whether coset spaces of the form  $SU(n)/F$  are  $H$ -spaces for any  $n$  and for  $F$  a cyclic group of any order  $m$  remains open. We do expect, however, that in addition to the  $SU(4)/\mathbb{Z}_3$  that we already found there are other examples of  $H$ -spaces of the form  $SU(n)/F$  that do not use the center of  $SU(n)$  as an ingredient.

Throughout this paper,  $\mathbb{Z}_n$  shall denote the cyclic group of order  $n$ .

## 2. The coset spaces $SU(p)/F$

The Borel structure theorem for finite  $H$ -spaces [Bo] states that for  $p$  an odd prime, the mod  $p$  cohomology algebra is a tensor product of exterior algebras on odd degree elements and a truncated polynomial algebra on even degree elements with truncation at a power of  $p$ . We use it in what follows.

**Theorem 2.1.** *Let  $p$  be an odd prime and let  $F \subset SU(p)$  be a cyclic subgroup of order  $p$ . Then  $H^*(SU(p)/F; \mathbb{F}_p)$  satisfies the Borel structure theorem for finite  $H$ -spaces if and only if  $F$  is the center of  $SU(p)$ .*

*Proof.* Let  $\omega = e^{2\pi i/p}$  and let  $g$  be a generator for  $F$ . We describe  $F \subset SU(p)$  in terms of the following diagram:

$$\begin{array}{ccccc}
 g & & F & \xrightarrow{t} & SU(p) \\
 \downarrow & & \downarrow & & \searrow \\
 \omega & & S^1 & \xrightarrow{\Delta} & S^1 \times \cdots \times S^1 & \xrightarrow{\theta} & S^1 \times \cdots \times S^1 & \hookrightarrow & U(p)
 \end{array}$$

$\underbrace{\hspace{10em}}_p \qquad \qquad \qquad \underbrace{\hspace{10em}}_p$

where  $\Delta(z) = (z, z, \dots, z)$  and  $\theta(z_1, \dots, z_p) = (z_1^{m_1}, \dots, z_p^{m_p})$  with  $\sum_{i=1}^p m_i \equiv 0 \pmod{p}$ . Then  $t(g)$  is the diagonal matrix

$$\begin{pmatrix} \omega^{m_1} & & & 0 \\ & \omega^{m_2} & & \\ & & \ddots & \\ 0 & & & \omega^{m_p} \end{pmatrix}.$$

We follow the argument on p. 314 of [BB] applied to the commutative diagram of principal fiber bundles

$$\begin{array}{ccc} SU(p) & \longrightarrow & SU(p) \\ \downarrow & & \downarrow \\ EF \times_F SU(p) & \longrightarrow & ESU(p) \\ \downarrow & & \downarrow \\ K(\mathbb{Z}_p, 1) & \xrightarrow{Bt} & BSU(p) \end{array}, \quad (1)$$

with the Borel construction in place of  $SU(p)/F$  in the left column. Computing  $(Bt)^*$  on the total Chern class  $c$  we have

$$(Bt)^*(c) = (1 + m_1 y_2)(1 + m_2 y_2) \cdots (1 + m_p y_2) = 1 + \sum_{k=1}^p \sigma_k(\overline{m}) y_2^k$$

and

$$(Bt)^* c_k = \sigma_k(\overline{m}) y_2^k,$$

where  $\sigma_k$  is the  $k$ -th elementary symmetric polynomial in  $p$  variables,  $\overline{m}$  is  $(m_1, m_2, \dots, m_p)$ ,  $c_k$  is the  $k$ -th Chern class, and  $y_2$  generates  $H^2(K(\mathbb{Z}_p, 1); \mathbb{F}_p)$ .

Now  $(Bt)^* c_1 = 0$  because  $\sigma_1(\overline{m}) = \sum m_i \equiv 0 \pmod{p}$  as  $F \subset SU(p)$ . The first non-trivial differential in the Serre spectral sequence for the left side of diagram (1) is determined by the smallest value of  $k$  where  $(Bt)^* c_k \neq 0$ . If 1 were an eigenvalue of  $t(g)$  (i.e some  $m_i \equiv 0 \pmod{p}$ ) then  $k \leq p - 1$ . Assuming  $k \leq p - 1$ , we next show that the Borel structure theorem is not satisfied by the mod  $p$  cohomology of  $SU(p)/F$ .

We have

$$\begin{aligned} E_2^{*,*} &= H^*(K(\mathbb{Z}_p, 1); \mathbb{F}_p) \otimes H^*(SU(p); \mathbb{F}_p) \\ &\equiv \Lambda(x_1) \otimes \mathbb{Z}_p[y_2] \otimes \Lambda(f_3, \dots, f_{2p-1}), \end{aligned}$$

and our hypothesis is

$$d_{2k-1}(f_{2k-1}) = \lambda y_2^k \quad \text{where } \lambda \not\equiv 0 \pmod{p}.$$

Then

$$E_{2k}^{*,*} = \Lambda(x_1) \otimes \mathbb{Z}_p[y_2]/y_2^k \otimes \Lambda(f_3, \dots, \hat{f}_{2k-1}, \dots, f_{2p-1})$$

omitting the generator  $f_{2k-1}$ , and

$$E_{2k}^{*,*} = E_\infty^{*,*} = E_\infty^{*,0} \otimes E_\infty^{0,*},$$

for dimension reasons and by inspection. The bidegrees of the generators are

$$x_1 \in (1, 0), \quad y_2 \in (2, 0), \quad f_i \in (0, 2i - 1).$$

Thus the mod  $p$  cohomology algebra of the total space is generated by elements in these bidegrees. It follows that this algebra is the homology of the free algebra on these generators subject to the differential  $d_{2k-1}f_{2k-1} = y_2^k$ . If  $k \neq p$ , this algebra fails to satisfy Borel's theorem. If all the differentials are 0 then  $E_\infty^{*,*}$  is not a finite algebra. Thus,  $f_{2p-1}$  transgresses to a non-zero multiple of  $y_2^p$  and  $m_i$  is not congruent to 0 mod  $p$  and likewise for the  $p$ -th symmetric product.

Thus 1 is not an eigenvalue of  $t(g)$ ,  $\sigma_k(\overline{m}) \equiv 0 \pmod{p}$  for  $1 \leq k \leq p-1$ , and  $\sigma_p(\overline{m}) \not\equiv 0 \pmod{p}$ .

It now follows from the unique factorization for polynomials in  $\mathbb{F}_p(x)$  that  $m_1 = m_2 = \dots = m_p$ , so that  $F$  is the center of  $SU(p)$ .  $\square$

**Corollary 2.2.** *The only quotient of  $SU(p)$  by a finite group of order  $p$  that yields an  $H$ -space is  $PSU(p)$ .*

### 3. Embeddings of $\mathbb{Z}_m$ in $SU(n)$ and Catalan numbers

Here we count the number of ways in which an element of order  $m$  can be embedded in  $SU(n)$  such that 1 is not an eigenvalue. For the case where  $m$  and  $n$  are both prime, we also count the number of ways the group  $\mathbb{Z}_m$  can be embedded in  $SU(n)$  with the condition that 1 is not an eigenvalue of any generator of  $\mathbb{Z}_m$ .

Let  $N'(SU(n), m)$  be the number of conjugacy classes of elements of  $SU(n)$  of order  $m$ , none of whose eigenvalues is 1.<sup>1</sup>

Let  $F_1, F_2$  be  $\mathbb{Z}_m$  subgroups of  $SU(n)$ . We say  $F_1$  and  $F_2$  are in the same conjugacy class if there exists an element  $g \in SU(n)$  such that for any  $h \in F_1$ ,  $ghg^{-1} \in F_2$ .

**Definition 3.1.** A conjugacy class of  $\mathbb{Z}_m$  subgroups of  $SU(n)$  such that no element of the subgroups has eigenvalue 1 is called *special*. The number of such classes is denoted  $SpCG(SU(n), \mathbb{Z}_m)$ .

Note that each conjugacy class, including special ones, has at least one representative all of whose elements are diagonal.

**Theorem 3.2.** *For any positive integers  $m$  and  $n$ , we have*

$$N'(SU(n), m) = \frac{1}{m} \left( \sum_{d|(m,n)} \phi(d) \binom{m/d + n/d - 1}{n/d} - \sum_{d|(m,n-1)} \phi(d) \binom{m/d + \frac{n-1}{d} - 1}{\frac{n-1}{d}} \right), \quad (2)$$

where  $\phi$  is Euler's totient function.

---

<sup>1</sup>The notation  $N(SU(n), m)$  is reserved for the number of conjugacy classes of all elements of  $SU(n)$  of order  $m$  (including those with eigenvalue 1); it has been computed in [Dj, FS].

*Proof.* Let

$$F(x, t) = \prod_{k=1}^{m-1} \left( \sum_{a=0}^{\infty} (t^k x)^a \right).$$

A typical term in  $F(x, t)$  is

$$x^{\sum n_k} t^{\sum kn_k},$$

where  $n_k$ ,  $k = 1, \dots, m-1$  are non-negative integers. If  $\sum n_k = n$  and  $\sum kn_k \equiv 0 \pmod{m}$  then the sequence  $\{n_k\}$  corresponds to a diagonal  $SU(n)$  matrix of order  $m$  with eigenvalue  $e^{2\pi ik/m}$  repeated  $n_k$  times. Since we excluded  $k = 0$ , the matrix does not have 1 as an eigenvalue. Thus, such a sequence  $\{n_k\}$  corresponds to a conjugacy class counted by  $N'(SU(n), m)$ .

To pick out the terms in  $F(x, t)$  for which  $\sum kn_k \equiv 0 \pmod{m}$ , thereby obtaining a generating function for  $N'(SU(n), m)$ , let  $\zeta = \exp 2\pi i/m$  and recall

$$\frac{1}{m} \sum_{j=0}^{m-1} \zeta^{jb} = \begin{cases} 1, & \text{if } m \mid b, \\ 0, & \text{else.} \end{cases}$$

Define

$$G(x) = \frac{1}{m} \sum_{j=0}^{m-1} F(x, \zeta^j).$$

A typical term in  $G(x)$  is now

$$\frac{1}{m} \left( x^{\sum n_k} \zeta^{j \sum kn_k} \right),$$

so the sum over  $j$  picks out the terms with  $\sum kn_k \equiv 0 \pmod{m}$ , giving

$$G(x) = \sum_n N'(SU(n), m) x^n.$$

We have then

$$G(x) = \frac{1}{m} \sum_{j=0}^{m-1} \prod_{k=1}^{m-1} \left( \sum_{a=0}^{\infty} (\zeta^{jk} x)^a \right) = \frac{1}{m} \sum_{j=0}^{m-1} \prod_{k=1}^{m-1} \left( \frac{1}{1 - \zeta^{jk} x} \right).$$

The factorization  $1 - x^d = \prod_{\ell=0}^{d-1} (1 - \zeta^{j\ell} x)$  for  $\zeta^j$  a primitive  $d$ -th root of unity gives

$$\prod_{\ell=0}^{m-1} (1 - \zeta^{j\ell} x) = (1 - x^d)^{m/d} = (1 - x) \prod_{\ell=1}^{m-1} (1 - \zeta^{j\ell} x).$$

Together with the fact that  $\zeta^j$ ,  $j = 0, \dots, m-1$  is a primitive  $d$ -th root of unity  $\phi(d)$  times, we obtain

$$\begin{aligned} G(x) &= \frac{1}{m} \sum_{d|m} \phi(d) \frac{1-x}{(1-x^d)^{m/d}} \\ &= \frac{1}{m} \sum_{d|m} \phi(d) (1-x) \sum_{b \geq 0} \binom{m/d + b - 1}{b} x^d b \end{aligned}$$

$$= \frac{1}{m} \sum_{d|m} \phi(d) \left[ \sum_{b \geq 0} \binom{m/d + b - 1}{b} x^{db} - \sum_{b \geq 0} \binom{m/d + b - 1}{b} x^{db+1} \right].$$

The coefficient of  $x^n$  in  $G(x)$  will come from values of  $d$  such that  $db = n$  in the first term and  $db + 1 = n$  in the second term, leading to  $d|n$  and  $d|n - 1$ , respectively, as in equation (2).  $\square$

**Corollary 3.3.** *If  $p$  is a prime and  $n$  is any positive integer, then*

$$N'(SU(n), p) = \frac{1}{p} \left[ \binom{p+n-2}{n} + (p-1)\alpha(n, p) \right],$$

where

$$\alpha(n, p) = \begin{cases} 1 & \text{if } p \mid n, \\ -1 & \text{if } p \mid n - 1, \\ 0 & \text{otherwise.} \end{cases}$$

*Proof.* Follows from equation (2).  $\square$

In particular, if  $p$  and  $q$  are distinct primes, we have

$$N'(SU(p), q) = \begin{cases} \frac{1}{q} \binom{q+p-2}{p} & \text{if } q \nmid (p-1), \\ \frac{1}{q} \left[ \binom{q+p-2}{p} - (q-1) \right] & \text{if } q \mid (p-1), \end{cases} \quad (3)$$

and if  $q = p$ , we have

$$N'(SU(p), p) = \frac{1}{p} \left[ p - 1 + \binom{2p-2}{p} \right].$$

**Theorem 3.4.** *If  $p$  and  $q$  are distinct primes, and  $p \nmid q - 1$  then*

$$SpCG(SU(p), \mathbb{Z}_q) = \begin{cases} \frac{1}{(q-1)q} \binom{q+p-2}{p} = \frac{(q+p-2)!}{p!q!} & \text{if } q \nmid (p-1), \\ \frac{1}{(q-1)q} \left[ \binom{q+p-2}{p} - (q-1) \right] & \text{if } q \mid (p-1). \end{cases} \quad (4)$$

Further, if  $p = 2$  and  $m$  is any positive integer,  $SpCG(SU(2), \mathbb{Z}_m) = 1$ .

*Proof.* We first show that all the  $(q-1)$  generators of all  $\mathbb{Z}_q$  groups are in distinct conjugacy classes of  $SU(p)$ . It follows that we may divide the expressions in equation (3) by  $(q-1)$  and obtain the result.

Let  $F$  be a  $\mathbb{Z}_q$  subgroup of  $SU(p)$ , and let  $h \in F$ . Let  $\zeta = \exp 2\pi i/q$  and let  $n_k, k = 1, \dots, q-1$ , be the multiplicity of  $\zeta^k$  as an eigenvalue of  $h$  (so  $\sum n_k = p, \sum kn_k \equiv 0 \pmod{q}$ ).

Suppose  $h^t$  is in the same conjugacy class as  $h$  for some integer  $t$ . Then the multiplicity of the eigenvalues of  $h^t$  is the same as that for  $h$ . It follows that  $n_{kt} = n_k$  for all  $k$  (all indices are taken mod  $q$ ). It also follows that  $h^{lt}$  is in the same conjugacy class as  $h$  for any integer  $l$  (if  $h^t = ghg^{-1}$  for some  $g \in SU(p)$ , then  $h^{lt} = g^l h g^{-l}$ ). Therefore,  $n_{kt^l} = n_k$  for any  $l$  and  $k$ .

Let  $c$  be the order of  $t$ , i.e. the smallest integer such that  $t^c \equiv 1 \pmod{q}$  ( $c$  exists because  $q$  is prime). We have  $n_1 = n_t = n_{t^2} = \dots = n_{t^{c-1}}$ , all indices being distinct mod  $q$ . Let  $A_{i_1} = \{1, t, t^2, \dots, t^{c-1}\}$ . Now pick any  $i_2 \notin A_{i_1}$  and let  $A_{i_2} = \{i_2, i_2 t, i_2 t^2, \dots, i_2 t^{c-1}\}$ . Then  $A_{i_2}$  has  $c$  distinct elements mod  $q$ , as  $i_2 t^{\ell_1} \equiv i_2 t^{\ell_2} \pmod{q}$  implies  $\ell_1 - \ell_2 \geq c$ . Further,  $A_{i_2}$  does not intersect with  $A_{i_1}$ , as  $t^{\ell_1} \equiv i_2 t^{\ell_2} \pmod{q}$  implies  $i_2 \equiv t^{\ell_1 - \ell_2} \pmod{q}$ , i.e.  $i_2 \in A_{i_1}$ , a contradiction. We see that the set of  $n_k$ 's can be partitioned into several subsets of size  $c$ , with all the  $n_k$ 's within each subset equal to each other.

Since  $\sum_k n_k = p$ , it follows that  $p$  is divisible by  $c$ , so  $c = 1$  or  $c = p$ . If  $c = 1$  then  $t \equiv 1 \pmod{q}$  and there are no conjugacies. If  $c = p$  then since  $c$  is the order of a subgroup of  $\mathbb{Z}_q^*$ , we have  $c \mid q - 1$ , so  $p \mid q - 1$  which we assumed is not the case.

For  $p = 2$ , if  $m$  is any positive integer there is exactly one conjugacy class of a  $\mathbb{Z}_m$  subgroup generated by  $\text{diag}(\zeta, \zeta^{m-1})$  where  $\zeta = e^{\frac{2\pi i}{q}}$ . So  $SpCG(SU(2), \mathbb{Z}_m) = 1$ .  $\square$

**Theorem 3.5.** *If  $p$  is prime, then*

$$SpCG(SU(p), \mathbb{Z}_p) = \frac{1}{p}(1 + C_{p-1}), \quad (5)$$

where

$$C_{p-1} = \frac{1}{p} \binom{2p-2}{p-1}$$

is the  $(p-1)$ -th Catalan number.

*Proof.* The same proof as the previous theorem holds, with  $q$  replaced by  $p$ .  $\square$

**Corollary 3.6.** *For any prime  $p$ , we have*

$$C_{p-1} \equiv -1 \pmod{p}.$$

*Proof.* Follows from Eq. (5).  $\square$

For all  $p \geq 5$ , the number given by Eq. (5) is larger than 1 and blows up quickly: for  $p = 5, 7, 11, \dots$  it is  $3, 19, 1527, \dots$ .<sup>2</sup> For  $p = 5$ , the resulting pair of subgroups are generated by  $\text{diag}(\zeta, \zeta, \zeta, \zeta^3, \zeta^4)$  and  $\text{diag}(\zeta, \zeta, \zeta^2, \zeta^2, \zeta^4)$ , where  $\zeta = e^{2\pi i/5}$ . Generators for the 18 groups at  $p = 7$  appear in Appendix A.

**Corollary 3.7.** *For any primes  $q$  and  $p$  such that  $q \mid p - 1$ ,*

$$\frac{1}{q-1} \binom{q+p-2}{p} \equiv 1 \pmod{q}.$$

*Proof.* Follows from Eq. (4) for  $q \mid p - 1$ .  $\square$

Both corollaries can be checked using Wilson's theorem,  $(p-1)! \equiv -1 \pmod{p}$ .

---

<sup>2</sup>As it happens, this series appears in the Online Encyclopaedia of Integer Sequences as A098796, submitted there by Chapoton in 2004 [Ch].

#### 4. Homotopy classes of maps

Our results may be applied to the study of homotopy classes of maps  $B\mathbb{Z}_p \rightarrow BSU(n)$ . Theorems proved independently by Dwyer and Wilkerson [DW] and by Zabrodsky [Z] and Notbohm [N] say that homotopy classes of essential maps are in one-to-one correspondence with conjugacy classes of non-constant homomorphisms of  $Z_p$  to  $SU(n)$ . These in turn are determined by conjugacy classes of non-identity elements of prime order  $p$ . To connect our results with this theory, we make a definition.

**Definition 4.1.** A homotopy class of essential maps from  $B\mathbb{Z}_m$  to  $BSU(n)$  is called special if it does not factor through  $BSU(n-1)$ .

Corresponding to the special conjugacy classes of elements counted by Theorem 3.2 and Corollary 3.3, we have the special homotopy classes of maps for  $m = p$ . Then the following is a restatement of Corollary 3.3.

**Corollary 4.2.** *For any prime  $p$  and any integer  $n$ , the number of special homotopy classes of essential maps from  $B\mathbb{Z}_p$  to  $BSU(n)$  is*

$$\frac{1}{p} \left[ \binom{p+n-2}{n} + (p-1)\alpha(n,p) \right],$$

where

$$\alpha(n,p) = \begin{cases} 1 & \text{if } p \mid n, \\ -1 & \text{if } p \mid n-1, \\ 0 & \text{otherwise.} \end{cases}$$

*In particular, the number of special homotopy classes of essential maps from  $B\mathbb{Z}_p$  to  $BSU(p)$  is*

$$\frac{1}{p} \left[ p-1 + \binom{2p-2}{p} \right].$$

#### 5. The space $SU(4)/\mathbb{Z}_3$

In this section, let  $F = \mathbb{Z}_3$  be the center of  $SU(3)$ . Embed  $F$  in  $SU(4)$  via the standard inclusion of  $SU(3)$  in  $SU(4)$  given in terms of matrices by

$$A \mapsto \begin{pmatrix} A & 0 \\ 0 & 1 \end{pmatrix}.$$

We have a diagram of inclusions of subgroups yielding  $S^5$  and  $S^7$  as homogeneous spaces in two ways,

$$\begin{array}{ccccc} S^3 & \hookrightarrow & SU(3) & \longrightarrow & S^5 \\ \downarrow & & \downarrow & & \\ Sp(2) & \hookrightarrow & SU(4) & \longrightarrow & S^5 \\ \downarrow & & \downarrow & & \\ S^7 & & S^7 & & \end{array}$$

We regard the spheres as right coset spaces. Taking left coset spaces with respect to  $F$ , we obtain the diagram

$$\begin{array}{ccccc}
 S^3 & \longrightarrow & PSU(3) & \longrightarrow & L(5; 1, 1, 1) \\
 \downarrow & & \downarrow & & \\
 Sp(2) & \hookrightarrow & SU(4)/F & \longrightarrow & L(5; 1, 1, 1) \\
 \downarrow & & \downarrow & & \\
 S^7 & & S^7 & & 
 \end{array} \tag{6}$$

where the lens space  $L(5; 1, 1, 1)$  is the quotient of the 5-sphere by the cyclic group of order 3.

In [H, p. 332], it is proven that after localizing at  $p = 3$ , the top row of diagram (6) splits; hence  $L(5; 1, 1, 1)$  is a 3-local  $H$ -space and  $SU(4)/F$  is 3-locally equivalent to  $Sp(2) \times L(5; 1, 1, 1)$  which is a 3-local  $H$ -space. Rationally,  $SU(4)/F$  is  $K(\mathbb{Q}, 3) \times K(\mathbb{Q}, 5) \times K(\mathbb{Q}, 7)$  which can only be primitively generated. Since the action by  $F$  is via maps homotopic to the identity, the spaces are simple and we may apply localization there to conclude that  $SU(4)/F$  is an  $H$ -space since each of its  $p$ -localizations are  $H$ -spaces and each rationalization map is an  $H$ -map. So we have proved:

**Theorem 5.1.** *The coset  $SU(4)/\mathbb{Z}_3$ , with  $\mathbb{Z}_3$  embedded via inclusion of the center of  $SU(3)$  in  $SU(4)$ , is an  $H$ -space.*

### Acknowledgments

We thank Richard Stanley and Frédéric Chapoton for helpful discussions. We also thank the referee for helpful comments.

### Appendix A. Generators for subgroups of $SU(7)$ of order 7

Below we list generators for the 18 subgroups of  $SU(7)$  of order 7 (not including the center). A set of 7 integers  $[a_1, \dots, a_7]$  corresponds to the generator  $\text{diag}(\zeta^{a_1}, \dots, \zeta^{a_7})$  where  $\zeta = e^{2\pi i/7}$ :

$[1, 2, 2, 2, 2, 2, 3],$	$[1, 1, 2, 2, 2, 3, 3],$	$[1, 1, 1, 2, 3, 3, 3],$	$[1, 1, 2, 2, 2, 2, 4],$
$[1, 1, 1, 2, 2, 3, 4],$	$[1, 1, 1, 1, 3, 3, 4],$	$[2, 3, 3, 3, 3, 3, 4],$	$[2, 2, 3, 3, 3, 4, 4],$
$[1, 3, 3, 3, 3, 4, 4],$	$[2, 2, 2, 3, 4, 4, 4],$	$[1, 2, 3, 3, 4, 4, 4],$	$[1, 1, 3, 4, 4, 4, 4],$
$[1, 1, 1, 1, 2, 3, 5],$	$[2, 2, 2, 3, 3, 4, 5],$	$[1, 2, 3, 3, 3, 4, 5],$	$[1, 2, 2, 3, 4, 4, 5],$
$[1, 1, 3, 3, 4, 4, 5],$	$[1, 1, 2, 3, 4, 5, 5].$		

### References

[AK] R. Alter and K. Kubota, “Prime and prime power divisibility of Catalan numbers,” *J. Combin. Theory Ser. A* **15** (1973), 243–256.

[Bo] A. Borel, “Sur la cohomologie des espaces fibrés principaux et des espaces homogènes de groupes de Lie compacts,” *Ann. of Math.* **57** (1953), 115–207.

- [Br] W. Browder, “On differential Hopf algebras,” *Trans. Amer. Math. Soc.* **107** (1963), 153–176.
- [BB] P.F. Baum and W. Browder, “The cohomology of quotients of classical groups,” *Topology* **3** (1965), 305–336.
- [Ch] The On-Line Encyclopedia of Integer Sequences, published electronically at <http://oeis.org>, 2010, Sequence A098796, submitted in 2004 by F. Chapoton.
- [Dj] D.Z. Djokovic, “On conjugacy classes of elements of finite order in compact or complex semisimple Lie groups,” *Proc. Amer. Math. Soc.* **80** (1980), no. 1, 181–184;  
D.Z. Djokovic, “On conjugacy classes of elements of finite order in complex semisimple Lie groups,” *J. Pure Appl. Algebra* **35** (1985), no. 1, 1–13.
- [DW] W.G. Dwyer and C.W. Wilkerson, “Maps of  $B\mathbb{Z}/p\mathbb{Z}$  to  $BG$ ,” Algebraic Topology – Rational Homotopy (Louvain-la-Neuve, 1986), 92–98, *Lecture Notes in Math.*, 1318, Springer, Berlin, 1988.
- [FS] T. Friedmann and R. Stanley, “Counting conjugacy classes of elements of finite order in Lie groups,” *European J. Combin.* **36** (2014), 86 [arXiv:1311.0599];  
T. Friedmann and R. Stanley, “The string landscape: On formulas for counting vacua,” *Nuclear Phys. B* **869** (2013), 74 [arXiv:1212.0583].
- [H] J. Harper, “Regularity of finite H-spaces,” *Illinois J. Math.* **23** (1979), no. 2, 330–333.
- [KK] D. Kishimoto and A. Kono, “Mod  $p$  decompositions of non-simply connected Lie groups,” *J. Math. Kyoto Univ.* **48** (2008), no. 1, 1–5.
- [N] D. Notbohm, “Maps between classifying spaces,” *Math. Z.* **207** (1991), 229–257.
- [Z] A. Zabrodsky, “On spaces of functions between classifying spaces,” *Israel J. Math.* **76** (1991) 1–26.

Tamar Friedmann `tfriedmann@smith.edu`

Department of Mathematics and Statistics, Burton Hall, Smith College, 44 College Lane, Northampton, MA 01060, USA

Department of Physics and Astronomy, University of Rochester, 500 Wilson Boulevard, Rochester, NY 14623, USA

John Harper `john.harper@rochester.edu`

Department of Mathematics, Hylan Building, University of Rochester, 500 Wilson Boulevard, Rochester, NY 14623, USA