

# Contact surgeries on Legendrian figure-eight knots

JAMES CONWAY

We show that all positive contact surgeries on every Legendrian figure-eight knot in  $(S^3, \xi_{\text{std}})$  result in an overtwisted contact structure. The proof uses convex surface theory and invariants from Heegaard Floer homology.

## 1. Introduction

Dehn surgery on knots has been a fruitful way to construct new contact structures on 3-manifolds, and in particular to try to construct new tight contact manifolds. When the knot in question is a Legendrian knot (*ie.* its tangent vectors lie in the contact planes), Dehn surgery with framing equal to the contact framing always results in an overtwisted contact manifold. The remaining framings break into two classes: those less than the contact framing, and those greater. Surgeries with these framings give rise to negative and positive contact surgery, respectively.

In [25], Wand showed that given a tight contact manifold, the result of negative contact surgery on any Legendrian knot is a tight contact manifold. Regarding positive contact surgery, much less is known; existing tightness results can be found in [2, 11, 16, 17, 19–21].

Most of the results for positive contact surgery prove tightness using various flavours of Heegaard Floer homology. In particular, the non-vanishing of the Heegaard Floer contact class shows that a contact manifold is tight, however its vanishing is not equivalent to a contact manifold being overtwisted. Several of the above results give conditions under which contact (+1)-surgery (*ie.* positive contact surgery with framing one more than the contact framing) has vanishing Heegaard Floer contact class.

There are fewer results that show that positive contact surgeries starting from a tight contact structure result in an overtwisted one. Lisca and Stipsicz showed in [18] that there exists a configuration in the front projection of a Legendrian knot that ensures contact (+1)-surgery on the knot is overtwisted. This configuration is not present in the figure-eight knot under

consideration in this paper (but it is present in the negative torus knots, for example). In [2], the author used versions of the Bennequin inequality (an inequality of Legendrian knot invariants that holds in tight contact manifolds) to give general results for when positive contact surgery on Legendrian knots is overtwisted.

After the unknot and the trefoils, the figure-eight knot is next natural knot to study (contact surgeries on the others were understood by [3, 16] for the unknot, [17] for the right-handed trefoil, and [2, 18] for the left-handed trefoil). The classification of Legendrian figure-eight knots in  $(S^3, \xi_{\text{std}})$  was undertaken by Etnyre and Honda in [8], who proved that all such Legendrian knots are classified up to isotopy by their Thurston–Bennequin number ( $tb$ ) and rotation class ( $rot$ ), and that all such knots destabilise to a Legendrian knot with  $tb = -3$  and  $rot = 0$ . Lisca and Stipsicz showed in [18] that the result of contact (+1)-surgery on any Legendrian figure-eight knot has vanishing Heegaard Floer contact class; we answer the natural follow-up question:

**Theorem 1.1.** *The results of all positive contact surgeries on any Legendrian figure-eight knot in  $(S^3, \xi_{\text{std}})$  are overtwisted.*

**Remark 1.2.** One should not conclude from Theorem 1.1 that the manifolds resulting from surgery on the figure-eight support no tight contact structure: in fact, they all support tight contact structures. However, they do not arise from positive contact surgery on a figure-eight knot in  $(S^3, \xi_{\text{std}})$ .

The proof uses convex surfaces and the Heegaard Floer contact class. In particular, given any Legendrian knot  $L$  we show that if any positive contact surgery on  $L$  is tight, then a particular contact structure on  $S^3 \setminus N(K)$  is also tight. For the figure-eight knot, we can show that this contact structure  $\xi$  on  $S^3 \setminus N(K)$  has vanishing Heegaard Floer contact class. We then use convex surfaces to classify all tight contact structures on  $S^3 \setminus N(K)$  that induce a particular set of dividing curves on a convex Seifert surface (the same set of curves can also be found in  $\xi$ ). We then construct these tight contact structures, and show that they have non-vanishing Heegaard Floer contact class. This shows that  $\xi$  is overtwisted, and proves Theorem 1.1.

Beyond the figure-eight knot, it is unclear how successful this approach will be. The facts that the figure-eight knot is fibred and has genus 1 play a large role in making the classification of relevant tight contact structures on  $S^3 \setminus N(K)$  possible. However, the approach of showing that a particular contact structure on  $S^3 \setminus N(K)$  is overtwisted is more widely applicable, as can be seen in [2].

In all known cases where the result of positive contact surgery on a Legendrian knot  $(S^3, \xi_{\text{std}})$  is tight, we also know that the Heegaard Floer contact invariant is non-vanishing. This paper, along with the results in [2], lend support toward a positive answer to this question:

**Question 1.3.** *Let  $(M, \xi)$  be the result of some positive contact surgery on a Legendrian knot in  $(S^3, \xi_{\text{std}})$ . Is  $\xi$  tight if and only if its Heegaard Floer contact class is non-vanishing?*

### Acknowledgements

The author is indebted to John Etnyre for many helpful discussions. He thanks Hyunki Min for pointing out a mistake in an earlier version of this paper. The author also thanks an anonymous referee for their comments. The results in this paper originally appeared as part of the author's doctoral thesis. This work was partially supported by NSF Grant DMS-1309073.

## 2. Contact geometric background

We begin with a brief reminder of standard theorems about contact structures on 3-manifolds which we will use throughout this paper. We assume a basic knowledge of contact structures at the level of [6, 7].

### 2.1. Farey graph

The Farey graph is the 1-skeleton of a tessellation of the hyperbolic plane by geodesic triangles shown in Figure 1, where the endpoints of the geodesics are labeled. The labeling, shown in Figure 1, is determined as follows: let the left-most point be labeled  $\infty = 1/0$  and the right-most point be labeled  $0$ . Given a geodesic triangle where two corners are already labeled  $a/b$  and  $c/d$ , then the third corner is labeled  $(a+c)/(b+d)$ . For triangles in the upper half of the plane, we treat  $0$  as  $0/(-1)$ , whereas for triangles in the lower half of the plane, we treat  $0$  as  $0/1$ . Thus, the labels on the upper half are all negative, and those on the lower half are all positive. Every rational number and infinity is found exactly once as a label on the Farey graph.

### 2.2. Convex surfaces

We introduce the basics of convex surfaces. See [5] for more details.




Figure 1: The Farey graph.

A surface  $\Sigma$  (possibly with boundary) in a contact manifold  $(M, \xi)$  is called *convex* if there exists a *contact vector field*  $v$  such that  $v$  is transverse to  $\Sigma$ . Here, a *contact vector field* is a vector field whose flow preserves the contact planes. Using the contact vector field  $v$ , it is not hard to see that convex surfaces have a neighbourhood contactomorphic to  $\Sigma \times \mathbb{R}$  with an  $\mathbb{R}$ -invariant contact structure, called a *vertically-invariant neighbourhood* of  $\Sigma$ .

Given a surface  $\Sigma$  in  $(M, \xi)$  and the characteristic foliation  $\mathcal{F}$  on  $\Sigma$  induced by  $\xi$ , we say that a multi-curve  $\Gamma$  on  $\Sigma$  *divides*  $\mathcal{F}$  if

- $\Sigma \setminus \Gamma = \Sigma_+ \sqcup \Sigma_-$ ,
- $\Gamma$  is transverse to the singular foliation  $\mathcal{F}$ , and

- there is a volume form  $\omega$  on  $\Sigma$  and a vector field  $w$  such that
  - $\pm \mathcal{L}_w \omega > 0$  on  $\Sigma_{\pm}$ ,
  - $w$  directs  $\mathcal{F}$ , and
  - $w$  points out of  $\Sigma_+$  along  $\Gamma$ .

**Theorem 2.1 (Giroux [10]).** *A closed surface  $\Sigma$  is  $C^\infty$ -close to a convex surface. If  $\Sigma$  is a surface with Legendrian boundary such that the twisting of the contact planes along each boundary component is non-positive when measured against the framing given by  $\Sigma$ , then  $\Sigma$  can be  $C^0$ -perturbed in a neighbourhood of the boundary and  $C^\infty$ -perturbed on its interior to be convex.*

*If  $\Sigma \subset (M, \xi)$  is an orientable surface, and its boundary (if it is non-empty) is Legendrian, then  $\Sigma$  is a convex surface if and only if its characteristic foliation has a dividing set. Given a convex surface  $\Sigma$  with dividing curves  $\Gamma$ , and any singular foliation  $\mathcal{F}$  on  $\Sigma$  divided by  $\Gamma$ , then  $\Sigma$  can be perturbed to a convex surface with characteristic foliation  $\mathcal{F}$ .*

In particular, convex surfaces are generic, and the germ of the contact structure at a convex surface is determined (up to a  $C^0$ -perturbation of the surface) by its dividing curves and the signs of the regions  $\Sigma_{\pm}$ .

A properly-embedded graph  $G$  on a convex surface  $\Sigma$  is *non-isolating* if  $G$  intersects the dividing curves  $\Gamma$  transversely, and each component of  $\Sigma \setminus G$  has non-trivial intersection with  $\Gamma$ .

**Theorem 2.2 (Honda [12]).** *If  $G$  is a non-isolating properly-embedded graph on a convex surface  $\Sigma$ , then there is an isotopy of  $\Sigma$  relative to its boundary such that  $G$  is contained in the new characteristic foliation. If  $G$  is a simple closed curve, then the twisting of the contact planes along  $L$  with respect to the framing on  $G$  given by  $\Sigma$  is equal to*

$$tw(G, \Sigma) = -\frac{|G \cap \Gamma|}{2}.$$

This is commonly called the *Legendrian realisation principle*. In particular, a simple closed curve in  $\Sigma$  that is non-separating can always be Legendrian realised on a convex surface. If  $L$  is a null-homologous Legendrian knot bounding a convex surface, then  $tw(L, \Sigma) = tb(L)$ , and so  $tb(L) = -|L \cap \Gamma|/2$ .

Giroux has shown that there are restrictions on dividing curves in tight manifolds. This result is often called *Giroux’s Criterion*.

**Theorem 2.3 (Giroux [10]).** *If  $\Sigma = S^2$  is convex, then a vertically-invariant neighbourhood of  $\Sigma$  is tight if and only if the dividing set  $\Gamma$  is connected. If*

$\Sigma \neq S^2$ , then a vertically-invariant neighbourhood of  $\Sigma$  is tight if and only if  $\Gamma$  has no contractible components.

Given two convex surfaces  $\Sigma_1$  and  $\Sigma_2$  that intersect in a Legendrian curve  $L$ , Kanda [14] and Honda [12] have shown that between each intersection of  $L$  with  $\Gamma_{\Sigma_1}$  is exactly one intersection of  $L$  with  $\Gamma_{\Sigma_2}$ , as in Figure 2. Honda further showed that there is a way to “round edges” at  $L$  and get a new convex surface. The dividing set on the new surface is derived from  $\Gamma_{\Sigma_i}$  as in Figure 3.




Figure 2: Two convex surfaces intersecting in a Legendrian curve. This figure is reproduced from [5, Figure 10].

A special case of “rounding edges” at the intersection of two convex surfaces is when  $\Sigma_2$  is a *bypass*. This is when  $\Sigma_2$  is a disc with Legendrian boundary with  $tb = -1$ , such that  $\Sigma_1 \cap \Sigma_2$  is an arc  $\alpha$  intersecting  $\Gamma_{\Sigma_1}$  in three points, two of which are the endpoints of  $\alpha$ ; we further require that the endpoints of  $\alpha$  are elliptic singularities of the characteristic foliation on  $\Sigma_2$ . By the above discussion, the dividing set  $\Gamma_{\Sigma_2}$  is a single arc with endpoints on  $\alpha$ . By Theorem 2.1, we can arrange for there to be a unique hyperbolic singularity on  $\partial\Sigma_2$  that lies on  $\alpha$  and is between the two points  $\alpha \cap \Gamma_{\Sigma_2}$ . The sign of this hyperbolic singularity is called the *sign* of the bypass.

Honda proved [12] that in a neighbourhood of  $\Sigma_1 \cup \Sigma_2$ , there is a one-sided neighbourhood  $\Sigma_1 \times [0, 1]$  of  $\Sigma_1$  such that  $\Sigma_1 \times \{0, 1\}$  is convex, the dividing curves on  $\Sigma_1 \times \{0\}$  are  $\Gamma_{\Sigma_1}$ , and the dividing curves on  $\Sigma_1 \times \{1\}$  are  $\Gamma_{\Sigma_1}$  changed along a neighbourhood of  $\alpha$  as in Figure 4. We say that the convex surface  $\Sigma_1 \times \{1\}$  is obtained from  $\Sigma_1$  by a bypass attachment along  $\Sigma_2$ .




Figure 3: “Rounding edges” of intersecting convex surfaces. This figure is reproduced from [5, Figure 11].




Figure 4: The result of performing a bypass on the dividing curves.

If  $\Sigma_1$  is a convex  $T^2$  (resp.  $T^2 \setminus D^2$ ) with 2 parallel dividing curves, then we can choose the characteristic foliation on  $\Sigma_1$  such that it consists of two curves called *Legendrian divides* parallel to the dividing curves along with a linear foliation of the torus by curves not parallel to the dividing curves, called *ruling curves*. Under these hypotheses, Honda proved [12] how the slopes of the dividing curves change under bypass attachments along a ruling curve. Denote the slope of curves parallel to  $(\frac{q}{p})$  by  $p/q$ , as in the Farey graph.

**Theorem 2.4 (Honda [12]).** *Let  $\Sigma_1$  have two dividing curves of slope  $s$  and ruling curves of slope  $r$ . Let  $\Sigma_2$  be a bypass attached to  $\Sigma_1$  along a ruling curve. Then the result  $\Sigma'_1$  of a bypass attachment along  $\Sigma_2$  has two dividing curves with slope  $s'$ , where  $s'$  is the label on the Farey graph clockwise of  $r$*

and counter-clockwise of  $s$ , and such that  $s'$  is the label closest to  $r$  with an edge to  $s$ .

**Remark 2.5.** If  $\Sigma_2$  is a bypass for  $\Sigma_1$  attached along the back of  $\Sigma_1$ , then the bypass attachment will change  $\Gamma_{\Sigma_1}$  in a manner similar to Figure 4 but reflected in the vertical axis. Theorem 2.4 will hold after reversing the words “clockwise” and “counter-clockwise”.

Bypasses are only useful if we can find them. To that effect, we have the Imbalance Principle, which allows us to find bypasses on annuli.

**Theorem 2.6 (Honda [12]).** *Let  $\Sigma$  and  $A = S^1 \times [0, 1]$  be two convex surfaces with Legendrian boundary, such that  $\Sigma \cap A = S^1 \times \{0\}$ . Then, if the twisting of the contact planes along the boundary of  $A$  satisfies*

$$tw(S^1 \times \{0\}, A) < tw(S^1 \times \{1\}, A) \leq 0,$$

*then there is a bypass for  $\Sigma$  along  $A$ , ie. some subsurface of  $A$  is a bypass for  $\Sigma$ .*

In particular, if  $S^1 \times \{1\}$  sits on a convex surface  $\Sigma'$ , and

$$|\Gamma_{\Sigma} \cap (S^1 \times \{0\})| > |\Gamma_{\Sigma'} \cap (S^1 \times \{0\})|,$$

then the hypotheses of Theorem 2.6 hold, and there is a bypass for  $\Sigma$  along  $A$ .

### 2.3. Basic slices

Consider the manifold  $(T^2 \times I, \xi)$ , with  $\xi$  tight. Let the two boundary components be convex with two dividing curves each, with slopes  $s_0$  and  $s_1$ . If  $s_0$  and  $s_1$  are labels on the Farey graph connected by a geodesic, then  $(T^2 \times I, \xi)$  is called a *basic slice* if the contact structure is *minimally twisting*, ie. if any boundary-parallel convex torus has dividing curves of slope clockwise of  $s_0$  and counter-clockwise of  $s_1$ . If not, then the manifold can be cut up into basic slices along boundary parallel convex tori, following the path between  $s_0$  and  $s_1$  along the Farey graph.

**Theorem 2.7 (Honda [12]).** *There are exactly two tight contact structures up to isotopy (and only one up to contactomorphism) on  $T^2 \times I$  with a fixed singular foliation on the boundary that is divided by two dividing curves on  $T^2 \times \{i\}$  for  $i = 0, 1$  each of slope  $s_i$ , where  $s_0$  and  $s_1$  are labels in the Farey graph connected by a geodesic.*

Given a contact structure  $\xi$  on  $M$  that is trivialised by  $v$  on  $\partial M$ , we can define a relative Euler class  $e(\xi, v) \in H^2(M, \partial M; \mathbb{Z})$ . Given a convex surface  $\Sigma$  with boundary on  $\partial M$ , whose oriented tangent vector on  $\partial\Sigma$  agrees with  $v$ , we can calculate

$$e(\xi, v)([\Sigma]) = \chi(\Sigma_+) - \chi(\Sigma_-).$$

For a basic slice with  $s_0 = -\infty$  and  $s_1 = -1$ , the relative Euler class acts as 0 on the annulus  $(\frac{1}{0}) \times [0, 1]$  and as  $\pm 1$  on the annulus  $(\frac{0}{-1}) \times [0, 1]$ , where the slope of  $(\frac{q}{p})$  is  $p/q$ . Every other basic slice can be put in this standard form by an element of  $SL_2(\mathbb{Z})$ . This calculation allows us to distinguish the basic slices by calling them *positive* and *negative* basic slices; this sign choice is also such that when gluing a negative (resp. positive) basic slice to the boundary of the complement of a standard neighbourhood of a Legendrian knot, the result is the complement of a standard neighbourhood of its negative (resp. positive) stabilisation.

In addition, this classification implies that if we have a basic slice  $(T^2 \times I, \xi)$  that can be broken up into two basic slices  $(T^2 \times [0, 1/2], \xi_1)$  and  $(T^2 \times [1/2, 1], \xi_2)$ , then the sign of each of the latter two basic slices agrees with the sign of  $(T^2 \times I, \xi)$ . Thus, if the signs disagree, then  $(T^2 \times I, \xi)$  is overtwisted (and hence by definition is not a basic slice).

## 2.4. Contact surgery

Given a null-homologous Legendrian knot  $L \subset (M, \xi)$ , we start by removing the interior of a standard neighbourhood  $N(L)$  of  $L$ , *ie.* the interior of a tight solid torus with convex boundary, where the dividing curves have the same slope as the contact framing  $tb(L)\mu + \lambda$ , where  $\mu$  is a meridian and  $\lambda$  is the Seifert framing of  $L$ .

To do *positive contact surgery* on  $L$ , we first glue a basic slice to  $\partial N(L)$  such that the new contact structure on  $M \setminus N(L)$  has convex boundary with two meridional dividing curves. Different sign choices on this basic slice in general give rise to distinct contact structures; we denote by  $\xi^+(L)$  (resp.  $\xi^-(L)$ ) the contact structure on  $M \setminus N(L)$  coming from gluing on a positive (resp. negative) basic slice. Finally, we then glue a solid torus to the boundary such that the desired topological surgery is achieved, and we extend the contact structure over the solid torus such that it is tight on the solid torus. Different choices of sign on the basic slice and different extensions over the solid torus will in general give rise to distinct contact structures on the surgered manifold, see [12, 14].

### 2.5. Heegaard Floer homology

We make use of invariants of contact structures coming from Heegaard Floer theory: for closed contact manifolds  $(M, \xi)$ , we have an element  $c(\xi) \in \widehat{HF}(-M)$  (see [22]), and for contact manifolds  $(M', \Gamma, \xi')$  with convex boundary, where  $\Gamma \subset \partial M'$  is the dividing set, we have an element  $EH(\xi) \in SFH(-M', -\Gamma)$  (see [13]). If  $(M', \Gamma, \xi') \subset (M, \xi)$  is a contact embedding, then there is a map  $SFH(-M', -\Gamma) \rightarrow \widehat{HF}(-M)$  that sends  $EH(\xi')$  to  $c(\xi)$ .

To a Legendrian knot  $L \subset (M, \xi)$ , we associate an element  $\widehat{\mathcal{L}}(L)$  (defined in [15]) in the knot Heegaard Floer group  $\widehat{HF\mathcal{K}}(-M, -L)$ . For knots in  $(S^3, \xi_{\text{std}})$ ,  $\widehat{\mathcal{L}}(L)$  was identified (up to an automorphism of the ambient group) in [1] with a more easily calculable invariant defined in [23]; this latter invariant can be shown to vanish for any Legendrian figure-eight knot  $L$  (as  $\widehat{HF\mathcal{K}}(-S^3, -L)$  is trivial in the required grading). In [24], the element  $\widehat{\mathcal{L}}(L)$  was also identified with the class  $EH(\xi_{\text{std}}^-(L))$  of  $(S^3 \setminus N(K), \xi_{\text{std}}^-(L))$ , under an isomorphism  $\widehat{HF\mathcal{K}}(-S^3, -L) \cong SFH(-S^3 \setminus N(K), -\Gamma_{\text{meridional}})$ .

### 3. Surgeries on the figure-eight knot

Consider the figure-eight knot  $K$  in  $S^3$  (see Figure 5). We will show that the result of any positive contact surgery on any Legendrian realisation of the figure-eight knot in  $(S^3, \xi_{\text{std}})$  is overtwisted.




Figure 5: On the left is a smooth figure-eight knot  $K$ . On the right is a Legendrian representative  $L$  of  $K$  with  $tb(L) = \overline{tb}(K) = -3$ . We omit choices of orientation, since  $K$  is amphichiral.

Let  $L$  be a Legendrian figure-eight knot in  $(S^3, \xi_{\text{std}})$ . Define a contact structure  $\xi^-(L)$  (resp.  $\xi^+(L)$ ) on  $S^3 \setminus N(K)$  by gluing a negative (resp. positive) basic slice to the complement of  $N(L) \subset (S^3, \xi_{\text{std}})$  such that  $\partial(S^3 \setminus N(K))$  is convex with two meridional dividing curves.

**Proposition 3.1.** *Let  $L$  be a Legendrian figure-eight knot in  $(S^3, \xi_{\text{std}})$ .*

- 1) *If  $tb(L) - rot(L) = -3$  and  $tb(L) < -3$ , then  $(S^3 \setminus N(K), \xi^+(L))$  is overtwisted.*
- 2) *If  $tb(L) + rot(L) = -3$  and  $tb(L) < -3$ , then  $(S^3 \setminus N(K), \xi^-(L))$  is overtwisted.*
- 3) *If  $tb(L) \pm rot(L) < -3$ , then  $(S^3 \setminus N(K), \xi^\pm(L))$  is overtwisted.*

*Proof.* For any Legendrian knot  $L$ ,  $(S^3 \setminus N(K), \xi^-(L))$  is contactomorphic to  $(S^3 \setminus N(K), \xi^+(\bar{L}))$ , where  $\bar{L}$  is the mirror Legendrian knot to  $L$ . Since the figure-eight knot is amphichiral,  $\bar{L}$  is also a figure-eight knot, and  $rot(\bar{L}) = -rot(L)$ . Thus, (1) and (2) are equivalent. Also, if  $L$  satisfies  $tb(L) \pm rot(L) < -3$ , then so does  $\bar{L}$ , so to prove the proposition, it suffices to consider  $\xi^-(L)$  for  $L$  satisfying the hypotheses of (2) and (3).

By [8], the figure-eight knot is a Legendrian simple knot (*ie.* Legendrian figure-eight knots are classified up to isotopy by their  $tb$  and  $rot$ ) with  $tb(L) - rot(L) \leq -3$ . Thus, any  $L$  satisfying the hypotheses of (2) or (3) is a positive stabilisation of some other Legendrian knot  $L'$ . By the discussion in Section 2.3, gluing a positive basic slice (with appropriate slopes of dividing curves) to the complement of a standard neighbourhood of  $L'$  recovers the complement of a standard neighbourhood of  $L$ . Thus, gluing a negative basic slice to the complement of a standard neighbourhood of  $L$  — which constructs the contact structure  $\xi^-(L)$  on  $S^3 \setminus N(K)$  — is the same as first gluing a positive basic slice to the complement of a standard neighbourhood of  $L'$ , and then gluing on a further negative basic slice to get to  $\xi^-(L)$ . These two basic slices (the positive and the negative) glue together to give a single  $T^2 \times I$ , but since the two basic slices have opposite signs, the contact structure on this  $T^2 \times I$  is overtwisted (see the discussion after Theorem 2.7). This  $T^2 \times I$  embeds into  $(S^3 \setminus N(K), \xi^-(L))$ , so we conclude that  $(S^3 \setminus N(K), \xi^-(L))$  is overtwisted. □

Let  $L_t$  have  $tb(L_t) = t \leq -3$  and  $tb(L_t) - rot(L_t) = -3$ . The negative basic slice with dividing curve slopes  $-3$  and  $\infty$  can be divided into two negative basic slices, one with dividing curve slopes  $-3$  and  $t$ , and one with

dividing curve slopes  $t$  and  $\infty$ . Hence,

$$(S^3 \setminus N(L_t), \xi^-(L_t)) = (S^3 \setminus N(L_{-3}), \xi^-(L_{-3}))$$

for all  $t \leq -3$ . A similar statement holds for  $\xi^+$  for  $L$  satisfying  $tb(L) + rot(L) = -3$ . Additionally, as in the proof of Proposition 3.1, the amphichirality of the figure-eight knot gives a contactomorphism between  $\xi^-(L_{-3})$  and  $\xi^+(L_{-3})$ . Thus, to prove Theorem 1.1, it is sufficient to show that  $(S^3 \setminus N(K), \xi^-(L_{-3}))$  is overtwisted.

For the rest of this section, let  $L$  denote the Legendrian figure-eight knot in  $(S^3, \xi_{\text{std}})$  with  $tb(L) = -3$  (called  $L_{-3}$  above). Recall that the knot invariant  $\widehat{\mathcal{L}}(L)$  coming from Heegaard Floer vanishes for all Legendrian figure-eight knots, which implies that the contact invariant  $EH(\xi^-(L)) = 0$  as well (see Section 2.5).

**Proposition 3.2.** *All positive contact surgeries on  $L$  are overtwisted.*

*Sketch of Proof.* Assuming  $\xi$  is tight, we will use convex surfaces to show that  $(S^3 \setminus N(K), \xi^-(L))$  is contactomorphic to a unique contact manifold (see Lemma 3.3 and Lemma 3.4). We will then construct this contact manifold, and show that it has non-vanishing Heegaard Floer contact class  $EH$ . However, since  $\widehat{\mathcal{L}}(L) = 0$ , we know that  $EH(\xi^-(L))$  vanishes, and so we arrive at a contradiction, and  $(S^3 \setminus N(K), \xi^-(L))$  is overtwisted. We are then done, by the discussion preceding the proposition.  $\square$

Given a Seifert surface  $\Sigma$  for  $L$ , we can think of  $\Sigma$  as sitting inside  $S^3 \setminus N(K)$  with boundary on  $\partial(S^3 \setminus N(K))$ . After perturbing  $\Sigma$  to be convex, we first wish to normalise the dividing curves of  $\Sigma$  in  $(S^3 \setminus N(K), \xi^-(L))$ . We will use the fact that  $S^3 \setminus N(K)$  is fibred over  $S^1$  with fibre  $\Sigma$ , and the monodromy (after choosing a basis for  $\Sigma$ ) is given by

$$\phi = \begin{pmatrix} 2 & 1 \\ 1 & 1 \end{pmatrix}$$

up to twisting along the boundary of  $\Sigma$ ; choose the representative without any boundary twisting.

**Lemma 3.3.** *If  $(S^3 \setminus N(K), \xi^-(L))$  is tight, there is an isotopic copy of  $\Sigma$  in  $(S^3 \setminus N(K), \xi^-(L))$  such that it is convex and the dividing curves consist of a single boundary-parallel arc.*




Figure 6: Possible dividing curves on the annulus  $A$ . The tops are identified with the bottoms, and the left-hand side sits on  $\partial(S^3 \setminus N(K))$ .

*Proof.* During this proof, we will perturb  $\Sigma$  and swing it around the fibration to get new surfaces isotopic to  $\Sigma$ ; we will call each new copy  $\Sigma$ .

Etnyre and Honda showed in [8] that there exists a convex copy of  $\Sigma$  in the complement of  $N(L)$  with dividing curves consisting of three arcs, parallel to  $\begin{pmatrix} 0 \\ 1 \end{pmatrix}$ ,  $\begin{pmatrix} 1 \\ 1 \end{pmatrix}$ , and  $\begin{pmatrix} 1 \\ 2 \end{pmatrix}$ .

After gluing on a negative basic slice to get  $(S^3 \setminus N(K), \xi^-(L))$ , we extend  $\Sigma$  to the new boundary by gluing on an annulus  $A$  whose dividing curves are of one of the forms given in Figure 6, a translate of one of those forms (*ie.* the right-hand side endpoints are shifted up/down in the  $S^1$ -direction from what is shown in the figure), or the image of one of those forms in a power of a Dehn twist along the core of the annulus. Note that we have already excluded from our list of possibilities the cases where the dividing curves on  $A$  trace a boundary-parallel curve along  $\partial(S^3 \setminus N(K))$ . In these cases, the dividing curves on  $\Sigma$  would consist of a boundary-parallel curve and a contractible curve. Since we are assuming that  $(S^3 \setminus N(K), \xi^-(L))$  is tight, these cases would contradict Theorem 2.3.

In any of the remaining cases, the resulting dividing curves on  $\Sigma \cup A$  consist either of a single boundary-parallel arc or one non-boundary-parallel arc and one closed curve. We claim that the second case cannot occur. Indeed, we claim that a the relative Euler class acting on  $\Sigma \cup A$  must be non-zero, whereas it would be zero in the second case.

Indeed, in the second case, the dividing curves divide  $\Sigma \cup A$  into positive and negative regions that have the same Euler characteristic. Since the relative Euler class is the difference of these Euler characteristics, it must vanish. On the other hand, using the description of the dividing curves of  $\Sigma$ , it is straightforward to calculate that its relative Euler class vanishes (alternatively, this is the value of  $rot(L)$ , which is 0). Then, by the additivity of this invariant, the relative Euler class of  $\Sigma \cup A$  is equal to the relative Euler class of  $A$  in the basic slice.

To calculate this, we convert our basic slice to the standard picture described in Section 2.3. The matrix  $\begin{pmatrix} 1 & 0 \\ 2 & 1 \end{pmatrix} \in SL_2(\mathbb{Z})$  takes the slopes  $-\infty$  and  $-3$  of our basic slice to the standard slopes  $-\infty$  and  $-1$ , respectively. Since

$$\begin{pmatrix} 1 & 0 \\ 2 & 1 \end{pmatrix} \begin{pmatrix} 1 \\ 0 \end{pmatrix} = \begin{pmatrix} 1 \\ 2 \end{pmatrix} = \begin{pmatrix} 1 \\ 0 \end{pmatrix} - 2 \cdot \begin{pmatrix} 0 \\ -1 \end{pmatrix},$$

we see that the relative Euler class of the basic slice evaluated on  $A$  is  $\mp 2$ , which is non-zero, as claimed. □

**Lemma 3.4.** *Up to contactomorphism, there is at most one tight contact structure on  $S^3 \setminus N(K)$  inducing a convex boundary with two meridional dividing curves and such that there exists a copy of  $\Sigma$  with dividing curves of the form described in Lemma 3.3.*

*Proof.* First, we claim we can switch the signs of the regions  $\Sigma_{\pm}$  of  $\Sigma$ . Indeed, since  $\phi = (-id) \circ \phi \circ (-id)^{-1}$ , we can apply  $-id$  to  $\Sigma$ , which keeps the same dividing curves, but switches the signs of the regions.

Given  $\Sigma$  with fixed dividing curves  $\Gamma$  and signs of the regions  $\Sigma \setminus \Gamma$ , this uniquely determines a tight vertically-invariant contact structure on some neighbourhood  $N(\Sigma)$  of  $\Sigma$ . We will show that there exists a unique tight contact structure on  $M \setminus N(\Sigma)$ . Then, given two tight contact structures on  $M$  inducing the same dividing curves on  $\Sigma$  with the same signs, a contactomorphism of  $N(\Sigma)$  can be extended to a contactomorphism on all of  $M$ .

Observe that  $M \setminus N(\Sigma) \cong \Sigma \times [0, 1]$  is a genus 2 handlebody. The contact structure has a convex boundary obtained by rounding the edges of  $\Sigma \times \{i\}$  and  $\partial\Sigma \times [0, 1]$ , where the dividing curves on  $\Sigma \times \{0\}$  are  $\Gamma$ , those on  $\Sigma \times \{1\}$  are  $\phi(\Gamma) = \Gamma$  (since  $\Gamma$  is boundary-parallel), and those on  $\partial\Sigma \times [0, 1]$  are two copies of  $\{pt\} \times [0, 1]$ . We will look for compressing discs  $D_1$  and  $D_2$  such that their boundaries are Legendrian with  $tb = -1$ . After making the compressing discs convex, there will be a unique choice of dividing curves for  $D_i$ , since their dividing curves intersect the boundary of the disc at exactly two points, by Theorem 2.2, and there can be no contractible dividing




Figure 7: In each picture, the top and bottom are identified, as are the left and right sides. The dotted lines represent the dividing curves. The solid lines represent the intersection of the boundaries  $\partial D_i$  of the compressing discs with  $\Sigma \times \{0, 1\}$ .

curves, by Theorem 2.3. This allows us to uniquely define the tight contact structure in a neighbourhood of  $\partial(M \setminus N(\Sigma)) \cup D_1 \cup D_2$ . The complement of this neighbourhood is diffeomorphic to  $B^3$ , and by [4], we can uniquely extend the tight contact structure over  $B^3$ .

The dividing curves on  $\Sigma \times \{0, 1\}$  are shown as dotted lines in Figure 7. The compressing discs are shown as solid lines. Figure 8 shows the dividing curves in  $\partial\Sigma \times [0, 1]$ . As the curves  $\partial D_i$  pass from  $\Sigma \times \{0\}$  to  $\Sigma \times \{1\}$  through the region  $\partial\Sigma \times [0, 1]$ , they do not intersect any dividing curves, but they do switch which side of the dividing curves they are on. Thus,  $\partial D_i$  intersects the dividing curves exactly twice for each  $i = 0, 1$ , as required.  $\square$

We now construct this tight contact manifold, and show that the Heegaard Floer contact class is non-vanishing:

Consider the open book for  $S^3$  given by the figure-eight knot. The supported contact structure  $\xi_{ot}$  on  $S^3$  is overtwisted, but given any Legendrian approximation  $L'$  of the binding of the open book, it was shown in [9] that  $\hat{\mathcal{L}}(L')$  is non-vanishing. After gluing a negative basic slice to the complement of a standard neighbourhood of  $L'$ , we arrive at  $S^3 \setminus N(K)$  with contact structure  $\xi_{ot}^-(L')$  (in the language of Section 2.4). By the discussion in Section 2.5,




Figure 8: The left and right sides are identified in this picture. The dotted lines represent the dividing curves. The annulus in the middle is the region  $\partial\Sigma \times [0, 1]$ , and the darker regions above and below are interpolating regions representing how the dividing curves get connected while smoothing the boundary of  $M \setminus N(\Sigma)$ .

the fact that  $\widehat{\mathcal{L}}(L') \neq 0$  implies that  $\xi_{\text{ot}}^-(L')$  is tight, and that the Heegaard Floer contact class satisfies  $EH(\xi_{\text{ot}}^-(L')) \neq 0$ . It is also shown in [9] that in  $(S^3 \setminus N(K), \xi_{\text{ot}}^-(L'))$ , there is a copy of  $\Sigma$  (which is a page of the open book) that is convex, with dividing set consisting of one boundary-parallel arc. Thus, the unique contactomorphism class from Lemma 3.4 of type (2) has non-vanishing Heegaard Floer contact invariant.

*Proof of Theorem 1.1.* By Proposition 3.1 and the discussion below it, it suffices to consider the case  $tb(L) = -3$ . The result of any positive contact surgery on  $L$  has a contact submanifold that can be identified with  $(S^3 \setminus N(K), \xi^-(L))$  or  $(S^3 \setminus N(K), \xi^+(L))$ . The Heegaard Floer contact class  $EH(\xi^\pm(L))$  vanishes, as  $\widehat{\mathcal{L}}(L) = 0$  and  $L$  is amphichiral. Since if tight,  $\xi^-(L)$  and  $\xi^+(L)$  would have to be contactomorphic to the contact structure on  $S^3 \setminus N(K)$  constructed above with non-vanishing Heegaard Floer contact class, we conclude that  $\xi^-(L)$  and  $\xi^+(L)$  are overtwisted. Thus, any manifold which contains them as a contact submanifold must also be overtwisted.  $\square$

## References

- [1] John A. Baldwin, David Shea Vela-Vick, and Véra Vertesi, *On the equivalence of Legendrian and transverse invariants in knot Floer homology*, *Geometry & Topology* **17** (2013), 925–974.
- [2] James Conway, *Transverse surgery on knots in contact 3-manifolds*, to appear in *Trans. Amer. Math. Soc.* [arXiv:1409.7077v2 \[math.GT\]](https://arxiv.org/abs/1409.7077v2), (2014).
- [3] Fan Ding, Hansjörg Geiges, and András I. Stipsicz, *Surgery diagrams for contact 3-manifolds*, *Turkish J. Math.* **28** (2004), 41–74.
- [4] Yakov Eliashberg, *Contact 3-manifolds, twenty years since J. Martinet's work*, *Ann. Inst. Fourier (Grenoble)* **42** (1992), no. 1-2, 165–192.
- [5] John B. Etnyre, *Convex surfaces in contact geometry: Class notes*.
- [6] John B. Etnyre, *Introductory lectures on contact geometry*, *Proc. Sympos. Pure Math.* **71** (2003), 81–107.
- [7] John B. Etnyre, *Lectures on contact geometry in low-dimensional topology*, *IAS/Park City Math. Ser.* **15** (2009), 229–264.
- [8] John B. Etnyre and Ko Honda, *Knots and contact geometry I: Torus knots and the figure eight knot*, *J. Sympl. Geom.* **1** (2001), no. 1, 63–120.
- [9] John B. Etnyre and David Shea Vela-Vick, *Torsion and open book decompositions*, *IMRN* (2010), 4385–4398.
- [10] Emmanuel Giroux, *Convexité en topologie de contact*, *Comment. Math. Helv.* **66** (1991), no. 4, 637–677.
- [11] Marco Golla, *Ozsváth-Szabó invariants of contact surgeries*, *Geom. Topol.* **19** (2015), no. 1, 171–235.
- [12] Ko Honda, *On the classification of tight contact structures I*, *Geom. Topol.* **4** (2000), 309–368.
- [13] Ko Honda, William H. Kazez, and Gordana Matić, *The contact invariant in sutured Floer homology*, *Invent. Math.* **176** (2009), no. 3, 637–676.
- [14] Yutaka Kanda, *The classification of tight contact structures on the 3-torus*, *Comm. Anal. Geom.* **5** (1997), no. 3, 413–438.

- [15] Paolo Lisca, Peter Ozsváth, András I. Stipsicz, and Zoltán Szabó, *Heegaard Floer invariants of Legendrian knots in contact three-manifolds*, J. Eur. Math. Soc. **11** (2009), no. 6, 1307–1363.
- [16] Paolo Lisca and András I. Stipsicz, *Heegaard Floer invariants and tight contact three-manifolds*, [arXiv:0303280](https://arxiv.org/abs/0303280), (2003).
- [17] Paolo Lisca and András I. Stipsicz, *Ozsváth-Szabó invariants and tight contact three-manifolds, I*, Geom. Topol. **8** (2004), 925–945.
- [18] Paolo Lisca and András I. Stipsicz, *Notes on the contact Ozsváth-Szabó invariants*, Pacific J. Math. **228** (2006), no. 2, 277–295.
- [19] Paolo Lisca and András I. Stipsicz, *On the existence of tight contact structures on Seifert fibered 3-manifolds*, Duke Math. J. **148** (2009), no. 2, 175–209.
- [20] Paolo Lisca and András I. Stipsicz, *Contact surgery and transverse invariants*, J. Topol. **4** (2011), no. 4, 817–834.
- [21] Thomas E. Mark and Bülent Tosun, *Naturality of Heegaard Floer invariants under positive rational contact surgery*, J. Differential Geom. **110** (2018), no. 2, 281–344.
- [22] Peter Ozsváth and Zoltán Szabó, *Heegaard Floer homologies and contact structures*, Duke Math. J. **129** (2005), no. 1, 39–61.
- [23] Peter Ozsváth, Zoltán Szabó, and Dylan Thurston, *Legendrian knots, transverse knots and combinatorial floer homology*, Geometry & Topology **12** (2008), no. 2, 941–980.
- [24] András I. Stipsicz and Véra Vertesi, *On invariants for Legendrian knots*, Pacific J. Math. **239** (2009), no. 1, 157–177.
- [25] Andy Wand, *Tightness is preserved by Legendrian surgery*, Ann. Math. **182** (2015), no. 2, 723–738.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, BERKELEY  
970 EVANS HALL #3840 BERKELEY, CA 94720, USA  
*E-mail address*: [conway@berkeley.edu](mailto:conway@berkeley.edu)

RECEIVED NOVEMBER 14, 2016

ACCEPTED MARCH 19, 2018