

## DYNAMICS OF RATIONAL MAPS: A CURRENT ON THE BIFURCATION LOCUS

LAURA DEMARCO

ABSTRACT. Let  $f_\lambda : \mathbf{P}^1 \rightarrow \mathbf{P}^1$  be a family of rational maps of degree  $d > 1$ , parametrized holomorphically by  $\lambda$  in a complex manifold  $X$ . We show that there exists a canonical closed, positive (1,1)-current  $T$  on  $X$  supported exactly on the bifurcation locus  $B(f) \subset X$ . If  $X$  is a Stein manifold, then the stable regime  $X - B(f)$  is also Stein. In particular, each stable component in the space  $\text{Poly}_d$  (or  $\text{Rat}_d$ ) of all polynomials (or rational maps) of degree  $d$  is a domain of holomorphy.

### 1. Introduction

It is well-known that for a rational map  $f : \mathbf{P}^1 \rightarrow \mathbf{P}^1$  of degree  $d > 1$ , there is a natural  $f$ -invariant measure  $\mu_f$  supported on the Julia set of  $f$  [B],[Ly]. This measure can be described as the weak limit of purely atomic measures,

$$\mu_f = \lim_{n \rightarrow \infty} \frac{1}{d^n} \sum_{\{z: f^n(z)=a\}} \delta_z,$$

for any  $a \in \mathbf{P}^1$  (with at most two exceptions).

There is also a potential-theoretic description of  $\mu_f$ , defined in terms of a homogeneous polynomial lift  $F : \mathbf{C}^2 \rightarrow \mathbf{C}^2$  of  $f$ . The potential function on  $\mathbf{C}^2$  is given by

$$(1) \quad h(z) = \lim_{m \rightarrow \infty} \frac{1}{d^m} \log \|F^m(z)\|,$$

and the (1,1)-current  $\partial\bar{\partial}h$  satisfies

$$\pi^* \mu_f = \frac{i}{\pi} \partial\bar{\partial}h$$

where  $\pi$  is the canonical projection  $\mathbf{C}^2 - \{0\} \rightarrow \mathbf{P}^1$  [HP]. In particular, when  $f$  is a monic polynomial, this definition reduces to

$$\mu_f = \frac{i}{\pi} \partial\bar{\partial}G = \frac{1}{2\pi} \Delta G \, dx \wedge dy,$$

where  $G : \mathbf{C} \rightarrow [0, \infty)$  is the Green's function for the complement of the filled Julia set  $K(f) = \{z : f^n(z) \not\rightarrow \infty \text{ as } n \rightarrow \infty\}$ .

---

Received March 2, 2000.

In this paper, we construct a (1,1)-current on the **parameter space** of a holomorphic family of rational maps, supported exactly on the bifurcation locus (just as  $\mu_f$  is supported exactly on the Julia set).

Let  $X$  be a complex manifold. A **holomorphic family of rational maps**  $f$  **over**  $X$  is a holomorphic map  $f : X \times \mathbf{P}^1 \rightarrow \mathbf{P}^1$ . For each parameter  $\lambda \in X$ , we obtain a rational map  $f_\lambda : \mathbf{P}^1 \rightarrow \mathbf{P}^1$  with Julia set  $J(f_\lambda)$ . The **bifurcation locus**  $B(f)$  of the family  $f$  over  $X$  is the set of all  $\lambda_0 \in X$  for which  $\lambda \mapsto J(f_\lambda)$  is a discontinuous function (in the Hausdorff topology) in any neighborhood of  $\lambda_0$  (§2).

**Theorem 1.1.** *Let  $f : X \times \mathbf{P}^1 \rightarrow \mathbf{P}^1$  be a holomorphic family of rational maps on  $\mathbf{P}^1$  of degree  $d > 1$ . Then there exists a canonical closed, positive (1,1)-current  $T(f)$  on  $X$  such that the support of  $T(f)$  is  $B(f)$ , the bifurcation locus of  $f$ .*

By general properties of positive currents (Lemma 3.3), we have

**Corollary 1.2.** *If  $X$  is a Stein manifold, then  $X - B(f)$  is also Stein.*

Let  $\text{Rat}_d$  and  $\text{Poly}_d$  denote the “universal families” of all rational maps and of all monic polynomials of degree exactly  $d > 1$ . We have  $\text{Poly}_d \simeq \mathbf{C}^d$  and  $\text{Rat}_d \simeq \mathbf{P}^{2d+1} - V$ , where  $V$  is a resultant hypersurface. In particular,  $\text{Rat}_d$  and  $\text{Poly}_d$  are Stein manifolds.

**Corollary 1.3.** *Every stable component in  $\text{Rat}_d$  and  $\text{Poly}_d$  is a domain of holomorphy (i.e. a Stein open subset).*

Corollary 1.3 answers a question posed by McMullen in [M2], motivated by analogies between rational maps and Teichmüller space. Bers and Ehrenpreis showed that finite-dimensional Teichmüller spaces are domains of holomorphy [BE].

*Sketch proof of Theorem 1.1.* Consider a holomorphic family of homogeneous polynomial maps  $\{F_\lambda\}$  on  $\mathbf{C}^2$ , locally lifting the holomorphic family  $f$  over  $X$ . Let  $\{h_\lambda\}$  be the corresponding potential functions on  $\mathbf{C}^2$  defined by equation (1). The function  $h_\lambda(z)$  is plurisubharmonic in both  $\lambda \in X$  and  $z \in \mathbf{C}^2$ , and it is pluriharmonic in  $z$  away from  $\pi^{-1}(J(f_\lambda))$ . Suppose for simplicity that we have holomorphic functions  $c_j : X \rightarrow \mathbf{P}^1$ ,  $j = 1, \dots, 2d - 2$ , parametrizing the critical points of  $f_\lambda$  in  $\mathbf{P}^1$ . We choose lifts  $\tilde{c}_j$  from a neighborhood in  $X$  to  $\mathbf{C}^2$  so that  $c_j = \pi \circ \tilde{c}_j$  and define the plurisubharmonic function

$$H(\lambda) = \sum_j h_\lambda(\tilde{c}_j(\lambda)).$$

The desired (1,1)-current on  $X$  is defined by

$$T(f) = \frac{i}{\pi} \partial \bar{\partial} H,$$

independent of the choices of  $\{F_\lambda\}$  and  $\tilde{c}_j$ . It is supported on  $B(f)$  since  $H$  fails to be pluriharmonic exactly when a critical point  $c_j(\lambda)$  passes through the Julia set  $J(f_\lambda)$ .  $\square$

I would like to thank C. McMullen, J.E. Fornæss, and X. Buff for helpful comments and ideas.

## 2. Stability

Let  $f : X \times \mathbf{P}^1 \rightarrow \mathbf{P}^1$  be a holomorphic family of rational maps of degree  $d > 1$ . The Julia sets of such a family are said to **move holomorphically** at a point  $\lambda_0 \in X$  if there is a family of injections  $\phi_\lambda : J_{\lambda_0} \rightarrow \mathbf{P}^1$ , holomorphic in  $\lambda$  near  $\lambda_0$  with  $\phi_{\lambda_0} = \text{id}$ , such that  $\phi_\lambda(J_{\lambda_0}) = J_\lambda$  and  $\phi_\lambda \circ f_{\lambda_0}(z) = f_\lambda \circ \phi_\lambda(z)$ . In other words,  $\phi_\lambda$  provides a conjugacy between  $f_{\lambda_0}$  and  $f_\lambda$  on their Julia sets. The family of rational maps  $f$  over  $X$  is **stable** at  $\lambda_0 \in X$  if any of the following equivalent conditions are satisfied [M1, Theorem 4.2]:

- (1) The number of attracting cycles of  $f_\lambda$  is locally constant at  $\lambda_0$ .
- (2) The maximum period of an attracting cycle of  $f_\lambda$  is locally bounded at  $\lambda_0$ .
- (3) The Julia set moves holomorphically at  $\lambda_0$ .
- (4) For all  $\lambda$  sufficiently close to  $\lambda_0$ , every periodic point of  $f_\lambda$  is attracting, repelling, or persistently indifferent.
- (5) The Julia set  $J_\lambda$  depends continuously on  $\lambda$  (in the Hausdorff topology) in a neighborhood of  $\lambda_0$ .

Suppose also that each of the  $2d - 2$  critical points of  $f_\lambda$  are parametrized by holomorphic functions  $c_j : X \rightarrow \mathbf{P}^1$ . Then the following conditions are equivalent to those above:

- (6) For each  $j$ , the family of functions  $\{\lambda \mapsto f_\lambda^n(c_j(\lambda))\}_{n \geq 0}$  is normal in some neighborhood of  $\lambda_0$ .
- (7) For all nearby  $\lambda$ ,  $c_j(\lambda) \in J_\lambda$  if and only if  $c_j(\lambda_0) \in J_{\lambda_0}$ .

We let  $S(f) \subset X$  denote the set of stable parameters and define the **bifurcation locus**  $B(f)$  to be the complement  $X - S(f)$ . Mañé, Sad, and Sullivan showed that  $S(f)$  is open and dense in  $X$  [MSS, Theorem A].

**Example.** In the family  $f_c(z) = z^2 + c$ , the bifurcation locus is  $B(f) = \partial M$ , where  $M = \{c \in \mathbf{C} : f_c^n(0) \not\rightarrow \infty \text{ as } n \rightarrow \infty\}$  is the Mandelbrot set [M1, Theorem 4.6].

**Lemma 2.1.** *If  $B(f)$  is contained in a complex hypersurface  $D \subset X$ , then  $B(f)$  is empty.*

*Proof.* Suppose there exists  $\lambda_0 \in B(f)$ . By characterization (4) of stability, any neighborhood  $U$  of  $\lambda_0$  must contain a point  $\lambda_1$  at which the multiplier  $m(\lambda)$  of a periodic cycle for  $f_\lambda$  is passing through the unit circle. In other words, the holomorphic function  $m(\lambda)$  defined in a neighborhood  $N$  of  $\lambda_1$  is non-constant with  $|m(\lambda_1)| = 1$ . The set  $\{\lambda \in N : |m(\lambda)| = 1\}$  lies in the bifurcation locus and cannot be completely contained in a hypersurface.  $\square$

### 3. Stein manifolds and positive currents

Let  $X$  be a paracompact complex manifold and  $\mathcal{O}(X)$  its ring of holomorphic functions. Then  $X$  is a **Stein manifold** if the following three conditions are satisfied:

- for any  $x \in X$  there exists a neighborhood  $U$  of  $x$  and  $f_1, \dots, f_n \in \mathcal{O}(X)$  defining local coordinates on  $U$ ;
- for any  $x \neq y \in X$ , there exists an  $f \in \mathcal{O}(X)$  such that  $f(x) \neq f(y)$ ; and
- for any compact set  $K$  in  $X$ , the holomorphic hull

$$\hat{K} = \{x \in X : |f(x)| \leq \sup_K |f| \text{ for all } f \in \mathcal{O}(X)\}$$

is also compact in  $X$ .

An open domain  $\Omega$  in  $X$  is **locally Stein** if every boundary point  $p \in \partial\Omega$  has a neighborhood  $U$  such that  $U \cap \Omega$  is Stein.

**Properties of Stein manifolds.** The Stein manifolds are exactly those which can be embedded as closed complex submanifolds of  $\mathbf{C}^N$ . If  $\Omega$  is an open domain in  $\mathbf{C}^n$  then  $\Omega$  is Stein if and only if  $\Omega$  is pseudoconvex if and only if  $\Omega$  is a domain of holomorphy. An open domain in a Stein manifold is Stein if and only if it is locally Stein. Also, an open domain in complex projective space  $\mathbf{P}^n$  is Stein if and only if it is locally Stein and not all of  $\mathbf{P}^n$ . See, for example, [H] and the survey article by Siu [S].

**Examples.** (1)  $\mathbf{C}^N$  is Stein. (2) The space of all monic polynomials of degree  $d$ ,  $\text{Poly}_d \simeq \mathbf{C}^d$ , is Stein. (3)  $\mathbf{P}^n - V$  for a hypersurface  $V$  is Stein. If  $V$  is the zero locus of degree  $d$  homogeneous polynomial  $F$  and  $\{g_j\}$  a basis for the vector space of homogeneous polynomials of degree  $d$ , then the map  $(g_1/F, \dots, g_N/F)$  embeds  $\mathbf{P}^n - V$  as a closed complex submanifold of  $\mathbf{C}^N$ . (4) The space  $\text{Rat}_d$  of all rational maps  $f(z) = P(z)/Q(z)$  on  $\mathbf{P}^1$  of degree exactly  $d$  is Stein. Indeed, parameterizing  $f$  by the coefficients of  $P$  and  $Q$  defines an isomorphism  $\text{Rat}_d \simeq \mathbf{P}^{2d+1} - V$ , where  $V$  is the resultant hypersurface given by the condition  $\gcd(P, Q) \neq 1$ .

A  $(p, q)$ -**current**  $T$  on a complex manifold of dimension  $n$  is an element of the dual space to smooth  $(n-p, n-q)$ -forms with compact support. See [HP], [Le], and [GH] for details. The wedge product of a  $(p, q)$ -current  $T$  with any smooth  $(n-p, n-q)$ -form  $\alpha$  defines a distribution by  $(T \wedge \alpha)(f) = T(f\alpha)$  for  $f \in C_c^\infty(X)$ . Recall that a distribution  $\delta$  is positive if  $\delta(f) \geq 0$  for functions  $f \geq 0$ . A  $(p, p)$ -current is **positive** if for any system of  $n-p$  smooth  $(1, 0)$ -forms with compact support,  $\{\alpha_1, \dots, \alpha_{n-p}\}$ , the product

$$T \wedge (i\alpha_1 \wedge \bar{\alpha}_1) \wedge \dots \wedge (i\alpha_{n-p} \wedge \bar{\alpha}_{n-p})$$

is a positive distribution.

An upper-semicontinuous function  $h$  on a complex manifold  $X$  is **plurisubharmonic** if  $h|_{\mathbf{D}}$  is subharmonic for any complex analytic disk  $\mathbf{D}^1$  in  $X$ . The current  $T = i\partial\bar{\partial}h$  is positive for any plurisubharmonic  $h$ , and  $T \equiv 0$  if and only

if  $h$  is pluriharmonic. The “ $\partial\bar{\partial}$ -Poincaré Lemma” says that any closed, positive (1,1)-current  $T$  on a complex manifold is locally of the form  $i\partial\bar{\partial}h$  for some plurisubharmonic function  $h$  [GH].

The next three Lemmas show that the “region of pluriharmonicity” of a plurisubharmonic function is locally Stein. See [C, Theorem 6.2], [U, Lemma 2.4], [FS, Lemma 5.3], and [R, Theorem II.2.3] for similar statements.

**Lemma 3.1.** *Suppose  $h$  is plurisubharmonic on the open unit polydisk  $\mathbf{D}^2$  in  $\mathbf{C}^2$  and  $h$  is pluriharmonic on the “Hartogs domain”*

$$\Omega_\delta = \{(z, w) : |z| < 1, |w| < \delta\} \cup \{(z, w) : 1 - \delta < |z| < 1, |w| < 1\}.$$

*Then  $h$  is pluriharmonic on  $\mathbf{D}^2$ .*

*Proof.* Let  $H$  be a holomorphic function on  $\Omega_\delta$  such that  $h = \operatorname{Re} H$ . Any holomorphic function on  $\Omega_\delta$  extends to  $\mathbf{D}^2$ , and extending  $H$  we have  $h \leq \operatorname{Re} H$  on  $\mathbf{D}^2$  since  $h$  is plurisubharmonic. The set  $A = \{z \in \mathbf{D}^2 : h = \operatorname{Re} H\}$  is closed by upper-semi-continuity of  $h$ . If  $A$  has a boundary point  $w \in \mathbf{D}^2$ , then for any ball  $B(w)$  about  $w$ , we have

$$\begin{aligned} h(w) &= \operatorname{Re} H(w) \\ &= \frac{1}{|B(w)|} \int_{B(w)} \operatorname{Re} H \\ &> \frac{1}{|B(w)|} \int_{B(w)} h \end{aligned}$$

since  $\operatorname{Re} H > h$  on a set of positive measure in  $B(w)$ . This inequality, however, contradicts the sub-mean-value property of the subharmonic function  $h$ . Therefore  $A = \mathbf{D}^2$  and  $h$  is pluriharmonic on the polydisk.  $\square$

**Lemma 3.2.** *Let  $X$  be a complex manifold. If an open subset  $\Omega \subset X$  is not locally Stein, there is a  $\delta > 0$  and an embedding*

$$e : \mathbf{D}^2 \rightarrow X$$

*so that  $e(\Omega_\delta) \subset \Omega$  but  $e(\mathbf{D}^2) \not\subset \Omega$ .*

*Proof.* Suppose  $\Omega$  is not locally Stein at  $x \in \partial\Omega$ . By choosing local coordinates in a Stein neighborhood  $U$  of  $x$  in  $X$ , we may assume that  $U$  is a pseudoconvex domain in  $\mathbf{C}^n$ . Then  $\Omega_0 = U \cap \Omega$  is not pseudoconvex and the function  $\phi(z) = -\log d_0(z)$  is not plurisubharmonic near  $x \in \partial\Omega$ . Here,  $d_0$  is the Euclidean distance function to the boundary of  $\Omega_0$ .

If  $\phi$  is not plurisubharmonic at the point  $z_0 \in U \cap \Omega$ , then there is a one-dimensional disk  $\alpha : \mathbf{D}^1 \rightarrow \Omega$  centered at  $z_0$  such that  $\int_{\partial\mathbf{D}^1} \phi < \phi(z_0)$  (identifying the disk with its image  $\alpha(\mathbf{D}^1)$ ). Let  $\psi$  be a harmonic function on  $\mathbf{D}^1$  so that  $\psi = \phi$  on  $\partial\mathbf{D}^1$ . Then  $\psi(z_0) < \phi(z_0)$ . Let  $\Psi$  be a holomorphic function on  $\mathbf{D}^1$  with  $\psi = \operatorname{Re} \Psi$ .

Now, let  $p \in \partial\Omega$  be such that  $d_0(z_0) = |z_0 - p|$ . Let  $e : \mathbf{D}^2 \rightarrow U$  be given by

$$e(z_1, z_2) = \alpha(z_1) + z_2(1 - \varepsilon)e^{-\Psi(z_1)}(p - z_0).$$

That is, the two-dimensional polydisk is embedded so that at each point  $z_1 \in \mathbf{D}^1$  there is a disk of radius  $|(1 - \varepsilon) \exp(-\Psi(z_1))|$  in the direction of  $p - z_0$ . If  $\varepsilon$  is small enough we have a Hartogs-type subset of the polydisk contained in  $\Omega$  but the polydisk is not contained in  $\Omega$  since  $d_0(z_0, \partial\Omega) = \exp(-\phi(z_0)) < \exp(-\psi(z_0))$ .  $\square$

**Lemma 3.3.** *Let  $T$  be a closed, positive  $(1,1)$ -current on a complex manifold  $X$ . Then  $\Omega = X - \text{supp}(T)$  is locally Stein.*

*Proof.* Let  $p$  be a boundary point of  $\Omega$ . Choose a Stein neighborhood  $U$  of  $p$  in  $X$  so that  $T = i\partial\bar{\partial}h$  for some plurisubharmonic function  $h$  on  $U$ . By definition of  $\Omega$ ,  $h$  is pluriharmonic on  $U \cap \Omega$ .

If  $\Omega$  is not locally Stein at  $p$ , then by Lemma 3.2, we can embed a two-dimensional polydisk into  $U$  so that a Hartogs-type domain  $\Omega_\delta$  lies in  $\Omega$ , but the polydisk is not contained in  $\Omega$ . By Lemma 3.1,  $h$  must be pluriharmonic on the whole polydisk, contradicting the definition of  $\Omega$ .  $\square$

**Corollary 3.4.** *If  $X$  is Stein, then so is  $X - \text{supp } T$ .*

**Example.** If  $X$  is a Stein manifold and  $V$  a hypersurface, then  $V = \text{supp } T$  for a positive  $(1,1)$ -current  $T$  given locally by  $T = \frac{i}{\pi} \partial\bar{\partial} \log |f|$ , where  $V$  is the zero set of  $f$ . Lemma 3.3 shows that  $X - V$  is locally Stein, and thus Stein. Similarly,  $\mathbf{P}^n - V$  is Stein for any hypersurface  $V$ .

#### 4. The potential function of a rational map

Let  $f : \mathbf{P}^n \rightarrow \mathbf{P}^n$  be a holomorphic map. Let  $F : \mathbf{C}^{n+1} \rightarrow \mathbf{C}^{n+1}$  be a lift of  $f$  to a homogeneous polynomial, unique up to scalar multiple, so that  $\pi \circ F = f \circ \pi$  where  $\pi$  is the projection  $\mathbf{C}^{n+1} \setminus 0 \rightarrow \mathbf{P}^n$ . Let  $d$  be the degree of the components of  $F$ ; then  $f$  has topological degree  $d^n$ .

Assume that  $d > 1$ . Following [HP], we define the **potential function** of  $F$  by

$$h_F(z) = \lim_{m \rightarrow \infty} \frac{1}{d^m} \log \|F^m(z)\|.$$

The limit converges uniformly on compact subsets of  $\mathbf{C}^{n+1} - 0$ , and  $h_F(z)$  is plurisubharmonic on  $\mathbf{C}^{n+1}$  since  $\log \|\cdot\|$  is plurisubharmonic. Let  $\Omega_F \subset \mathbf{C}^{n+1}$  be the basin of attraction of the origin for  $F$ ; that is,

$$\Omega_F = \{x \in \mathbf{C}^{n+1} : F^m(x) \rightarrow 0 \text{ as } m \rightarrow \infty\}.$$

Note that  $\Omega_F$  is open and bounded.

From the definition, we obtain the following properties of the potential function  $h_F$  [HP]:

- (1)  $h_F(\alpha z) = h_F(z) + \log |\alpha|$  for  $\alpha \in \mathbf{C}^*$ ;
- (2)  $\Omega_F = \{z : h_F(z) < 0\}$ ; and
- (3)  $h_F$  is independent of the choice of norm  $\|\cdot\|$  on  $\mathbf{C}^{n+1}$ .

**Theorem 4.1.** (Hubbard-Papadopol, Ueda, Fornaess-Sibony) *The support of the positive  $(1,1)$ -current*

$$\omega_f = \frac{i}{\pi} \partial \bar{\partial} h_F$$

*on  $\mathbf{C}^{n+1} - 0$  is equal to the preimage of the Julia set  $\pi^{-1}(J(f))$ . If  $n = 1$ , then the Brodin-Lyubich measure  $\mu_f$  satisfies  $\pi^* \mu_f = \omega_f$ .*

*Proof.* See [HP, Theorem 4.1] for  $n = 1$  and [U, Theorem 2.2], [FS, Theorem 2.12] for  $n > 1$ .  $\square$

From Corollary 3.4, we obtain the following ([U, Theorem 2.3], [FS, Theorem 5.2]):

**Corollary 4.2.** (Ueda, Fornaess-Sibony) *The Fatou components of  $f : \mathbf{P}^n \rightarrow \mathbf{P}^n$  are Stein.*

## 5. The bifurcation current

In this section we complete the proof of Theorem 1.1. Let  $f : X \times \mathbf{P}^1 \rightarrow \mathbf{P}^1$  be a holomorphic family of rational maps on  $\mathbf{P}^1$  of degree  $d > 1$ . Let  $\{F_\lambda\}$  be a holomorphic family of homogeneous polynomials on  $\mathbf{C}^2$ , locally lifting the family  $f$ , and let  $h_\lambda$  denote the potential function of  $F_\lambda$  (§4). The potential function  $h_\lambda(z)$  is plurisubharmonic as a function of the pair  $(\lambda, z)$ .

Fix  $\lambda_0 \in X$ . In a neighborhood  $U$  of  $\lambda_0$ , we can choose coordinates on  $\mathbf{P}^1$  so that  $\infty$  is not a critical point of  $f_\lambda$ ,  $\lambda \in U$ . For  $z \in \mathbf{P}^1 - \{\infty\}$ , let  $\tilde{z} = (z, 1) \in \mathbf{C}^2$ . Define a function  $H$  on  $U$  by

$$H(\lambda) = \sum_{\{c: f'_\lambda(c)=0\}} h_\lambda(\tilde{c}),$$

where the critical points are counted with multiplicity. Now, let  $N(\lambda)$  be the number of critical points of the rational map  $f_\lambda$  (counted without multiplicity). Let

$$D(f) = \{\lambda_0 \in X : N(\lambda) \text{ does not have a local maximum at } \lambda = \lambda_0\}.$$

Then  $D(f)$  is a complex hypersurface in  $X$ , since it is defined by the vanishing of a discriminant. If  $\lambda_0 \notin D(f)$ , there exists a neighborhood  $U$  of  $\lambda_0$  and holomorphic functions  $c_j : U \rightarrow \mathbf{P}^1$ ,  $j = 1, \dots, 2d-2$ , parametrizing the critical points of  $f_\lambda$ , such that  $\infty \notin c_j(U)$  for all  $j$ . In this case, we can express  $H$  as the sum

$$H(\lambda) = \sum_j H_j(\lambda)$$

of the plurisubharmonic functions

$$\begin{aligned} H_j(\lambda) &= h_\lambda \circ \tilde{c}_j(\lambda) \\ &= \lim_{m \rightarrow \infty} \frac{1}{d^m} \log \|F_\lambda^m(\tilde{c}_j(\lambda))\|. \end{aligned}$$

For any  $\lambda_0 \in X$ , then,  $H$  is defined and continuous in a neighborhood  $U$  of  $\lambda_0$  and plurisubharmonic on  $U - D(f)$ ; therefore  $H$  is plurisubharmonic on  $U$ .

The **bifurcation current**  $T$  is the positive (1,1)-current on parameter space  $X$  given locally by

$$T = \frac{i}{\pi} \partial \bar{\partial} H.$$

The next Lemma shows that  $T$  is globally well-defined on  $X$ .

**Lemma 5.1.** *The current  $T = \frac{i}{\pi} \partial \bar{\partial} H$  is independent of (a) the choice of lifts  $\tilde{c}_j$  of  $c_j$  and (b) the choice of lifts  $F_\lambda$  of  $f_\lambda$ .*

*Proof.* Suppose we define a new lift  $\hat{c}_j(\lambda) = t(\lambda) \cdot \tilde{c}(\lambda)$  for some holomorphic function  $t$  taking values in  $\mathbf{C}^*$ . Property (1) of the potential function  $h_\lambda$  (§4) implies that  $h_\lambda(\hat{c}(\lambda)) = h_\lambda(\tilde{c}(\lambda)) + \log |t(\lambda)|$  and  $i\partial\bar{\partial}H$  is unchanged since  $\log |t(\lambda)|$  is pluriharmonic, proving (a). If the lifted family  $\{F_\lambda\}$  is similarly replaced by  $\{t(\lambda) \cdot F_\lambda\}$ , a computation shows that  $h_\lambda$  is changed only by the addition of the pluriharmonic term  $\frac{1}{d-1} \log |t(\lambda)|$  where  $d$  is the degree of the  $f_\lambda$ . This proves (b).  $\square$

**Lemma 5.2.** *A parameter  $\lambda_0$  lies in the stable regime  $S(f) \subset X$  if and only if the function  $H$  is pluriharmonic in a neighborhood of  $\lambda_0$ .*

*Proof.* Let us first suppose that  $\lambda_0 \in S(f)$  is not in  $D(f)$  (in the notation above). By characterization (6) of stability (§2), for each  $j$ , the family of functions  $\{\lambda \mapsto f_\lambda^m(c_j(\lambda))\}$  is normal in a neighborhood  $V$  of  $\lambda_0$ ; hence, there exists a subsequence converging uniformly on compact subsets to a holomorphic function  $g_j(\lambda)$ . As in [HP, Prop 5.4], we can shrink our neighborhood  $V$  if necessary to find a norm  $\|\cdot\|$  on  $\mathbf{C}^2$  so that  $\log \|\cdot\|$  is pluriharmonic on  $\pi^{-1}(g_j(V))$ ; e.g., if  $g_j(V)$  is disjoint from  $\{|x| = |y|\}$ , we can choose norm  $\|(x, y)\| = \max\{|x|, |y|\}$ . Then, on any compact set in  $V$ , the functions

$$\lambda \mapsto \frac{1}{d^{m_k}} \log \|F_\lambda^{m_k}(\tilde{c}_j(\lambda))\|$$

are pluriharmonic if  $k$  is large enough. By property (3) of the potential function  $h_\lambda$  (§4), this subsequence converges uniformly to  $H_j$ . Therefore,  $H$  is pluriharmonic on  $V$ .

If  $\lambda_0$  lies in  $D(f) \cap S(f)$ , then  $H$  is defined and continuous on a neighborhood  $V$  of  $\lambda_0$  and pluriharmonic on  $V - D(f)$ . As  $D(f)$  has codimension 1,  $H$  must be pluriharmonic on all of  $V$ .

For the converse, let us suppose again that  $\lambda_0 \notin D(f)$  and that  $H$  is pluriharmonic in a neighborhood of  $\lambda_0$ . Each  $H_j$  is pluriharmonic and so we may write  $H_j = \operatorname{Re} G_j$  in a neighborhood  $V$  of  $\lambda_0$ . In analogy with [U, Prop. 2.1], we define new lifts  $\hat{c}_j(\lambda) = e^{-G_j(\lambda)} \cdot \tilde{c}(\lambda)$  of the  $c_j$  and compute

$$\begin{aligned} h_\lambda(\hat{c}_j(\lambda)) &= h_\lambda(\tilde{c}(\lambda)) + \log |e^{-G_j(\lambda)}| \\ &= h_\lambda(\tilde{c}(\lambda)) - \operatorname{Re} G_j \\ &= H_j - H_j \\ &= 0. \end{aligned}$$



By property (2) of  $h_\lambda$ , this implies that  $\hat{c}_j(\lambda)$  lies in  $\partial\Omega_\lambda$  for all  $\lambda \in V$ . If  $V$  is small enough, the set  $\cup_{\lambda \in V}(\{\lambda\} \times \partial\Omega_\lambda)$  has compact closure in  $X \times \mathbf{C}^2$ . As the functions  $F_\lambda$  preserve  $\partial\Omega_\lambda$ , the family  $\{\lambda \mapsto F_\lambda^n(\hat{c}_j(\lambda))\}$  is uniformly bounded and thus normal. Of course,  $f_\lambda^n \circ c_j = \pi \circ F_\lambda^n \circ \hat{c}_j$  demonstrating that  $\lambda_0$  is a stable parameter by (6) of Section 2.

Finally suppose that  $H$  is pluriharmonic in a neighborhood  $U$  of parameter  $\lambda_0 \in D(f)$ . Then  $U - D(f)$  lies in the stable regime and Lemma 2.1 shows that all of  $U$  must belong to  $S(f)$ .  $\square$

*Proof of Theorem 1.1.* Let  $T$  be the bifurcation current defined above for the family of rational maps  $f$  over  $X$ . By Lemma 5.2, the support of  $T$  is the bifurcation locus  $B(f)$ .  $\square$

Corollaries 1.2 and 1.3 now follow immediately from Corollary 3.4.

## 6. Examples

**Example 6.1.** In the family  $\{f_\lambda(z) = z^d + \lambda\}$ ,  $\lambda \in \mathbf{C}$ , the bifurcation current  $T$  takes the form

$$T = \frac{d-1}{d} \left( \frac{i}{\pi} \partial \bar{\partial} G \right)$$

where  $G$  is the Green's function for the complement of the “degree  $d$  Mandelbrot set”  $M_d = \{\lambda : f_\lambda^n(0) \not\rightarrow \infty \text{ as } n \rightarrow \infty\}$ . That is,  $T$  is a multiple of harmonic measure supported on  $\partial M_d$ . The  $T$ -mass of  $\partial M_d$  is  $(d-1)/d$ .

*Proof.* If  $G_\lambda$  denotes the Green's function for the complement of the filled Julia set  $K(f_\lambda) = \{z : f_\lambda^n(z) \not\rightarrow \infty \text{ as } n \rightarrow \infty\}$ , then  $G(\lambda) = G_\lambda(\lambda)$  (see e.g. [CG, VIII.4]). By [HP, Prop 8.1], we have

$$h_\lambda(x, y) = G_\lambda(x/y) + \log |y|$$

where  $(x, y)$ ,  $y \neq 0$ , is a point of  $\mathbf{C}^2$ . Note that  $d-1$  of the critical points of  $f_\lambda$  are at  $z = 0$  and the other  $d-1$  are at  $z = \infty$ . Computing, we find

$$\begin{aligned} T &= \frac{i}{\pi} \sum_j \partial \bar{\partial} h_\lambda(\tilde{c}_j(\lambda)) \\ &= (d-1) \frac{i}{\pi} \partial \bar{\partial} h_\lambda(0, 1) \\ &= (d-1) \frac{i}{\pi} \partial \bar{\partial} G_\lambda(0) \\ &= \frac{d-1}{d} \frac{i}{\pi} \partial \bar{\partial} G_\lambda(\lambda). \end{aligned}$$

$\square$

**Example 6.2.** Let  $f$  be a polynomial of degree  $d$  and  $G_f$  the Green's function for the complement of the filled Julia set. The Lyapunov exponent of  $f$  (for the Brolin-Lyubich measure) satisfies ([Prz],[Mn])

$$L(f) = \log d + \sum_{\{c \in \mathbf{C} : f'(c)=0\}} G_f(c).$$

If  $\{f_\lambda\}$  is any holomorphic family of polynomials, the Lyapunov exponent as a function of the parameter is a potential function for the bifurcation current; that is,

$$T = \frac{i}{\pi} \partial \bar{\partial} L.$$

In the sequel, we examine further the connection between the bifurcation current and the Lyapunov exponent.

### References

- [BE] L. Bers and L. Ehrenpreis, *Holomorphic convexity of Teichmüller spaces*, Bull. Amer. Math. Soc. **70** (1964), 761–764.
- [B] H. Brolin, *Invariant sets under iteration of rational functions*, Ark. Mat. **6** (1965), 103–144.
- [CG] L. Carleson and T. W. Gamelin, *Complex dynamics*, Springer-Verlag, New York, 1993.
- [C] U. Cegrell, *Plurisubharmonic functions*, Proc. London Math. Soc. **36** (1978), 310–336.
- [FS] J. E. Fornæss and N. Sibony, *Complex dynamics in higher dimension. II*. Modern methods in complex analysis (Princeton, NJ, 1992), Princeton Univ. Press, Princeton, NJ, 1995, pp. 135–182.
- [GH] P. Griffiths and J. Harris, *Principles of algebraic geometry*, John Wiley & Sons Inc., New York, 1994.
- [H] L. Hörmander, *An introduction to complex analysis in several variables*, North-Holland Publishing Co., Amsterdam, 1990.
- [HP] J. Hubbard and P. Papadopol, *Superattractive fixed points in  $\mathbf{C}^n$* , Indiana Univ. Math. J. **43** (1994), 321–365.
- [Le] P. Lelong, *Fonctions plurisousharmoniques et formes différentielles positives*, Gordon & Breach, Paris, 1968.
- [Ly] M. Lyubich, *Entropy properties of rational endomorphisms of the Riemann sphere*, Ergodic Theory Dynamical Systems **3** (1983), 351–385.
- [Mn] R. Mañé, *The Hausdorff dimension of invariant probabilities of rational maps*, Dynamical systems, Valparaiso 1986, Lecture Notes in Math. **1331**, Springer, Berlin, 1988, pp. 86–117.
- [MSS] R. Mañé, P. Sad, and D. Sullivan, *On the dynamics of rational maps*, Ann. Sci. cole Norm. Sup. **16** (1983), 193–217.
- [M1] C. McMullen, *Complex dynamics and renormalization*, Annals of Mathematics Studies **135**, Princeton University Press, Princeton, NJ, 1994.
- [M2] ———, *Rational maps and Teichmüller space*, Lecture Notes in Mathematics **1574**, Springer-Verlag, 1994, pp. 430–433.
- [Prz] F. Przytycki, *Hausdorff dimension of harmonic measure on the boundary of an attractive basin for a holomorphic map*, Invent. Math. **80** (1985), 161–179.
- [R] R. M. Range, *Holomorphic functions and integral representations in several complex variables*, Graduate Texts in Mathematics **108**, Springer-Verlag, New York, 1986.
- [S] Y.-T. Siu, *Pseudoconvexity and the Problem of Levi*, Bull. Amer. Math. Soc. **84** (1978), 481–512.
- [U] T. Ueda, *Fatou sets in complex dynamics on projective spaces*, J. Math. Soc. Japan **46** (1994), 545–555.

DEPARTMENT OF MATHEMATICS, HARVARD UNIVERSITY, ONE OXFORD STREET CAMBRIDGE, MA 02138.

*E-mail address:* demarco@math.harvard.edu