

COMPONENT GROUPS OF PURELY TORIC QUOTIENTS

BRIAN CONRAD AND WILLIAM A. STEIN

ABSTRACT. Suppose $\pi : J \rightarrow A$ is an optimal quotient of abelian varieties over a p -adic field, optimal in the sense that $\ker(\pi)$ is connected. Assume that J is equipped with a symmetric principal polarization θ (e.g., any Jacobian of a curve has such a polarization), that J has semistable reduction, and that A has purely toric reduction. In this paper, we express the group of connected components of the Néron model of A in terms of the monodromy pairing on the character group of the torus associated to J . We apply our results in the case when A is an optimal quotient of the modular Jacobian $J_0(N)$. For each prime p that exactly divides N , we obtain an algorithm to compute the order of the component group of A at p .

1. Introduction

Let A be an abelian variety over the rational numbers \mathbf{Q} . Birch and Swinnerton-Dyer found a conjectural formula for the order of the Shafarevich-Tate group of A . The Tamagawa numbers c_p of A are among the quantities that appear in this formula. We now recall the definition of the Tamagawa numbers of an abelian variety (the definition of Néron model and component groups is given in Section 2).

Definition 1.1 (Tamagawa number). Let p be a prime, let \mathcal{A} be the Néron model of A over the p -adic integers \mathbf{Z}_p , and let $\Phi_{A,p}$ be the component group of \mathcal{A} at p . Then the *Tamagawa number* c_p of A at p is the order of the subgroup $\Phi_{A,p}(\mathbf{F}_p)$ of \mathbf{F}_p -rational points in $\Phi_{A,p}(\overline{\mathbf{F}}_p)$.

Remark 1.2. The Tamagawa number is defined in a different way in some other papers, but the definitions are equivalent.

When A has dimension one, A is called an elliptic curve, and A can be defined by a Weierstrass equation $y^2 = x^3 + ax + b$. Using that elliptic curves (and their related integral models) can be described by simple equations, Tate found an efficient algorithm to compute all of the Tamagawa numbers of A (see [18]). In the case when A is the Jacobian of a genus 2 curve, [7] discusses a method for computing the Tamagawa numbers of A . In this paper, we consider the situation in which A has purely toric reduction at p , with no constraint on the dimension of A . For such A we give an explicit description of the order of the group of connected components of the closed fiber of the Néron model

Received April 20, 2001.

Both authors were partially supported by the NSF and the Clay Mathematics Institute.

of A . In the case when $A = A_f$ is a quotient of $J_0(N)$ attached to a newform $f \in S_2(\Gamma_0(N))$ and $p \nmid N$, our method is completely explicit, and yields an algorithm to compute the Tamagawa number c_p of A (up to a bounded power of 2).

This paper is structured as follows. In Sections 2–6 we state and prove an explicit formula involving component groups of fairly general abelian varieties. Then in Section 7 we turn to quotients of modular Jacobians $J_0(N)$. We give some tables and discuss the arithmetic of quotients of $J_0(N)$ when N is prime. In Section 8 we prove a couple of facts about toric reduction that are used in the proof of Theorem 6.1.

2. The Main Results

In this section, we summarize the main contributions of this paper. First we recall the precise definition of the component group of an abelian variety, then we state our main theorem.

Let R be a discrete valuation ring with field of fractions K and maximal ideal \mathfrak{m} , and let $k = R/\mathfrak{m}$ be the residue class field. Let A be an abelian variety over K .

Definition 2.1 (Néron model). A *Néron model* of A is a smooth commutative group scheme \mathcal{A} over R such that A is its generic fiber and \mathcal{A} satisfies the Néron mapping property: the restriction map

$$\mathrm{Hom}_R(S, \mathcal{A}) \longrightarrow \mathrm{Hom}_K(S_K, A)$$

is bijective for all smooth schemes S over R .

The Néron mapping property implies that \mathcal{A} is unique up to a unique isomorphism, so we will refer without hesitation to “the” Néron model of A . Néron models are separated and of finite type as opposed to just locally of finite type, even though their universal property is on the category of arbitrary smooth R -schemes. For more about Néron models see [2].

The closed fiber \mathcal{A}_k of \mathcal{A} is a group scheme over k , which need not be connected. Denote by \mathcal{A}_k^0 the connected component of \mathcal{A}_k that contains the identity. We have an exact sequence

$$0 \longrightarrow \mathcal{A}_k^0 \longrightarrow \mathcal{A}_k \longrightarrow \Phi_A \longrightarrow 0,$$

where Φ_A is a finite étale group scheme over k . Equivalently, Φ_A is a commutative finite group equipped with a continuous action of $\mathrm{Gal}(\bar{k}/k)$.

Definition 2.2 (Component group). The *component group* of an abelian variety A over K is the group scheme $\Phi_A = \mathcal{A}_k/\mathcal{A}_k^0$.

2.1. Statement of the Theorem. We now state our main result, suppressing some of the definitions of the terms used until later (see Section 6 below for a more complete statement and the proof). Let K be as above, and suppose $\pi : J \rightarrow A$ is an optimal quotient. Assume that J is equipped with a symmetric principal polarization λ , in the sense of Definition 5.1. For example, the θ polarization of the Jacobian of a curve is a symmetric principal polarization. Also assume that J has semistable reduction, and that A has purely toric reduction.

We express the component group of A in terms of the monodromy pairing associated to J . Let $m_A = \sqrt{\deg(\theta_A)}$, where $\theta_A : A^\vee \rightarrow A$ is induced by the principal polarization λ . Let X_J be the character group of the toric part of the closed fiber of the Néron model of J . Let \mathcal{L} be the saturation of the image of X_A in X_J . The monodromy pairing induces a map $\alpha : X_J \rightarrow \text{Hom}(\mathcal{L}, \mathbf{Z})$. Let Φ_X be the cokernel of α and $m_X = [\alpha(X_J) : \alpha(\mathcal{L})]$ be the order of the finite group $\alpha(X_J)/\alpha(\mathcal{L})$. The main result of this paper is that

$$\frac{\#\Phi_A}{m_A} = \frac{\#\Phi_X}{m_X},$$

and this is recorded as Theorem 6.1 below.

Using the snake lemma, one sees that Φ_X is isomorphic to the image of the natural map $\Phi_J \rightarrow \Phi_A$, and the above formula implies that the cokernel of the map $\Phi_J \rightarrow \Phi_A$ has order m_A/m_X . A non-obvious consequence of this is that $m_X \mid m_A$.

In the context of modular forms, if the optimal quotient $J \rightarrow A$ arises from a newform on $\Gamma_0(N)$, then the quantities m_A , m_X and Φ_X can be explicitly computed, hence we can compute $\#\Phi_A$. Note that the authors have not computed the structure of Φ_A as a group.

3. Optimal Quotients

Let K be as in Section 2, let J be an abelian variety equipped with a symmetric principal polarization θ_J (see Definition 5.1). For example, J could be the Jacobian of a curve equipped with the canonical principal polarization arising from the θ -divisor.

Definition 3.1 (Optimal quotient). An *optimal quotient* of J is an abelian variety A and a smooth surjective morphism $\pi : J \rightarrow A$ whose kernel is connected (i.e., an abelian variety).

Remark 3.2. Any connected scheme of finite type over a field is geometrically connected if it contains a rational point (e.g., if it is a group scheme). See [8, IV₂, §4.5.13].

Let $\pi : J \rightarrow A$ be an optimal quotient. Denote by J^\vee and A^\vee the abelian varieties dual to J and A , respectively. Upon composing the dual of π with $\theta_J^\vee = \theta_J$, we obtain a map

$$A^\vee \xrightarrow{\pi^\vee} J^\vee \xrightarrow{\theta_J} J.$$

Proposition 3.3. *The map $\theta_J \circ \pi^\vee : A^\vee \rightarrow J$ is a closed immersion.*

Proof. Since θ_J is an isomorphism, we want to prove that π^\vee is a closed immersion. It is a general fact that duals to surjections of abelian varieties with abelian variety kernel are closed immersions, but for lack of an adequate reference we recall the proof. Since a monomorphism between smooth finite type group schemes over a field is necessarily a closed immersion, it suffices to show that the commutative proper group scheme $\ker(\pi^\vee)$ vanishes. Since a non-zero commutative proper group scheme G over a field F necessarily has a non-zero finite subgroup scheme $G[n]$ for some n (since either $(G/\overline{F})_{\text{red}}^0$ is an abelian variety or else G is finite and non-zero), it suffices to show that $\ker(\pi^\vee)[n]$ vanishes for all positive integers n . In other words, it suffices to show that the induced map $A^\vee[n] \rightarrow J^\vee[n]$ is a closed immersion for all n .

Since Cartier duality interchanges faithfully flat maps and closed immersions, and the scheme-theoretic Weil pairing identifies the Cartier dual of the map induced by π^\vee on n -torsion with $\pi : J[n] \rightarrow A[n]$, we just have to show that these latter maps are faithfully flat for all integers n . Using the short exact sequence

$$0 \rightarrow \ker(\pi) \rightarrow J \rightarrow A \rightarrow 0$$

in the abelian category of fppf abelian sheaves over $\text{Spec}(K)$, the snake lemma gives an exact sequence

$$0 \rightarrow \ker(\pi)[n] \rightarrow J[n] \rightarrow A[n] \rightarrow 0$$

because $n : \ker(\pi) \rightarrow \ker(\pi)$ is a faithfully flat map (hence fppf surjective), as $\ker(\pi)$ is an abelian variety. This gives an isomorphism of group schemes

$$J[n]/\ker(\pi)[n] \simeq A[n]$$

compatible with the maps from $J[n]$, whence $J[n] \rightarrow A[n]$ is faithfully flat. \square

Henceforth we will abuse notation and denote the injection $A^\vee \rightarrow J$ by π^\vee . We define θ_A to be the composite $\pi \circ \pi^\vee$, so the kernel of θ_A equals the scheme-theoretic intersection of A^\vee and $B = \ker(\pi)$, as depicted in the following diagram:

$$\begin{array}{ccc} A^\vee \cap B & \longrightarrow & B \\ \downarrow & & \downarrow \\ A^\vee & \xrightarrow{\pi^\vee} & J \\ & \searrow \theta_A & \downarrow \pi \\ & & A. \end{array}$$

Since θ_A is a polarization (due to how its definition uses the polarization θ_J) the degree of θ_A is a perfect square (see [16, §16, p. 150]).

Definition 3.4 (Degree). Define the *degree* of A as a quotient of J to be the integer

$$m_A = \sqrt{\#\ker(\theta_A)}.$$

4. The Closed Fiber of the Néron Model

In this section we recall some terminology associated with closed fibers of Néron models. Let K , R , and k be as in Section 2, and let $\Phi_A = \mathcal{A}_k / \mathcal{A}_k^0$ be the group scheme of connected components of the closed fiber \mathcal{A}_k . By Chevalley's structure theorem (see [3], or [4] for a modern account), if K is a perfect extension field of k (e.g., $K = \bar{k}$) then there is a unique short exact sequence

$$0 \rightarrow \mathcal{C} \rightarrow \mathcal{A}_K^0 \rightarrow \mathcal{B} \rightarrow 0$$

with \mathcal{C} a smooth affine algebraic K -group and \mathcal{B} an abelian variety. Moreover, there is a unique exact sequence

$$0 \rightarrow \mathcal{T} \rightarrow \mathcal{C} \rightarrow \mathcal{U} \rightarrow 0$$

with \mathcal{T} a torus and \mathcal{U} unipotent.

Using the rigidity of tori, one can show that \mathcal{T} is induced by a unique torus in \mathcal{A}_k^0 . In particular, the condition that $\mathcal{B} = \mathcal{U} = 0$ is equivalent to the condition that \mathcal{A}_k^0 be a torus, and the condition that $\mathcal{U} = 0$ is equivalent to the condition that \mathcal{A}_k^0 be the extension of an abelian variety by a torus (i.e., be a semi-abelian variety). These conditions can be checked on a geometric closed fiber.

Definition 4.1. The abelian variety A is said to have *purely toric reduction* if \mathcal{A}_k^0 is torus, and to have *semistable reduction* if \mathcal{A}_k^0 is a semi-abelian variety (i.e., \mathcal{A}_k^0 has vanishing unipotent part).

4.1. The Monodromy Pairing on the Character Group.

Definition 4.2 (Character group of torus). The *character group*

$$X_A = \mathrm{Hom}_{\bar{k}}(\mathcal{T}_{\bar{k}}, \mathbf{G}_{m\bar{k}})$$

is a free abelian group contravariantly associated to A .

As discussed in [9], if A is semistable there is a *monodromy pairing* $X_A \times X_{A^\vee} \rightarrow \mathbf{Z}$ and an exact sequence

$$0 \rightarrow X_{A^\vee} \rightarrow \mathrm{Hom}(X_A, \mathbf{Z}) \rightarrow \Phi_A \rightarrow 0.$$

Also, the canonical isomorphism $(A^\vee)^\vee \cong A$ induces an isomorphism

$$X_{A^\vee} \times X_{(A^\vee)^\vee} \cong X_A \times X_{A^\vee},$$

which identifies the monodromy pairing associated to A^\vee with that associated to A .

Example 4.3 (Tate curve). Suppose $E = \mathbf{G}_m/q^{\mathbf{Z}}$ is a Tate curve over $\mathbf{Q}_p^{\mathrm{ur}}$. The monodromy pairing on $X_E = q^{\mathbf{Z}}$ is

$$\langle q, q \rangle = \mathrm{ord}_p(q) = -\mathrm{ord}_p(j).$$

Thus Φ_E is cyclic of order $-\mathrm{ord}_p(j)$.

Suppose J is an abelian variety equipped with a symmetric principal polarization. Since J is self dual via the given symmetric principal polarization, we can view the monodromy pairing on J as a pairing $X_J \times X_J \rightarrow \mathbf{Z}$. Because the principal polarization on J is symmetric the resulting pairing $X_J \times X_J \rightarrow \mathbf{Z}$ is symmetric, so there is no ambiguity about left versus right definitions of $X_J \rightarrow \text{Hom}(X_J, \mathbf{Z})$. The above exact sequence then becomes

$$0 \rightarrow X_J \rightarrow \text{Hom}(X_J, \mathbf{Z}) \rightarrow \Phi_J \rightarrow 0.$$

5. The Degree of a Symmetric Isogeny

We next relate the degree of the isogeny $A^\vee \rightarrow A$ defined at the end of Section 3 to the order of the cokernel of the induced map on the character groups of tori defined in Section 4.1. Let K be as in Section 2, and let A be an abelian variety over K .

Definition 5.1 (Symmetric isogeny). A *symmetric isogeny* $\varphi : A^\vee \rightarrow A$ is an isogeny such that the map

$$\varphi^\vee : A^\vee \rightarrow (A^\vee)^\vee = A$$

is equal to φ .

If J and A are as in Section 3 then the principal polarization θ_J of J is symmetric, so the natural map $A^\vee \rightarrow A$ is a symmetric isogeny.

Lemma 5.2. *Suppose that A is a purely toric abelian variety over K and that $\varphi : A^\vee \rightarrow A$ is a symmetric isogeny. Let $\varphi_a : X_A \rightarrow X_{A^\vee}$ denote the induced map on character groups. Then*

$$\deg(\varphi) = \# \text{coker}(\varphi_a)^2.$$

Proof. By Corollary 8.7 applied to our isogeny φ (so what we are presently calling A^\vee and A are respectively called A and B in the discussion surrounding Theorem 8.6), we deduce that

$$\deg(\varphi) = \# \ker(\varphi) = \# \ker(\varphi_t) \cdot \# \ker(\varphi_t^\vee)$$

where φ_t and φ_t^\vee are the maps induced by φ and φ^\vee on closed fiber tori.

Since the character group X_A is, by definition, $\text{Hom}_{\bar{k}}(\mathcal{T}_{\bar{k}}, \mathbf{G}_{m_{\bar{k}}})$, where \mathcal{T} is the toric part of the closed fiber of A , it follows that $\# \ker(\varphi_t) = \# \text{coker}(\varphi_a)$. Since $\varphi = \varphi^\vee$, this proves the lemma. \square

6. Statement and Proof of the Main Theorem

Let K be as in Section 2, and let $\pi : J \rightarrow A$ be an optimal quotient. Assume that J is equipped with a symmetric principal polarization λ , that J has semistable reduction, and that A has purely toric reduction. Let X_A , X_{A^\vee} , and X_J denote the character groups of the toric parts of the closed fibers of the abelian varieties A , A^\vee , and J , respectively.

Let $\pi : J \rightarrow A$ be an optimal quotient, and let $\theta : A^\vee \rightarrow A$ denote the induced polarization. Let π_* , π^* , θ_* , and θ^* be the maps induced on character groups by the various functorialities, as indicated in the following two key diagrams:

$$\begin{array}{ccc} A^\vee \hookrightarrow J & \xrightarrow{\pi^\vee} & J \\ & \searrow \theta & \downarrow \pi \\ & & A \end{array} \qquad \begin{array}{ccc} X_A \hookrightarrow X_J & \xrightarrow{\pi^*} & X_J \\ & \searrow \theta^* & \downarrow \pi_* \\ & \searrow \theta_* & X_{A^\vee} \end{array}$$

The surjectivity of π_* is proved in Theorem 8.2. The injectivity of π^* follows because

$$\theta_* \pi_* \pi^* = \theta_* \theta^* = \deg(\theta) \neq 0,$$

and multiplication by a nonzero integer on a free abelian group is injective.

Let \mathcal{L} be the *saturation* of $\pi^* X_A$ in X_J ; thus $\pi^* X_A$ is a finite-index subgroup of \mathcal{L} and the quotient X_J/\mathcal{L} is torsion free. Let

$$\alpha : X_J \rightarrow \text{Hom}(\pi^* X_A, \mathbf{Z})$$

be the map defined by the monodromy pairing restricted to $X_J \times \pi^* X_A$. For L of finite index in \mathcal{L} , define the *degree* of L to be

$$m_L = [\alpha(X_J) : \alpha(L)],$$

and the *component group* of L to be

$$\Phi_L = \text{coker}(X_J \rightarrow \text{Hom}(L, \mathbf{Z})).$$

When $L = \mathcal{L}$ and A is fixed, for simplicity we write $m_X = m_{\mathcal{L}}$ and $\Phi_X = \Phi_{\mathcal{L}}$.

Recall that Φ_A is the component group of A and m_A is the square root of the degree of the induced map $A^\vee \rightarrow A$.

Theorem 6.1. *For any subgroup L of finite index in \mathcal{L} , the following relation holds:*

$$\frac{\#\Phi_A}{m_A} = \frac{\#\Phi_L}{m_L}.$$

6.1. Proof of the Main Theorem. The notation in this section is as in previous section.

Lemma 6.2. *Let $\pi_* : X_J \rightarrow X_{A^\vee}$ and $\alpha : X_J \rightarrow \text{Hom}(\pi^* X_A, \mathbf{Z})$ be as in previous section. Then*

$$\ker(\pi_*) = \ker(\alpha).$$

Proof. Suppose $x \in \ker(\pi_*)$, and let $y = \pi^* z$ with $z \in X_A$. Then

$$\langle x, y \rangle = \langle x, \pi^* z \rangle = \langle \pi_* x, z \rangle = 0,$$

so $x \in \ker(\alpha)$. Next let $x \in \ker(\alpha)$. Then for all $z \in X_A$,

$$0 = \langle x, \pi^* z \rangle = \langle \pi_* x, z \rangle,$$

so π_*x is in the kernel of the monodromy map

$$X_{A^\vee} \rightarrow \mathrm{Hom}(X_A, \mathbf{Z}).$$

Since X_{A^\vee} and $\mathrm{Hom}(X_A, \mathbf{Z})$ are free of the same finite rank and the cokernel is torsion, the monodromy map is injective. Thus $\pi_*x = 0$ and $x \in \ker(\pi_*)$. \square

Let $\pi^* : X_A \rightarrow X_J$ be as in previous section.

Lemma 6.3. *The monodromy-pairing map $X_J \rightarrow \mathrm{Hom}(X_J, \mathbf{Z})$ composed with restriction $\mathrm{Hom}(X_J, \mathbf{Z}) \rightarrow \mathrm{Hom}(\pi^*X_A, \mathbf{Z})$ gives rise to an exact sequence*

$$X_J \rightarrow \mathrm{Hom}(\pi^*X_A, \mathbf{Z}) \rightarrow \Phi_A \rightarrow 0.$$

Proof. Lemma 6.2 gives the following commutative diagram with exact rows

$$\begin{array}{ccccccc} 0 & \longrightarrow & X_J / \ker(\alpha) & \longrightarrow & \mathrm{Hom}(\pi^*X_A, \mathbf{Z}) & \longrightarrow & \mathrm{coker}(\alpha) \longrightarrow 0 \\ & & \downarrow \cong & & \downarrow \cong & & \downarrow \\ 0 & \longrightarrow & X_{A^\vee} & \longrightarrow & \mathrm{Hom}(X_A, \mathbf{Z}) & \longrightarrow & \Phi_A \longrightarrow 0. \end{array}$$

By Lemma 6.2, the first vertical map is an isomorphism. The second is an isomorphism because it is induced by the isomorphism $\pi^* : X_A \rightarrow \pi^*X_A$. It follows that $\mathrm{coker}(\alpha) \cong \Phi_A$, as claimed. \square

Recall that \mathcal{L} denotes the saturation of π^*X_A in X_J , and that $L \subset \mathcal{L}$ denotes a subgroup of finite index.

Lemma 6.4. *The rational number $\frac{\#\Phi_L}{m_L}$ is independent of the choice of L .*

Proof. Suppose L' is another finite index subgroup of \mathcal{L} , and let $n = [L : L']$. Here n is a rational number, the lattice index of L' in L . Since α is injective when restricted to \mathcal{L} , it follows that

$$m_{L'} = [\alpha(X_J) : \alpha(L')] = [\alpha(X_J) : \alpha(L)] \cdot [\alpha(L) : \alpha(L')] = m_L \cdot n.$$

Similarly, $\#\Phi_{L'} = \#\Phi_L \cdot n$. \square

Recall that $m_A = \sqrt{\deg(\theta)}$ and

$$\Phi_A \cong \mathrm{coker}(X_{A^\vee} \rightarrow \mathrm{Hom}(X_A, \mathbf{Z})),$$

where m_A is the degree of A and Φ_A is the component group of A .

Proof of Theorem 6.1. By Lemma 6.4 we may assume that $L = \pi^*X_A$. With this choice of L , Lemma 6.3 asserts that $\Phi_L \cong \Phi_A$. By Lemma 6.2, properties

of the index, and Lemma 5.2 we have

$$\begin{aligned}
 m_L &= [\alpha(X_J) : \alpha(L)] \\
 &= [\pi_*(X_J) : \pi_*(L)] \\
 &= [X_{A^\vee} : \pi_*(\pi^*X_A)] \\
 &= [X_{A^\vee} : \theta^*X_A] \\
 &= \# \operatorname{coker}(\theta^*) \\
 &= \sqrt{\deg(\theta)} = m_A.
 \end{aligned}$$

□

Recall that $\Phi_{\mathcal{L}}$ denotes the cokernel of the natural map $X_J \rightarrow \operatorname{Hom}(\mathcal{L}, \mathbf{Z})$ induced by composing the monodromy map $X_J \rightarrow \operatorname{Hom}(X_J, \mathbf{Z})$ with the natural restriction map $\operatorname{Hom}(X_J, \mathbf{Z}) \rightarrow \operatorname{Hom}(\mathcal{L}, \mathbf{Z})$.

Proposition 6.5. *The group $\Phi_{\mathcal{L}}$ is canonically isomorphic to the image of the map from Φ_J to Φ_A induced by $\pi : J \rightarrow A$. Thus*

$$\operatorname{image}(\Phi_J \rightarrow \Phi_A) \cong \Phi_{\mathcal{L}}.$$

Proof. Since $\pi^*X_A \subset \mathcal{L} \subset X_J$, an application of Lemma 6.3 gives the following commutative diagram with exact rows:

$$\begin{array}{ccccccc}
 X_J & \longrightarrow & \operatorname{Hom}(X_J, \mathbf{Z}) & \longrightarrow & \Phi_J & \longrightarrow & 0 \\
 \parallel & & \downarrow & & \downarrow & & \\
 X_J & \longrightarrow & \operatorname{Hom}(\mathcal{L}, \mathbf{Z}) & \longrightarrow & \Phi_{\mathcal{L}} & \longrightarrow & 0 \\
 \parallel & & \downarrow & & \downarrow & & \\
 X_J & \longrightarrow & \operatorname{Hom}(\pi^*X_A, \mathbf{Z}) & \longrightarrow & \Phi_A & \longrightarrow & 0.
 \end{array}$$

The map $\operatorname{Hom}(\mathcal{L}, \mathbf{Z}) \rightarrow \operatorname{Hom}(\pi^*X_A, \mathbf{Z})$ is an isomorphism, so the map $\Phi_{\mathcal{L}} \rightarrow \Phi_A$ is injective. Thus

$$\operatorname{image}(\Phi_J \rightarrow \Phi_A) \cong \operatorname{image}(\Phi_J \rightarrow \Phi_{\mathcal{L}}).$$

The cokernel of $\operatorname{Hom}(X_J, \mathbf{Z}) \rightarrow \operatorname{Hom}(\mathcal{L}, \mathbf{Z})$ surjects onto the cokernel of $\Phi_J \rightarrow \Phi_{\mathcal{L}}$. Using the exact sequence

$$0 \rightarrow \mathcal{L} \rightarrow X_J \rightarrow X_J/\mathcal{L} \rightarrow 0,$$

we find that

$$\operatorname{coker}(\operatorname{Hom}(X_J, \mathbf{Z}) \rightarrow \operatorname{Hom}(\mathcal{L}, \mathbf{Z})) \subset \operatorname{Ext}^1(X_J/\mathcal{L}, \mathbf{Z}).$$

Because \mathcal{L} is saturated, the quotient X_J/\mathcal{L} is torsion free, so the indicated Ext^1 group vanishes. Thus the map $\Phi_J \rightarrow \Phi_{\mathcal{L}}$ is surjective, from which the proposition follows. □

Corollary 6.6. *The cokernel of the map from Φ_J to Φ_A induced by $\pi : J \rightarrow A$ has order $m_A/m_{\mathcal{L}}$. Thus*

$$\# \operatorname{coker}(\Phi_J \rightarrow \Phi_A) = \frac{m_A}{m_{\mathcal{L}}}.$$

Proof. Combine Theorem 6.1 and Proposition 6.5. \square

7. Optimal Quotients of $J_0(N)$

In this section we specialize the general results of the rest of this paper to the concrete case in which $J = J_0(N)$ is the Jacobian of a modular curve, and $A = A_f$ is an optimal quotient of J attached to a modular forms. The paper [12] contains more computations like these.

7.1. Modular Curves and Semistable Reduction. Let $X_0(N)$ be the modular curve associated to the subgroup $\Gamma_0(N)$ of $\operatorname{SL}_2(\mathbf{Z})$ that consists of those matrices which are upper triangular modulo N . The algebraic curve $X_0(N)_{\mathbf{C}}$ can be constructed as a Riemann surface as the quotient

$$\Gamma_0(N) \backslash (\{z : z \in \mathbf{C}, \operatorname{Im}(z) > 0\} \cup \mathbf{P}^1(\mathbf{Q})),$$

and $X_0(N)$ has a canonical structure of algebraic curve over \mathbf{Q} .

It is well known that the p -new part of the Jacobian $J_0(N)$ of $X_0(N)$ has purely toric reduction at p when $p \nmid N$. Let us briefly recall the reason, writing $N = Mp$. Using the description of closed fibers of modular curves [10, Ch. 13] and Raynaud's result relating Néron models and Picard functors (as summarized in [2, Ch. 9]), the standard finite *flat* degeneracy maps $X_0(Mp) \rightarrow X_0(M)$ over $\mathbf{Z}_{(p)}$ induce a “*pushforward*” map on Néron model connected components

$$\operatorname{Pic}_{X_0(Mp)/\mathbf{Z}_{(p)}}^0 \longrightarrow \operatorname{Pic}_{X_0(M)/\mathbf{Z}_{(p)}}^0 \times \operatorname{Pic}_{X_0(M)/\mathbf{Z}_{(p)}}^0$$

which on the closed fiber is the map induced by *pullback* to the two components $X_0(M)_{/\mathbf{F}_p}$ in $X_0(Mp)_{/\mathbf{F}_p}$. The kernel of this latter map is a torus [2, Ex. 9.2.8], yet this kernel is visibly isogenous to the semistable mod p fiber of the dual of $J_0(Mp)^{\text{new}}$, whence the purely toric conclusion.

7.2. Newforms and Optimal Quotients. The Hecke algebra

$$\mathbf{T} = \mathbf{Z}[\dots T_n \dots] \subset \operatorname{End}(J_0(N))$$

is a commutative ring of endomorphisms of $J_0(N)$ of \mathbf{Z} -rank equal to the dimension of $J_0(N)$. The character group $X_{J,p}$ of $J_0(N)$ at p is equipped with a functorial action of \mathbf{T} . The Hecke algebra \mathbf{T} also acts on the complex vector space $S = S_2(\Gamma_0(N), \mathbf{C})$ of cusp forms.

Let f be a newform, and associate to f the ideal I_f of the Hecke algebra \mathbf{T} of elements which annihilate f . Then $\mathcal{O}_f = \mathbf{T}/I_f$ is an order in the ring of integers of the totally real number field K_f obtained by adjoining the Fourier coefficients of f to \mathbf{Q} . The quotient

$$A_f = J_0(N)/I_f J_0(N)$$

is an optimal quotient of $J_0(N)$ of dimension equal to $[K_f : \mathbf{Q}]$. As discussed in the previous section, A_f is purely toric at p .

7.3. Tamagawa Numbers. Let $\text{Frob}_p : X_J \rightarrow X_J$ denote the map induced by the Frobenius automorphism. We have $\text{Frob}_p = -W_p$, where W_p is the map induced by the Atkin-Lehner involution on $J_0(p)$. Let f be a newform, $A = A_f$ the corresponding optimal quotient, and w_p the sign of the eigenvalue of W_p on f .

Proposition 7.1.

$$\Phi_A(\mathbf{F}_p) = \begin{cases} \Phi_A(\overline{\mathbf{F}}_p) & \text{if } w_p = -1, \\ \Phi_A(\overline{\mathbf{F}}_p)[2] & \text{if } w_p = 1. \end{cases}$$

Proof. If $w_p = -1$, then $\text{Frob}_p = 1$ and the $\text{Gal}(\overline{\mathbf{F}}_p/\mathbf{F}_p)$ -action of $\Phi_A(\overline{\mathbf{F}}_p)$ is trivial. In this case $\Phi(\mathbf{F}_p) = \Phi(\overline{\mathbf{F}}_p)$. Next suppose $w_p = 1$. Recall that we have an exact sequence

$$0 \rightarrow X_{A^\vee} \rightarrow \text{Hom}(X_A, \mathbf{Z}) \rightarrow \Phi_A \rightarrow 0.$$

Since W_p acts as $+1$ on f , it also acts as $+1$ on each of the modules A , X_A , $\text{Hom}(X_A, \mathbf{Z})$, and Φ_A . Thus $\text{Frob}_p = -W_p$ acts as -1 on Φ_A . Since the subgroup of 2-torsion elements of a finite abelian group equals the subgroup of elements fixed under -1 , it follows that $\Phi_A(\mathbf{F}_p) = \Phi_A(\overline{\mathbf{F}}_p)[2]$. \square

Warning. When extending this result to the whole of $J_0(N)$, be careful. The action of $\text{Frob}_p = T_p$ need not be by ± 1 , even though it must be by an involution of order 2. For example, the component group of $J_0(65)$ at 5 is cyclic of order 42. The action of Frob_5 is by multiplication by -13 . Note that $(-13)^2 = 1 \pmod{42}$. The fixed points of multiplication by -13 is the order 14 subgroup of $\mathbf{Z}/42\mathbf{Z}$.

7.4. Computing Component Groups. Using modular symbols, we can enumerate the optimal quotients A_f of $J_0(N)$ (see, e.g., [1]) and compute the degree m_A (see [12, §3.1]). Suppose p is a prime that exactly divides N . As explained in [12], the method of graphs (see [14]) or the ideal theory of quaternion algebras (see [11]) can be used to compute $X = X_{J_0(N),p}$ with its \mathbf{T} -action and the monodromy pairing. We can then compute the following three modules:

1. the saturated submodule $\mathcal{L} = \bigcap_{t \in I_f} \ker(t)$ of X ,
2. the character group degree $m_X = m_{\mathcal{L}}$, and
3. $\Phi_X = \Phi_{\mathcal{L}}$.

By Theorem 6.1 we obtain

$$\#\Phi_{A,p} = \#\Phi_X \cdot \frac{m_A}{m_X}.$$

7.5. The Eisenstein Nature of Component Groups. The theorem below, which generalizes some of the results of [13] and [15], was conjectured by the second author after computing many component groups of quotients of $J_0(p)$ using the results of this paper. M. Emerton read an early version of this paper and subsequently announced a proof of the theorem below (see [6]).

Theorem 7.2 (Emerton). *Let p be a prime and let f_1, \dots, f_n be a set of representatives for the Galois-conjugacy classes of newforms in $S_2(\Gamma_0(p))$. Let A_1, \dots, A_n be the optimal quotients associated to f_1, \dots, f_n , respectively. Then for each i , $i = 1, \dots, n$, we have*

$$\#A_i(\mathbf{Q})_{\text{tor}} = \#\Phi_{A_i}(\overline{\mathbf{F}}_p) = \#\Phi_{A_i}(\mathbf{F}_p).$$

Furthermore,

$$\#\Phi_{J_0(p)}(\overline{\mathbf{F}}_p) = \prod_{i=1}^n \#\Phi_{A_i}(\overline{\mathbf{F}}_p).$$

Before Emerton proved the above assertion, the second author verified it using the algorithm of this paper for all $p \leq 757$, and, up to a power of 2, for all $p < 2000$.

Remark 7.3. It is tempting to guess that, e.g., the natural map

$$\Phi_{J_0(113)}(\overline{\mathbf{F}}_{113}) \rightarrow \prod_{i=1}^4 \Phi_{A_i}(\overline{\mathbf{F}}_{113})$$

is an isomorphism, but this is incorrect. Two of the $\Phi_{A_i}(\overline{\mathbf{F}}_{113})$ have order 2, so the product $\prod_{i=1}^4 \Phi_{A_i}(\overline{\mathbf{F}}_{113})$ is not a cyclic group. However, Mazur proved that the groups $\Phi_{J_0(p)}(\overline{\mathbf{F}}_p)$ are cyclic for all primes p .

7.6. Examples. In this section we give some examples of the numbers involved in computing component groups of quotients of $J_0(N)$. For more examples, see [12]. We use the notation for abelian varieties that is described in [1]. For example, **65A** is the “first” abelian variety quotient of $J_0(65)$ attached to a newform.

7.6.1. Quotients of $J_0(N)$. Table 1 contains many of the quantities involved in the computation of component groups for each of the newform optimal quotients for $N \in \{65, 66, 68, 69\}$.

7.6.2. Quotients of $J_0(p)^-$. We computed the quantities m_A , m_X , and Φ_X for each abelian variety A_f associated to a newform of prime level p with $p \leq 631$. Table 2 lists those A_f for which $w_p = -1$, along with the order of the corresponding component group. The first column, which is labeled “ A ” contains a description of A_f , the second column, labeled “ d ”, contains the dimension of A_f , and the third column, labeled “ $\#\Phi_A$ ”, contains the order $\#\Phi_{A_f,p}(\overline{\mathbf{F}}_p)$ of the component group.

TABLE 1. Component groups of quotients of $J_0(N)$

A	\dim	p	w_p	$\#\Phi_X$	m_X	m_A	$\#\Phi_A$
65A	1	5	+	1	2	2	1
		13	+	1	2		1
65B	2	5	+	3	2^2	2^2	3
		13	−	3	2^2		3
65C	2	5	−	7	2^2	2^2	7
		13	+	1	2^2		1
66A	1	2	+	1	2	2^2	2
		3	−	3	2^2		3
		11	+	1	2^2		1
66B	1	2	−	2	2	2^2	2^2
		3	+	1	2^2		1
		11	+	1	2^2		1
66C	1	2	−	1	2	$2^2 \cdot 5$	$2 \cdot 5$
		3	−	1	2^2		5
		11	−	1	$2^2 \cdot 5$		1
68A	2	17	+	2	$2 \cdot 3$	$2 \cdot 3$	2
69A	1	3	−	2	2	2	2
		23	+	1	2		1
69B	2	3	+	2	2	$2 \cdot 11$	$2 \cdot 11$
		23	−	2	$2 \cdot 11$		2

Remark 7.4. Theorem 7.2 together with [13, Prop. II.17.10] imply that the component groups of the A_f for which $w_p = +1$ are trivial, so we do not list them. An optimal quotient A_f of $J_0(p)$ with nonzero component group has nonzero rational torsion (by Theorem 7.2), so it factors through the Eisenstein quotient of $J_0(p)$. Also w_p acts as -1 on the Eisenstein quotient of $J_0(p)$, which is [13, Prop. II.17.10], and which is a deep result because of subtleties at the prime 2 (see the discussion in [13, III.1]).

TABLE 2. Component groups of quotients of $J_0(p)^-$

A	d	$\#\Phi_A$	A	d	$\#\Phi_A$	A	d	$\#\Phi_A$	A	d	$\#\Phi_A$
11A	1	5	157B	7	13	313A	2	1	487B	2	3
17A	1	2^2	163C	7	3^3	313C	12	$2 \cdot 13$	487C	3	1
19A	1	3	167B	12	83	317B	15	79	487D	16	3^3
23A	2	11	173B	10	43	331D	16	$5 \cdot 11$	491C	29	$5 \cdot 7^2$
29A	2	7	179A	1	1	337B	15	$2^2 \cdot 7$	499C	23	83
31A	2	5	179C	11	89	347D	19	173	503B	1	1
37B	1	3	181B	9	$3 \cdot 5$	349B	17	29	503C	1	1
41A	3	$2 \cdot 5$	191B	14	$5 \cdot 19$	353A	1	2	503D	3	1
43B	2	7	193C	8	2^4	353B	3	2	503F	26	251
47A	4	23	197C	10	7^2	353D	14	$2 \cdot 11$	509B	28	127
53B	3	13	199A	2	1	359D	24	179	521B	29	$2 \cdot 5 \cdot 13$
59A	5	29	199C	10	$3 \cdot 11$	367B	19	61	523C	26	$3 \cdot 29$
61B	3	5	211A	2	5	373C	17	31	541B	24	$3^2 \cdot 5$
67A	1	1	211D	9	7	379B	18	$3^2 \cdot 7$	547C	25	$7 \cdot 13$
67C	2	11	223C	12	37	383C	24	191	557B	1	1
71A	3	5	227B	2	1	389A	1	1	557D	26	139
71B	3	7	227C	2	1	389E	20	97	563A	1	1
73A	1	2	227E	10	113	397B	2	1	563E	31	281
73C	2	3	229C	11	19	397C	5	11	569B	31	$2 \cdot 71$
79B	5	13	233A	1	2	397D	10	3	571A	1	1
83B	6	41	233C	11	29	401B	21	$2^2 \cdot 5^2$	571B	1	1
89B	1	2	239B	17	$7 \cdot 17$	409B	20	$2 \cdot 17$	571C	2	1
89C	5	11	241B	12	$2^2 \cdot 5$	419B	26	$11 \cdot 19$	571D	2	1
97B	4	2^3	251B	17	5^3	421B	19	$5 \cdot 7$	571F	4	1
101B	7	5^2	257B	14	2^6	431B	1	1	571I	18	$5 \cdot 19$
103B	6	17	263B	17	131	431D	3	1	577A	2	3
107B	7	53	269C	16	67	431F	24	$5 \cdot 43$	577B	2	1
109A	1	1	271B	16	$3^2 \cdot 5$	433A	1	1	577C	3	1
109C	4	3^2	277B	3	1	433B	3	1	577D	18	2^4
113A	1	2	277D	9	23	433D	16	$2^2 \cdot 3^2$	587C	31	293
113B	2	2	281B	16	$2 \cdot 5 \cdot 7$	439C	25	73	593B	1	2
113D	3	7	283B	14	47	443C	1	1	593C	2	1
127B	7	$3 \cdot 7$	293B	16	73	443E	22	$13 \cdot 17$	593E	27	$2 \cdot 37$
131B	10	$5 \cdot 13$	307A	1	1	449B	23	$2^4 \cdot 7$	599C	37	$13 \cdot 23$
137B	7	$2 \cdot 17$	307B	1	1	457C	20	$2 \cdot 19$	601B	29	$2 \cdot 5^2$
139A	1	1	307C	1	1	461D	26	$5 \cdot 23$	607D	31	101
139C	7	23	307D	1	1	463B	22	$7 \cdot 11$	613C	27	$3 \cdot 17$
149B	9	37	307E	2	3	467C	26	233	617B	28	$2 \cdot 7 \cdot 11$
151B	3	1	307F	9	17	479B	32	239	619B	30	103
151C	6	5^2	311B	22	$5 \cdot 31$	487A	2	1	631B	32	$3 \cdot 5 \cdot 7$

8. Appendix: Some Facts Concerning Toric Reduction

Let R be a discrete valuation ring with fraction field K and residue field k . For any abelian variety A over K , with Néron model \mathcal{A} over R , we denote by

X_A the character group of the toric part of \mathcal{A}_k^0 (the connected component of the closed fiber of \mathcal{A}). All group schemes below are understood to be commutative.

Our aim in this appendix is to prove a couple of facts (Theorem 8.2 and Theorem 8.6) which are no doubt well-known to experts but for which published proofs do not appear to be readily available. We begin with a simple and basic lemma.

Lemma 8.1. *Let $f : G \rightarrow G'$ be a map between multiplicative (resp. étale) finite flat group schemes over R . The map f is a closed immersion (resp. faithfully flat) if and only if the generic fiber map f_K is a closed immersion (resp. faithfully flat).*

Proof. Cartier duality interchanges étaleness and multiplicativeness, as well as closed immersions and faithfully flat maps (as the latter two properties may be checked on the closed fiber, for which one is reduced to the standard case of finite commutative group schemes over a field). Thus, it suffices to consider the étale case. By faithfully flat base change to a strict henselization of R , we are reduced to the case where our finite étale group schemes are constant. Since faithful flatness is equivalent to surjectivity (for maps between étale schemes over a base), the lemma is now physically clear. \square

Now we turn to the first of the two main results we want to prove. Let $\pi : J \rightarrow A$ be an optimal quotient of abelian varieties over K (i.e., we assume that $\ker \pi$ is an abelian variety over K), and assume that J has semistable reduction over R (so A does too). We do not yet make any hypotheses of purely toric reduction. The dual abelian varieties A^\vee and J^\vee again have semistable reduction, as they are isogenous to A and J respectively.

Theorem 8.2. *With notation as above, the map $X_{J^\vee} \rightarrow X_{A^\vee}$ induced by π is surjective.*

Proof. The underlying idea comes down to two facts: Lemma 8.1 and the fact that we can lift tori on the level of ℓ -divisible groups for any prime ℓ . More precisely, we argue as follows. By Proposition 3.3, the map $\pi^\vee : A^\vee \rightarrow J^\vee$ is a closed immersion of abelian varieties. We will use this to prove that the induced map π_t^\vee on closed fiber tori of Néron models is a closed immersion. Since the “character group” functor sets up an anti-equivalence of categories between tori over a field F and finite free \mathbf{Z} -modules with continuous action of $\text{Gal}(F_s/F)$, identifying closed immersions of tori with surjections of character groups and surjections of tori with “saturated injections” of character groups (i.e., injections with torsion-free cokernel), the closed immersion property for π_t^\vee on the closed fiber tori will yield the desired surjection of character groups.

In general the “Néron model” functor doesn’t behave well for closed immersions. That is, just because π^\vee is a closed immersion, it does not follow purely formally that π^\vee induces a closed immersion on Néron models. Nevertheless, we claim quite generally that if $B \rightarrow B'$ is a closed immersion of abelian varieties

over K with semistable Néron models, then the induced map $T \rightarrow T'$ on closed fiber tori is a closed immersion. For this it is sufficient to prove that the induced map on ℓ -divisible groups $T[\ell^\infty] \rightarrow T'[\ell^\infty]$ is a closed immersion for *all* primes ℓ (i.e., all maps $T[\ell^n] \rightarrow T'[\ell^n]$ are closed immersions). Indeed, suppose we verify this closed immersion property on torsion, and let H be the kernel of $T \rightarrow T'$, so $H[\ell^n] = 0$ for all primes ℓ and positive integers n . The torus $(H_{/k}^0)_{\text{red}}$ must vanish (as it has no non-trivial torsion) and hence H is finite. If N is the order of H , then $H = H[N] = 0$. The map $T \rightarrow T'$ is then a monomorphism between algebraic groups over a field and hence is a closed immersion, as desired.

In order to verify that the ℓ -divisible group maps $T[\ell^\infty] \rightarrow T'[\ell^\infty]$ are closed immersions for all ℓ , we can make the faithfully flat base change to the henselization of R (which commutes with formation of Néron models) to reduce to the case where R is henselian. Now we recall the following basic result of Grothendieck:

Lemma 8.3. *Let R be a henselian local ring, G a quasi-finite separated R -scheme of finite presentation. There is a unique decomposition*

$$G = G_f \coprod G'$$

into disjoint clopen pieces with G_f finite over R (called the “finite part” of G) and G' having empty closed fiber. The formation of G_f is functorial in G and is compatible with henselian local base change and formation of fiber products over R .

If moreover G is a group scheme over R , then G_f is a clopen subgroup scheme and there exists a unique multiplicative closed R -subgroup scheme G_μ inside of G whose closed fiber is the multiplicative part of the closed fiber of G (G_μ is called the “multiplicative part” of G). The formation of G_μ is functorial in G .

Proof. For the first part, see [8, IV₄, 18.5.11(c)] (aside from the functorial properties, which are obvious). The second part, concerning group schemes, is a mechanical consequence of the first part (including the functoriality of the finite part). For example, the existence of G_μ follows from considering the connected-étale sequence of the Cartier dual of G_f over the henselian local base R , and the uniqueness and functoriality follows from the functoriality of $G \rightsquigarrow G_f$ and the functoriality of the connected-étale sequence. □

Remark 8.4. Assuming R in Lemma 8.3 is a discrete valuation ring (with fraction field K and residue field k), let us make some observations concerning the behavior of Lemma 8.3 with respect to primary components, as this will be useful later. Let's suppose that N and M are relatively prime integers with NM divisible by the order of G_K , and hence killing G . Thus, by functoriality we have $G = G[N] \times_R G[M]$ where $G[N]$ and $G[M]$ are quasi-finite separated R -group schemes. We claim that $G[N]$ and $G[M]$ are also *flat* over R , whence it follows that the formation of G_f and G_μ is compatible with passage to “primary components”.

In other words, if ℓ is a prime and ℓ^n is divisible by the ℓ -part of the order of G_K , then we claim that $G[\ell^n]$ is R -flat. From the clopen decomposition $G = G_f \coprod G'$, it is easy to see that $G[\ell^n] = G_f[\ell^n] \coprod X_n$ for some finite K -scheme X_n , so for the issue of R -flatness we can replace G with G_f . We are thereby reduced to the finite flat case, so we can use the proof of [10, 1.7.2].

The significance of Lemma 8.3 for our purposes is the following standard consequence.

Corollary 8.5. *Let A be an abelian variety over the fraction field K of a henselian discrete valuation ring R with residue field k . Let \mathcal{A} be the Néron model of A , and assume that \mathcal{A} has semistable reduction. For every prime ℓ , there exists a unique multiplicative ℓ -divisible group Γ_ℓ inside of \mathcal{A} whose closed fiber is the ℓ -divisible group of the torus T of \mathcal{A}_k^0 . The formation of Γ_ℓ is functorial in A .*

Proof. Fix ℓ . By the semistability hypothesis, the multiplication maps $\ell^n : \mathcal{A} \rightarrow \mathcal{A}$ are quasi-finite flat, so $\mathcal{A}[\ell^n]$ is a quasi-finite flat separated R -group scheme. Let $\mathcal{A}[\ell^n]_\mu$ denote its multiplicative part (as in Lemma 8.3), so the multiplicative $T[\ell^n] \hookrightarrow \mathcal{A}[\ell^n]_\mu$ lies inside of $(\mathcal{A}[\ell^n]_\mu)_k$. The “closed fiber” functor is an equivalence of categories between finite flat multiplicative group schemes over R and k (since Cartier duality reduces this to the étale case, and the “closed fiber” functor is an equivalence of categories between finite étale R -schemes and finite étale k -schemes [8, IV₄, 18.5.12]). Thus, there exists a unique multiplicative closed R -subgroup scheme $\Gamma^{(n)} \hookrightarrow \mathcal{A}[\ell^n]_\mu$ whose closed fiber is $T[\ell^n]$.

Moreover, using the equivalence of categories just mentioned, $\mathcal{A}[\ell^n]_\mu$ lies inside of $\mathcal{A}[\ell^{n+1}]_\mu$ and $\Gamma^{(n)}$ lies inside of $\Gamma^{(n+1)}$. The resulting system $\Gamma_\ell = \{\Gamma^{(n)}\}$ over R forms an ℓ -divisible group on the closed fiber and hence is an ℓ -divisible group over R . This settles the desired existence, as well as the desired uniqueness. The functoriality of Γ_ℓ in A follows from the functoriality of toric parts on the closed fiber of Néron models. \square

Returning to the proof of Theorem 8.2, recall that we were studying the map of toric parts $j_t : T \rightarrow T'$ induced by a closed immersion $j : B \hookrightarrow B'$ of semistable abelian varieties over K , with R henselian. We wanted the map

$$j_t[\ell^\infty] : T[\ell^\infty] \rightarrow T'[\ell^\infty]$$

to be a closed immersion for all primes ℓ (as we have seen that this forces $T \rightarrow T'$ to be a closed immersion, which is what we really want to show). Fix ℓ . By Corollary 8.5 there exist unique multiplicative ℓ -divisible groups Γ and Γ' over R in the respective Néron models \mathcal{B} and \mathcal{B}' such that Γ and Γ' respectively lift the ℓ -divisible groups of the tori of the closed fibers. Hence, it suffices to show that the R -map $\gamma : \Gamma \rightarrow \Gamma'$ induced by the Néron functoriality map $N(j)$ is a closed immersion. The generic fiber map γ_K is a closed immersion since it “sits inside” the generic fiber ℓ -divisible groups of B and B' , the map between which is a closed immersion since $j : B \rightarrow B'$ is a closed immersion. Now we

use Lemma 8.1 (applied at all finite torsion levels) to conclude that γ is a closed immersion. This completes the proof of Theorem 8.2. \square

We now turn to a result which requires a stronger hypothesis on the closed fiber. Note that we retain the hypothesis that R is henselian (this hypothesis arose in the proof of Theorem 8.2, even though it wasn't needed for the statement). Let A and B be abelian varieties over K with purely toric reduction (i.e., their Néron models have closed fiber connected components which are tori). Let $\varphi : A \rightarrow B$ be an isogeny, and let $\varphi_t : T_A \rightarrow T_B$ be the induced map on the closed fiber toric parts (i.e., connected components) of the Néron models. We denote by $\varphi_t^\vee : T_{B^\vee} \rightarrow T_{A^\vee}$ the analogous map induced by the dual isogeny φ^\vee . Since the map φ_t is an isogeny (by functoriality), the kernel $\ker(\varphi_t)$ is a finite multiplicative k -group scheme.

For any finite multiplicative k -group scheme G , we let \widetilde{G} denote the (unique) multiplicative finite flat R -group scheme with closed fiber G . For example, $\widetilde{\ker(\varphi_t)}$ is a multiplicative R -group scheme which lies inside of

$$\ker(N(\varphi))_\mu$$

(where $N(\varphi)$ is the map induced by Néron functoriality). Thus, we have a natural closed immersion

$$\widetilde{(\ker \varphi_t)_K} \hookrightarrow \ker \varphi$$

and likewise we have a natural quotient map

$$\ker(\varphi^\vee)^\vee \rightarrow \widetilde{\ker(\varphi_t^\vee)_K}^\vee$$

dual to the natural closed immersion using the isogeny φ^\vee .

By the duality theory for abelian varieties (particularly the adjointness of φ and φ^\vee with respect to the scheme-theoretic Weil pairing over K), there is a canonical perfect duality K -group scheme duality between $\ker(\varphi)$ and $\ker(\varphi^\vee)$ over K , whence there is a natural quotient map of K -group schemes

$$\ker(\varphi) \simeq \ker(\varphi^\vee)^\vee \rightarrow \widetilde{\ker(\varphi_t^\vee)_K}^\vee.$$

Theorem 8.6. *The diagram of K -group schemes*

$$0 \rightarrow \widetilde{\ker(\varphi_t)_K} \rightarrow \ker(\varphi) \rightarrow \widetilde{\ker(\varphi_t^\vee)_K}^\vee \rightarrow 0$$

is exact.

The content of the proof is the Grothendieck Orthogonality Theorem. Moreover, Theorem 8.6 is implicit in Grothendieck's construction of the monodromy pairing for semiabelian varieties.

Proof. The exact sequence of the theorem says that the finite flat K -group schemes

$$\ker(\varphi)/\widetilde{\ker(\varphi_t)_K} \quad \text{and} \quad \widetilde{\ker(\varphi_t^\vee)_K}^\vee$$

are canonically Cartier dual to each other compatibly with the perfect duality between $\ker(\varphi)$ and $\ker(\varphi^\vee)$. More precisely, let \mathcal{A} and \mathcal{B} denote the Néron models of A and B , respectively, let

$$G = \ker(\mathcal{A} \rightarrow \mathcal{B}), \quad G^\vee = \ker(\mathcal{B}^\vee \rightarrow \mathcal{A}^\vee),$$

so G and G^\vee are both quasi-finite flat separated R -group schemes whose generic fibers are the $\ker(\varphi)$ and $\ker(\varphi^\vee)$ in the theorem (the R -flatness of G and G^\vee arises from the semiabelian condition, since any quasi-finite morphism between semi-abelian schemes is necessarily flat, as can be checked on geometric fibers). Being quasi-finite flat and separated, the R -group schemes G and G^\vee have canonical respective “finite parts” G_f and G_f^\vee and “multiplicative parts” G_μ and G_μ^\vee (as in Lemma 8.3). Beware that we do *not* claim G_f^\vee (resp. G_μ^\vee) is the Cartier dual to G_f (resp. G_μ); usually such duality does not hold.

Since G_μ and G_μ^\vee are finite flat R -group schemes, the quotients G/G_μ and G^\vee/G_μ^\vee make sense as quasi-finite flat separated R -group schemes. The theorem almost says that there is a canonical duality between $(G/G_\mu)_K = G_K/(G_\mu)_K$ and $(G_\mu^\vee)_K$, induced by the duality between $G_K = \ker(\varphi)$ and $G_K^\vee = \ker(\varphi^\vee)$, except for the mild problem that G_μ might be larger than $\widetilde{\ker(\varphi_t)}$ (i.e., possibly $(G_\mu)_k$ is not entirely inside of \mathcal{A}_k^0) and likewise G_μ^\vee might be larger than $\widetilde{\ker(\varphi_t^\vee)}$.

We will work on ℓ -primary components for each prime ℓ individually. In order to permit this, we use Remark 8.4. We will first treat the more subtle case when ℓ is the residue characteristic, and then we’ll handle the case when it isn’t. The advantage of working with the case in which ℓ is the residue characteristic is that multiplicative $\widetilde{\text{finite } k\text{-group schemes}}$ are automatically *connected*. Thus, in this case $(G_\mu)_\ell = \widetilde{\ker(\varphi_t)_\ell}$ and $(G_\mu^\vee)_\ell = \widetilde{\ker(\varphi_t^\vee)_\ell}$.

Since

$$0 \rightarrow G_K \rightarrow A \rightarrow B \rightarrow 0$$

is an exact sequence of abelian sheaves on the fppf site over $\text{Spec}(K)$, by the usual snake lemma argument (and the fact that the ℓ -part $(G_K)_\ell$ of G_K is killed by a big power of ℓ) we obtain an exact sequence

$$0 \rightarrow (G_K)_\ell \rightarrow A[\ell^\infty] \rightarrow B[\ell^\infty] \rightarrow 0.$$

Arguing as in Corollary 8.5, there is an exact sequence over $\text{Spec}(R)$

$$0 \rightarrow (G_\mu)_\ell \rightarrow \mathcal{A}[\ell^\infty]_t \rightarrow \mathcal{B}[\ell^\infty]_t \rightarrow 0$$

which lifts the exact sequence involving ℓ -divisible groups of tori on the closed fiber (as $(G_\mu)_\ell$ *must* be in the relative connected component of \mathcal{A}). Passing to the generic fiber over K gives us a commutative diagram with exact rows and closed immersions along columns

$$\begin{array}{ccccccc} 0 & \longrightarrow & ((G_\mu)_K)_\ell & \longrightarrow & A[\ell^\infty]_t & \longrightarrow & B[\ell^\infty]_t \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & (G_K)_\ell & \longrightarrow & A[\ell^\infty] & \longrightarrow & B[\ell^\infty] \longrightarrow 0 \end{array}$$

where $A[\ell^\infty]_t$ denotes the K -fiber of the ℓ -divisible group $\{\mathcal{A}[\ell^n]_t\}$, and likewise for $B[\ell^\infty]_t$.

Using the snake lemma in the category of fppf abelian sheaves over $\mathrm{Spec}(K)$, we get a short exact sequence of cokernels

$$0 \rightarrow ((G/G_\mu)_K)_\ell \rightarrow A[\ell^\infty]/(\mathrm{toric}) \rightarrow B[\ell^\infty]/(\mathrm{toric}) \rightarrow 0$$

where all maps are the natural ones, and right two terms are ℓ -divisible groups over K .

The Grothendieck Orthogonality Theorem (see [9, Exp. IX, Prop 5.6]) asserts that the perfect scheme-theoretic Weil pairing between $A[\ell^n]$ and $A^\vee[\ell^n]$ makes $A[\ell^n]_t$ and $A^\vee[\ell^n]_f$ exact annihilators, where $A[\ell^n]_f$ denotes the K -fiber of the finite part of the $\mathcal{A}^0[\ell^n]$ and $A[\ell^n]_t$ denotes the K -fiber of the unique R -subgroup scheme in $\mathcal{A}^0[\ell^n]$ lifting the ℓ^n -torsion on the closed fiber torus. By the purely toric condition applied to A^\vee , we see $A^\vee[\ell^n]_f = A^\vee[\ell^n]_t$. Thus, the orthogonality theorem says that $A[\ell^n]/A[\ell^n]_t$ and $A^\vee[\ell^n]_t$ are in perfect duality via the scheme-theoretic Weil pairing over K .

Passing to the limit, we get a canonical isomorphism of ℓ -divisible groups

$$A[\ell^\infty]/(\mathrm{toric}) = (A^\vee[\ell^\infty]_t)^\vee.$$

But φ and φ^\vee are adjoint with respect to Weil pairing, so we conclude that the diagram

$$\begin{array}{ccc} A[\ell^\infty]/(\mathrm{toric}) & \xlongequal{\quad} & (A^\vee[\ell^\infty]_t)^\vee \\ \downarrow \varphi & & \downarrow (\varphi_t^\vee)^\vee \\ B[\ell^\infty]/(\mathrm{toric}) & \xlongequal{\quad} & (B^\vee[\ell^\infty]_t)^\vee \end{array}$$

commutes. Thus, we get an isomorphism between the kernels of these vertical isogenies. The kernel of the left column is $((G/G_\mu)_K)_\ell$, as we saw above. Meanwhile, the kernel of the right is (by duality theory of ℓ -divisible groups) exactly the dual of $\ker(\varphi_t^\vee) = (G_\mu^\vee)_K$. This gives the desired perfect duality between $(G/G_\mu)_K$ and $(G_\mu^\vee)_K$ on ℓ -primary parts for ℓ equal to the residue characteristic.

Now we consider the case when ℓ is not equal to the residue characteristic. There is no loss of generality in passing to the case of a strictly henselian base R . Thus, the closed fiber tori have *constant* ℓ -divisible groups. Also, we can work with \mathbf{Z}_ℓ -modules of geometric points (over K) via Tate's construction. The “toric” part of the ℓ -adic Tate module $T_\ell(A)$ is exactly the (saturated) maximal submodule with trivial Galois action, since a compatible system of ℓ -power torsion points in $A(K) = \mathcal{A}(R)$ must lie entirely inside of $\mathcal{A}^0(R)$ (thanks to the finiteness of the component group) and we can identify $\mathcal{A}^0(R)[\ell^n]$ with the (constant) ℓ^n -torsion on the split torus \mathcal{A}_k^0 over the separably closed k .

Using inverse limits, we see that $T_\ell(A) \rightarrow T_\ell(B)$ is injective with cokernel $(G_K)_\ell$ (= geometric points of ℓ -part), and this cokernel is exactly $\ker(\varphi)_\ell$. Likewise, the cokernel of the map

$$T_\ell(A)_t \rightarrow T_\ell(B)_t$$

on “toric” parts (i.e., ℓ -adic Tate module $\widetilde{\ker(\varphi_t)}$ generic fibers of the lifts of the ℓ -divisible groups of closed fiber tori) is $(\widetilde{\ker(\varphi_t)_K})_\ell$.

Thus, we get a commutative diagram with horizontal exact sequences

$$\begin{array}{ccccccc} 0 & \longrightarrow & T_\ell(A)_t & \longrightarrow & T_\ell(B)_t & \longrightarrow & (\widetilde{\ker(\varphi_t)_K})_\ell \longrightarrow 0 \\ & & \downarrow & & \downarrow & & \downarrow \\ 0 & \longrightarrow & T_\ell(A) & \longrightarrow & T_\ell(B) & \longrightarrow & \ker(\varphi)_\ell \longrightarrow 0 \end{array}$$

with columns given by the natural maps. These vertical maps are all injective, so by the snake lemma we get a short exact sequence of cokernels. We can now use the exact same Weil pairing arguments with the Grothendieck orthogonality theorem (now in the easier “ $\ell \neq p$ ” form of the orthogonality theorem [9, Exp IX, 2.4]), essentially just as we argued in the previous case. One distinction is that the use of inverse limit Tate modules rather than direct limits causes some maps to switch direction.

More specifically we have a commutative square with horizontal isomorphisms (thanks to the orthogonality theorem)

$$\begin{array}{ccc} T_\ell(A)/T_\ell(A)_t & \xlongequal{\quad} & T_\ell(A^\vee)_t^\vee \\ \varphi \downarrow & & \downarrow T_\ell(\varphi_t^\vee)^\vee \\ T_\ell(B)/T_\ell(B)_t & \xlongequal{\quad} & T_\ell(B^\vee)_t^\vee \end{array}$$

This induces an isomorphism

$$((\ker \varphi)/\widetilde{\ker(\varphi_t)_K})_\ell \simeq ((\ker \varphi_t^\vee)_K)_\ell^\vee$$

between the vertical cokernels, and by construction this isomorphism is compatible with Weil pairings, whence the desired perfect pairing has been shown. \square

Let R be an arbitrary dvr (not necessarily henselian), let $\varphi : A \rightarrow B$ be an isogeny, and let $\varphi_t : T_A \rightarrow T_B$ be the induced map on the closed fiber toric parts, as above.

Corollary 8.7. *The order of $\ker(\varphi)$ is the product of the orders of $\ker \varphi_t$ and $\ker \varphi_t^\vee$.*

Proof. Pass to the henselization of R and use Theorem 8.6. \square

Acknowledgements

This paper was inspired by lectures of R. Coleman and K. Ribet, and a letter from Ribet to Mestre (see [17]), which contains some of the results of the present paper in the case when A has dimension 1. The second author would like to thank A. Abbes, A. Agashe, D. Kohel, and D. Lorenzini, for helpful conversations.

References

- [1] A. Agashe and W. A. Stein, *Visibility of Shafarevich-Tate groups of abelian varieties: Evidence for the Birch and Swinnerton-Dyer conjecture*, (2001).
- [2] S. Bosch, W. Lütkebohmert, and M. Raynaud, *Néron models*. Ergebnisse der Mathematik und ihrer Grenzgebiete (3), 21. Springer-Verlag, Berlin, 1990.
- [3] C. Chevalley, *Une démonstration d'un théorème sur les groupes algébriques*, J. Math. Pures Appl. (9) **39** 1960, 307–317.
- [4] B. Conrad, *A modern proof of Chevalley's theorem on algebraic groups*, <http://www-math.mit.edu/~dejong/papers/chev.dvi>
- [5] B. Edixhoven, *L'action de l'algèbre de Hecke sur les groupes de composantes des jacobiniennes des courbes modulaires est "Eisenstein"*. Courbes modulaires et courbes de Shimura (Orsay, 1987/1988). Astérisque No. 196-197, (1991), 7–8, 159–170 (1992).
- [6] M. Emerton, *Optimal quotients of modular Jacobians*, (2001), preprint.
- [7] E. V. Flynn, F. Leprévost, E. F. Schaefer, W. A. Stein, M. Stoll, and J. L. Wetherell, *Empirical evidence for the Birch and Swinnerton-Dyer conjectures for modular Jacobians of genus 2 curves*, Math. Comp. **70** (2001), no. 236, 1675–1697 (electronic).
- [8] A. Grothendieck, *Éléments de géométrie algébrique*, Publications Mathématiques IHES, **4,8,11,17,20,24,28,32**, 1960–7.
- [9] A. Grothendieck, *Groupes de monodromie en géométrie algébrique*, Lecture Notes in Math **288**, Springer-Verlag, Heidelberg (1972).
- [10] N. Katz, B. Mazur, *Arithmetic moduli of elliptic curves*. Annals of Mathematics Studies, 108. Princeton University Press, Princeton, NJ, 1985.
- [11] D. R. Kohel, *Hecke module structure of quaternions*. Class field theory—its centenary and prospect (Tokyo, 1998), 177–195, Adv. Stud. Pure Math., 30, Math. Soc. Japan, Tokyo, 2001.
- [12] D. R. Kohel and W. A. Stein, *Component Groups of Quotients of $J_0(N)$* , Proceedings of the 4th International Symposium (ANTS-IV), Leiden, Netherlands, July 2–7, 2000 (Berlin), Springer, 2000.
- [13] B. Mazur, *Modular curves and the Eisenstein ideal*. Inst. Hautes Études Sci. Publ. Math. No. 47 (1977), 33–186 (1978).
- [14] J.-F. Mestre, *La méthode des graphes. Exemples et applications*. Proceedings of the international conference on class numbers and fundamental units of algebraic number fields (Katata, 1986), 217–242, Nagoya Univ., Nagoya, 1986.
- [15] J.-F. Mestre and J. Oesterlé, *Courbes de Weil semi-stables de discriminant une puissance m -ième*, J. Reine Angew. Math. **400** (1989), 173–184.
- [16] D. Mumford, *Abelian varieties*. Tata Institute of Fundamental Research Studies in Mathematics, No. 5, Published for the Tata Institute of Fundamental Research, Bombay; Oxford University Press, London 1970.
- [17] K. A. Ribet, *Letter about component groups of elliptic curves*, [arXiv:math.AG/0105124v1](https://arxiv.org/abs/math/0105124v1) (2001).
- [18] J. Tate, *Algorithm for determining the type of a singular fiber in an elliptic pencil*. Modular functions of one variable, IV (Proc. Internat. Summer School, Univ. Antwerp, Antwerp, 1972), 33–52. Lecture Notes in Math., Vol. 476, Springer, Berlin, 1975.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MICHIGAN, 2074 EAST HALL, ANN ARBOR, MI 48109-1109, USA.

E-mail address: bdconrad@math.lsa.umich.edu

HARVARD UNIVERSITY, DEPARTMENT OF MATHEMATICS, ONE OXFORD STREET, CAMBRIDGE, MA 02138, USA.

E-mail address: was@math.harvard.edu