

HILBERT’S TENTH PROBLEM AND MAZUR’S CONJECTURES IN COMPLEMENTARY SUBRINGS OF NUMBER FIELDS

KIRSTEN EISENTRÄGER, GRAHAM EVEREST AND ALEXANDRA SHLAPENTOKH

ABSTRACT. We show that Hilbert’s Tenth Problem is undecidable for complementary subrings of number fields and that the \mathfrak{p} -adic and archimedean ring versions of Mazur’s conjectures do not hold in these rings. More specifically, given a number field K , a positive integer $t > 1$, and t nonnegative computable real numbers $\delta_1, \dots, \delta_t$ whose sum is one, we prove that the nonarchimedean primes of K can be partitioned into t disjoint recursive subsets S_1, \dots, S_t of densities $\delta_1, \dots, \delta_t$, respectively such that Hilbert’s Tenth Problem is undecidable for each corresponding ring \mathcal{O}_{K, S_i} . We also show that we can find a partition as above such that each ring \mathcal{O}_{K, S_i} possesses an infinite Diophantine set which is discrete in every topology of the field. The only assumption on K we need is that there is an elliptic curve of rank one defined over K .

1. Introduction

Hilbert’s Tenth Problem in its original form was to find an algorithm to decide, given a polynomial equation $f(x_1, \dots, x_n) = 0$ with coefficients in the ring \mathbb{Z} of integers, whether it has a solution with $x_1, \dots, x_n \in \mathbb{Z}$. In 1969 Matiyasevich [8], using work by Davis *et al.* (see [5]), proved that no such algorithm exists, i.e. Hilbert’s Tenth Problem is undecidable. Since then, analogs of this problem have been studied by asking the same question for polynomial equations with coefficients and solutions in other recursive commutative rings R . We will refer to this analog of the original problem as *Hilbert’s Tenth Problem over R* . Perhaps the most important unsolved problem in this area is the case of $R = \mathbb{Q}$. One natural approach to showing that Hilbert’s Tenth Problem is undecidable for a ring R of characteristic 0 is to show that \mathbb{Z} admits a Diophantine definition over R , or more generally that there is a Diophantine model of the ring \mathbb{Z} over R . We define these notions below.

Definition 1.1. Let R be a commutative ring. Suppose $A \subseteq R^k$ for some $k \in \mathbb{N}$. We say that A has a *Diophantine definition over R* if there exists a polynomial

$$f(t_1, \dots, t_k, x_1, \dots, x_n) \in R[t_1, \dots, t_k, x_1, \dots, x_n]$$

such that for any $(t_1, \dots, t_k) \in R^k$,

$$(t_1, \dots, t_k) \in A \iff \exists x_1, \dots, x_n \in R, f(t_1, \dots, t_k, x_1, \dots, x_n) = 0.$$

In this case we also say that A is a *Diophantine subset* of R^k , or that A is *Diophantine over R* .

Received by the editors January 23, 2011.

1991 *Mathematics Subject Classification.* 11G05, 11U05.

Sadly, the second author passed away before the final version of the paper was completed.

Remark 1.1. Suppose that R is a domain whose quotient field is not algebraically closed. Then

- (a) Relaxing Definition 1.1 to allow an arbitrary finite conjunction of equations in place of the single equation on the right-hand side does not enlarge the collection of Diophantine sets.
- (b) Finite unions and finite intersections of Diophantine sets are Diophantine.

See [25] for details.

Definition 1.2. A *Diophantine model of \mathbb{Z} over a ring R* is a Diophantine subset $A \subseteq R^k$ for some k together with a bijection $\phi: \mathbb{Z} \rightarrow A$ such that the graphs of addition and multiplication (subsets of \mathbb{Z}^3) correspond under ϕ to Diophantine subsets of $A^3 \subseteq R^{3k}$.

In 1992, Mazur formulated a conjecture that would imply that a Diophantine definition of \mathbb{Z} over \mathbb{Q} does not exist, and which also ruled out the existence of a Diophantine model of \mathbb{Z} over \mathbb{Q} [4]. One form of Mazur's conjecture was that for a variety X over \mathbb{Q} , the closure of $X(\mathbb{Q})$ in the topological space $X(\mathbb{R})$ should have at most finitely many connected components [9, 10]. This conjecture also implied that no infinite set which is discrete in the archimedean topology has a Diophantine definition over \mathbb{Q} .

Mazur also formulated a version of his conjecture applying to both archimedean and nonarchimedean completions of arbitrary number fields [11, p. 257]:

Question 1.1. Let V be any variety defined over a number field K . Let \mathcal{S} be a finite set of places of K , and consider $K_{\mathcal{S}} = \prod_{v \in \mathcal{S}} K_v$ viewed as locally compact topological ring. Let $V(K_{\mathcal{S}})$ denote the topological space of $K_{\mathcal{S}}$ -rational points. For every point $p \in V(K_{\mathcal{S}})$ define $W(p) \subset V$ to be the subvariety defined over K that is the intersection of Zariski closures of the subsets $V(K) \cap U$, where U ranges through all open neighborhoods of p in $V(K_{\mathcal{S}})$. As p ranges through the points of $V(K_{\mathcal{S}})$, are there only a finite number of distinct subvarieties $W(p)$?

Fix a number field K and a place \mathfrak{p} . If Question 1.1 has a positive answer for K and $\mathcal{S} := \{\mathfrak{p}\}$, then there does not exist an infinite, \mathfrak{p} -adically discrete, Diophantine subset of K . See [15, Proof of Proposition 1.5] for the proof.

So one way to answer Question 1.1 (negatively) for K would be to construct a Diophantine definition of an infinite discrete \mathfrak{p} -adic set over a number field K . Unfortunately, at the moment such a construction seems out of reach. So instead we consider analogs in which K is replaced by one of its large integrally closed subrings $\mathcal{O}_{K,S}$:

Definition 1.3. For a number field K , let \mathcal{P}_K denote the set of finite primes of K , and let \mathcal{O}_K denote the ring of integers. Given a set S of prime ideals, not necessarily finite, the ring $\mathcal{O}_{K,S}$ is defined to be the subring of K defined by

$$\mathcal{O}_{K,S} = \{x \in K : \text{ord}_{\mathfrak{p}} x \geq 0 \text{ for all } \mathfrak{p} \notin S\}.$$

Observe that if $S = \emptyset$, then $\mathcal{O}_{K,S} = \mathcal{O}_K$ and if $S = \mathcal{P}_K$, then $\mathcal{O}_{K,S} = K$. If S is finite, $\mathcal{O}_{K,S}$ is called a *ring of S -integers*. In the case where the complement of S is finite, the rings $\mathcal{O}_{K,S}$ are semi-local. We will call all rings $\mathcal{O}_{K,S}$ with infinite S *big rings*.

To measure the “size” of a set of primes one can use natural density defined below.

Definition 1.4. Let $S \subseteq \mathcal{P}_K$. The *natural density* of S is defined to be the limit

$$\lim_{X \rightarrow \infty} \frac{\#\{\mathfrak{p} \in S : N\mathfrak{p} \leq X\}}{\#\{\text{all } \mathfrak{p} : N\mathfrak{p} \leq X\}}$$

if it exists. If the limit above does not exist, one can talk about *upper* density by substituting \limsup for \lim , or *lower* density by substituting \liminf for \lim .

The study of Hilbert’s Tenth Problem and of the archimedean version of Mazur’s conjecture over rings of S -integers has produced Diophantine definitions of \mathbb{Z} and discrete archimedean sets over large subrings of some number fields [20–24, 26]. In 2003, Poonen proved that there exists a recursive set S of primes of natural density one such that Hilbert’s Tenth Problem is undecidable for $\mathbb{Z}[S^{-1}]$. He also constructed an infinite discrete Diophantine set (in the archimedean topology) in this ring. In [15] Poonen and Shlapentokh prove that, if there exists an elliptic curve E over a number field K with $\text{rank}(E(K)) = 1$, then there exists a recursive set S of primes of density one such that Hilbert’s Tenth Problem is undecidable for $\mathcal{O}_{K,S}$. They also show that there is an infinite Diophantine subset A of $\mathcal{O}_{K,S}$ such that for all places v of K , the set A is discrete when viewed as a subset of the completion K_v .

In [6], Eisenträger and Everest reconsidered the original result of Poonen from a different point of view, looking for a “covering” of \mathbb{Q} by big rings that come from complementary sets of primes. More specifically, they proved that the rational primes can be partitioned into two disjoint sets S_1, S_2 such that Hilbert’s Tenth Problem is undecidable over both \mathcal{O}_{K,S_1} and \mathcal{O}_{K,S_2} . These results were improved by Perlega in [13] to show that the two sets can be of arbitrary computable densities.

In this paper, we generalize the results of [15], [6] and [13] to prove the following theorems:

Theorem 1.1. *Let K be a number field, and assume there is an elliptic curve defined over K with K -rank equal to 1. For every $t > 1$ and every collection $\delta_1, \dots, \delta_t$ of nonnegative computable real numbers adding up to 1, the set of the nonarchimedean valuations of K may be partitioned into t mutually disjoint recursive subsets S_1, \dots, S_t of natural densities $\delta_1, \dots, \delta_t$, respectively, with the property that each ring \mathcal{O}_{K,S_i} contains a Diophantine subset discrete under any valuation of K (archimedean or nonarchimedean).*

Theorem 1.2. *Assume there is an elliptic curve defined over K with K -rank equal to 1. For every $t > 1$ and every collection $\delta_1, \dots, \delta_t$ of nonnegative computable real numbers adding up to 1, the set of the nonarchimedean valuations of K may be partitioned into t mutually disjoint recursive subsets S_1, \dots, S_t of natural densities $\delta_1, \dots, \delta_t$, respectively, with the property that \mathbb{Z} admits a Diophantine model in each ring \mathcal{O}_{K,S_i} . In particular, Hilbert’s Tenth Problem is undecidable for each ring \mathcal{O}_{K,S_i} .*

Recently, Mazur and Rubin [12] showed that if the Shafarevich–Tate conjecture holds, then there always exists an elliptic curve defined over K whose K -rank is one.

When proving Theorems 1.1 and 1.2, we will show that given any partition of the nonarchimedean primes into sets W_1, \dots, W_t of densities $\delta_1, \dots, \delta_t$, the sets S_i can be constructed by changing the W_i ’s by sets of density zero. So our results can be seen as answering the following fundamental questions *up to sets of density zero*:

Questions 1.2.

- (1) For which number fields K and which subsets S of \mathcal{P}_K is Hilbert's Tenth Problem (un)decidable over $\mathcal{O}_{K,S}$?
- (2) For which number fields K and which subsets S of \mathcal{P}_K is there a Diophantine model of \mathbb{Z} over $\mathcal{O}_{K,S}$?
- (3) For which number fields K and subsets S of \mathcal{P}_K is there an infinite subset of $\mathcal{O}_{K,S}$ which is Diophantine over $\mathcal{O}_{K,S}$ and discrete in every topology of the field K ?

One question which is not addressed by this paper is for which number fields K and which subsets S of \mathcal{P}_K there is a Diophantine definition of \mathbb{Z} (or \mathcal{O}_K) over $\mathcal{O}_{K,S}$.

1.1. Overview of proof. The goal is to prove Theorems 1.1 and 1.2 by partitioning \mathcal{P}_K into t disjoint sets S_1, \dots, S_t , so that each ring \mathcal{O}_{K,S_r} admits a Diophantine model of the integers or has discrete infinite Diophantine subsets. In Sections 6 and 7, we first show how to find t not necessarily disjoint sets, whose union is \mathcal{P}_K such that the corresponding big rings have desirable properties. In Section 8, we show that these sets can also be chosen to be mutually disjoint and of the required density.

To construct infinite discrete Diophantine sets we will proceed as in [15] and construct a Diophantine set containing only the elements of a sequence converging (in all topologies of the number field) to a limit not in the set.

To construct a Diophantine model of \mathbb{Z} inside \mathcal{O}_{K,S_r} , it is enough to construct a model of the structure

$$\mathcal{Z} := (\mathbb{Z}_{\geq 1}, 1, +, B),$$

where B is a unary predicate for the set $\{2^n + n^2 : n \in \mathbb{Z}_{\geq 1}\}$ (see [15, Lemma 3.16]). A *Diophantine model of \mathcal{Z}* over a ring R is a Diophantine subset $A \subseteq R^m$ for some m together with a bijection $\phi : \mathbb{Z}_{\geq 1} \rightarrow A$ such that $\phi(B)$ is Diophantine over A and such that the graph of addition (a subset of $\mathbb{Z}_{\geq 1}^3$) corresponds under ϕ to a Diophantine subset of A^3 .

In order to find suitable sets S_r we work with an elliptic curve E of rank one over K and a point P of infinite order that is a suitable multiple of the generator for the nontorsion part. We will construct t (infinite) sequences of primes

$$\{\ell_{1,1}, \ell_{2,1}, \dots\}, \dots, \{\ell_{1,t}, \ell_{2,t}, \dots\}$$

such that for each $r \in \{1, \dots, t\}$, we have that $E(\mathcal{O}_{K,S_r}) \cap zE(K)$ for a suitable positive integer z , is the union of $\{\pm \ell_{1,r}P, \pm \ell_{2,r}P, \dots\}$ and some finite set. We then show that $A_r := \{x_{\ell_{i,r}} : i \in \mathbb{Z}_{\geq 1}\}$ is a Diophantine model of \mathcal{Z} in \mathcal{O}_{K,S_r} via the bijection $\phi : \mathbb{Z}_{\geq 1} \rightarrow A_r$ sending i to $x_{\ell_{i,r}}$. To prove Theorem 1.1 we construct t different sequences of primes and sets S_r and show that A_r as above is a discrete Diophantine set.

The paper is organized as follows. In Section 2, we review recursive presentations of primes of number fields, in Section 3 we give some background about primitive divisors and their properties, and then use these properties to prove that certain terms in divisibility sequences have many prime ideal divisors. Section 4 describes the technical changes in the assumptions and proofs in this paper relative to proofs and assumptions in [15]. Section 5 reviews and extends some density results from [15]. In Sections 6 and 7, we construct the rings and the sets with the required properties.

Finally, Section 8 shows how to adjust the sets of primes constructed in Sections 6 and 7 to make them complementary.

2. Computable sets of primes in number fields

In this section, we briefly discuss a presentation of primes in number fields and a way to define recursive sets of primes. We assume that a number field K of degree n over \mathbb{Q} is presented in terms of its integral basis over \mathbb{Q} . (Such a basis always exists and can be constructed given an irreducible polynomial over \mathbb{Q} of a field generator. See, for example, Section 7.3 of [16].) Elements of the field will be presented via n -tuples of the coordinates with respect to the basis. Given a K -prime \mathfrak{p} , we will present this prime by a pair $(p, \alpha_{\mathfrak{p}})$, where p is the \mathbb{Q} -prime below \mathfrak{p} and $\alpha_{\mathfrak{p}} \in K$ is an algebraic integer such that $\text{ord}_{\mathfrak{p}} \alpha_{\mathfrak{p}} = 1$ but $\text{ord}_{\mathfrak{q}} \alpha_{\mathfrak{p}} = 0$ for any prime $\mathfrak{q} \neq \mathfrak{p}$ conjugate to \mathfrak{p} over \mathbb{Q} . Since the choice $\alpha_{\mathfrak{p}}$ is not unique we can choose the first suitable $\alpha_{\mathfrak{p}}$ under some ordering of the field. Given an integral basis for K , the map $p \mapsto (\alpha_{\mathfrak{p}_1}, \dots, \alpha_{\mathfrak{p}_m})$, where $p = \prod_{i=1}^k \mathfrak{p}_i^{e_i}$ is the factorization of p in K , is recursive. Further, given an element of K , one can effectively determine the factorization of the divisor of this element, and given a prime compute its norm. Given a set of primes we can now say that it is computable if the corresponding set of $(n+1)$ -tuples $(p, \alpha_{\mathfrak{p}})$ is computable. It is also not hard to see that for any set of K -primes \mathcal{W} , the ring $\mathcal{O}_{K, \mathcal{W}}$ from Definition 1.3 is computable if and only if \mathcal{W} is computable. For more details see Section 4 of [2].

3. Primitive divisors

Let E denote an elliptic curve in Weierstrass form,

$$(3.1) \quad E: y^2 = x^3 + a_4x + a_6,$$

defined over \mathcal{O}_K . For background, definitions and the properties of elliptic curves used in this paper, consult [27, 29]. Let K denote an algebraic number field of degree $d = [K : \mathbb{Q}]$ over \mathbb{Q} . Throughout the paper, $E(K)$ denotes the group of K -rational points of E and \mathbf{O} denotes the point at infinity, the identity for the group of K -rational points. Suppose P denotes a K -rational point, $P \in E(K)$, which is not torsion. Write $nP = (x_n, y_n)$. The assumptions on E allow the factorization

$$(3.2) \quad (x_n) = (x(nP)) = \mathfrak{a}_n(P) / \mathfrak{b}_n^2(P)$$

of the principal fractional ideal $(x(nP))$ into relatively prime integral ideals \mathfrak{a}_n and \mathfrak{b}_n . Assuming P is nontorsion guarantees that all of the terms in the sequence $\mathfrak{b} = (\mathfrak{b}_n)$ are nonzero.

In the rational case, we may take \mathfrak{b}_n to be a positive integer. Silverman [28] proved that when P is a rational point, for all sufficiently large n , we have that \mathfrak{b}_n has a *primitive divisor*, that is, a divisor of \mathfrak{b}_n which is coprime to \mathfrak{b}_m for all positive integers $m < n$. In general, the expression *primitive ideal divisor* of a term \mathfrak{b}_n is used to describe an ideal \mathcal{I} which divides \mathfrak{b}_n but no \mathfrak{b}_m with $m < n$. Cheon and Hahn [3] extended Silverman's result from [28] to algebraic number fields, showing that for all sufficiently large n , it is the case that \mathfrak{b}_n has a primitive ideal divisor.

Results about primitive divisors have a long and fine tradition for certain sequences which satisfy a linear recurrence relation. An interested reader can find more results concerning the existence of primitive divisors in [1, 7, 17–19, 30].

For a point P on E and a nonzero integer n , define $\mathcal{S}_n(P)$ to be the set of all prime ideals of \mathcal{O}_K that divide the ideal $\mathfrak{b}_n(P)$.

We will use the following properties of the sequence $\mathfrak{b}_n(P)$ and the sets $\mathcal{S}_n(P)$:

Lemma 3.1. *Let P be a point of infinite order on an elliptic curve E defined over a number field K as above.*

- (1) *Let $n, m \in \mathbb{Z} - \{0\}$ and let (m, n) be their gcd. Then $\mathcal{S}_m(P) \cap \mathcal{S}_n(P) = \mathcal{S}_{(m, n)}(P)$. In particular, if P is an integral point of infinite order, and $(m, n) = 1$, then $\mathcal{S}_m(P) \cap \mathcal{S}_n(P) = \emptyset$.*
- (2) *The sequence \mathfrak{b}_n is a divisibility sequence, meaning that $\mathfrak{b}_m \mid \mathfrak{b}_n$ as ideals, whenever $m \mid n$.*

Proof. The proof follows from the standard local theory of elliptic curves, see for example Chapters 4 and 7 in [27]: for $\mathfrak{p} \in \mathcal{P}_K$, let $K_{\mathfrak{p}}$ be the completion of K at \mathfrak{p} and let

$$E_1(K_{\mathfrak{p}}) = \{O\} \cup \{R \in E(K_{\mathfrak{p}}) : \text{ord}_{\mathfrak{p}}(x(R)) \leq -2\}.$$

From [27, Proposition VII.2.2], we have that $E_1(K)$ with the elliptic curve addition is isomorphic to $\hat{E}(\mathcal{M})$ under the formal group addition, where \hat{E} is the formal group associated to E and \mathcal{M} is the valuation ideal of $K_{\mathfrak{p}}$. Note that while the proposition in [27] assumes that the Weierstrass equation is minimal, this assumption is not used in the proof. The first assertion of the lemma now follows from the fact that $E_1(K_{\mathfrak{p}})$ is a group.

Furthermore, using again the fact that the elliptic curve addition on $E_1(K_{\mathfrak{p}})$ corresponds to the addition in the formal group [27, Proposition VII.2.2], together with [27, Corollary IV.4.4], we have that for all $R \in E_1(K_{\mathfrak{p}}) - \{O\}$

$$\text{ord}_{\mathfrak{p}}(x(nR)) \leq \text{ord}_{\mathfrak{p}}(x(R)) - 2\text{ord}_{\mathfrak{p}}(n)$$

and the second assertion of the lemma follows at once. \square

To carry out our construction we need to prove that certain terms in the sequence $\mathfrak{b}_n(P)$ have *many* primitive ideal divisors. This is made precise in the next theorem.

Theorem 3.1. *Let p denote a prime and write $q = p^{t-1}$ for some fixed $t \geq 2$. Suppose Q is a K -rational point of infinite order and $P = qQ$. Let $\{(\mathfrak{b}_m)(P)\}$ be the sequence of ideals coming from the multiples of P as in equation (3.2). For every large enough n , which is coprime to p , the term $\mathfrak{b}_n(P)$ has at least t primitive ideal divisors. The same is true for the terms of the sequence $\mathfrak{b}_n(pP)$.*

Proof. Let n be an integer coprime to p and assume that n is large enough so that $\mathfrak{b}_k(Q)$ has a primitive divisor for all $k > n$. Let $\mathfrak{p}_{p^i n}$ be a primitive prime ideal divisor of $\mathfrak{b}_{p^i n}(Q)$, for $i = 0, \dots, t-1$. Observe that for $i \neq j$ we have that $\mathfrak{p}_{p^i n} \neq \mathfrak{p}_{p^j n}$. We claim that

$$\mathfrak{p}_{p^i n} \in \mathcal{S}_{p^{t-1}n}(Q) - \mathcal{S}_{p^{t-1}m}(Q) = \mathcal{S}_n(P) - \mathcal{S}_m(P)$$

for any positive $m < n$. Indeed, since $p^i n$ divides $p^{t-1}n$ we have that $\mathfrak{p}_{p^i n} \in \mathcal{S}_{p^{t-1}n}(Q)$. Suppose also $\mathfrak{p}_{p^i n} \in \mathcal{S}_{p^{t-1}m}(Q)$, where $m < n$. By Lemma 3.1, part (1), we can assume without loss of generality that m divides n and thus is prime to p . We now also have that $\mathfrak{p}_{p^i n} \in \mathcal{S}_{p^{t-1}m}(Q) \cap \mathcal{S}_{p^i n}(Q) = \mathcal{S}_{p^i m}(Q)$ contradicting the assumption that $\mathfrak{p}_{p^i n}$ is a primitive prime ideal divisor of $\mathfrak{b}_{p^i n}(Q)$. Thus $\mathfrak{p}_{p^i n}, i = 0, \dots, t-1$ are primitive ideal divisors of $\mathfrak{b}_n(p^{t-1}Q)$.

Similarly, $\mathfrak{p}_{p^i n} \in \mathcal{S}_{p^t n}(Q) - \mathcal{S}_{p^t m}(Q)$ for any positive $m < n$. Indeed, as above, since $p^i n$ divides $p^t n$ we have that $\mathfrak{p}_{p^i n} \in \mathcal{S}_{p^t n}(Q)$. Suppose also $\mathfrak{p}_{p^i n} \in \mathcal{S}_{p^t m}(Q)$, where $0 < m < n$. Again, by Lemma 3.1, part (1), we can assume without loss of generality that m divides n and thus is prime to p . We now also have that $\mathfrak{p}_{p^i n} \in \mathcal{S}_{p^t m}(Q) \cap \mathcal{S}_{p^i n}(Q) = \mathcal{S}_{p^i m}(Q)$, contradicting the assumption that $\mathfrak{p}_{p^i n}$ is a primitive prime ideal divisor of $\mathfrak{b}_{p^i n}(Q)$. Thus $\mathfrak{p}_{p^i n}, i = 0, \dots, t-1$ are primitive ideal divisors of $\mathfrak{b}_n(p^t Q)$. \square

4. Some technical matters

Below we construct two collections of rings \mathcal{O}_{K,S_i} : one to produce infinite discrete Diophantine sets and the other to construct a Diophantine model of the integers. The rings \mathcal{O}_{K,S_i} are constructed by generalizing the techniques from [15]. For the most part we use the same notation as in [15], but with the following modifications:

In [15], the authors define $\mathcal{S}_{\text{bad}} \subseteq \mathcal{P}_K$ to be the set of primes that ramify in K/\mathbb{Q} , the primes for which the reduction of the chosen Weierstrass model is singular (this includes all primes above 2), and the primes at which the coordinates of P are not integral. In the rings in [15] for which undecidability is then shown the primes in \mathcal{S}_{bad} are always inverted. That is, the rings are of the form $\mathcal{O}_{K,S}$ with $\mathcal{S}_{\text{bad}} \subseteq S$. We have to avoid inverting the primes in \mathcal{S}_{bad} in each ring, otherwise the sets S_i will not be mutually disjoint. That means that in our paper the fractional ideal generated by the x -coordinate of nP is of the form $x(nP) = \mathfrak{a}_n/\mathfrak{d}_n$ (with $\mathfrak{a}_n, \mathfrak{d}_n$ coprime integral ideals) and we do not have a separate ideal \mathfrak{b}_n that includes the contribution from the primes in \mathcal{S}_{bad} as in [15].

In view of the above, we need to show that the undecidability results in [15] can be proved without inverting the primes in \mathcal{S}_{bad} . Below we note that (1) P can be chosen to be integral, that (2) we can avoid inverting the primes that ramify in K/\mathbb{Q} and (3) that we can avoid inverting the primes for which the reduction of the Weierstrass model of E is singular:

- (1) We assume that the point $P := zQ$ has coordinates in \mathcal{O}_K . Here Q generates $E(K)/E(K)_{\text{tors}}$ and $z = 2^{t-1}3^{t-1}\#E(K)_{\text{tors}}$. This assumption is possible by Lemma 4.1 below. Our assumption implies that the point P does not contribute any primes to \mathcal{S}_{bad} .
- (2) Not inverting the primes that ramify in K/\mathbb{Q} . The fact that \mathcal{S}_{bad} contains the primes that ramify in K/\mathbb{Q} is used in [15] to prove Lemma 3.3, which is then used to prove Proposition 3.5 in [15]. Our proof below replaces Lemma 3.3 and Proposition 3.5 from [15] with Lemma 3.1 and Theorem 3.1.
- (3) Not inverting the primes of bad reduction. Our definitions of $\mathcal{T}_1, \mathcal{S}_n, \mathfrak{p}_n, \mathcal{T}_2$ differ from those in [15]: our set \mathcal{T}_1 is contained in the set \mathcal{T}_1 defined in [15], and it differs from it by at most finitely many primes (the primes in \mathcal{S}_{bad}). Our set \mathcal{S}_n contains *all* prime ideals dividing the denominator ideal of $x(nP)$,

and $\mathfrak{p}_n = \mathfrak{p}_n^{(1)}$ denotes a *primitive* prime ideal divisor of the largest norm in \mathcal{S}_n . This also affects the definition of \mathcal{T}_2 . See Notation 4.1 and the sets that are defined before Lemmas 6.2 and 7.2 below.

The primes of bad reduction are relevant in Lemma 3.1 and Corollary 3.2 of [15]. Since we have a different definition of $\mathfrak{p}_\ell^{(1)}$ we do not need to use these two results. The only other place in [15] where primes of bad reduction are relevant is Lemma 3.10, and we state below why this lemma still holds (see Lemmas 6.2 and 7.2 and their proofs).

Lemma 4.1. *If E is an elliptic curve and $P \in E(K)$, then there exists a curve E' that is isomorphic to E over K via an isomorphism ϕ such that $P' := \phi(P)$ has coordinates in \mathcal{O}_K .*

Proof. If E is given by a Weierstrass equation $E : y^2 = x^3 + ax + b$ and $P \in E(K)$ has coordinates $(\alpha, \beta) \in K$, we can choose an element $u \in K$ such that $u\alpha, u\beta \in \mathcal{O}_K$. We can then consider the curve E' whose Weierstrass equation is given by

$$E' : (y')^2 = (x')^3 + au^4(x') + u^6b,$$

which is isomorphic to E under $\phi : E \rightarrow E', (x, y) \mapsto (u^2x, u^3y)$. The point $P' := \phi(P)$ on E' has coordinates in \mathcal{O}_K . \square

Now we can fix some of our notation:

4.1. Notation.

- Let K be a number field.
- Let E be an elliptic curve of rank 1 over K , given by a Weierstrass equation with coefficients in the ring of integers \mathcal{O}_K . (In particular, we assume that K is such that such an E exists).
- Let $E(K)_{\text{tors}}$ be the torsion subgroup of $E(K)$.
- For any set S of K -primes let $\tilde{E}(\mathcal{O}_{K,S})$ be the set of affine points with coordinates in $\mathcal{O}_{K,S}$.
- Let $z = 2^{t-1}3^{t-1}\#E(K)_{\text{tors}}$ with $t \geq 1$.
- $P := zQ$, where Q generates $E(K)/E(K)_{\text{tors}}$. As explained above, we may assume $P = (x, y)$ with $x, y \in \mathcal{O}_K$.
- Let $\mathcal{P}_{\mathbb{Q}} = \{2, 3, 5, \dots\}$ be the set of rational primes.
- Let \mathcal{P}_K be the set of all finite primes of K .
- For $\mathfrak{p} \in \mathcal{P}_K$, let
 - $K_{\mathfrak{p}}$ be the completion of K at \mathfrak{p} .
 - $R_{\mathfrak{p}}$ be the valuation ring of $K_{\mathfrak{p}}$
 - $\mathbb{F}_{\mathfrak{p}}$ be the residue field of $R_{\mathfrak{p}}$,
 - $\mathbf{N}\mathfrak{p} = \#\mathbb{F}_{\mathfrak{p}}$ be the absolute norm of \mathfrak{p}
- For $n \neq 0$ write $nP = (x_n, y_n)$ where $x_n, y_n \in K$.
- Write the fractional ideal generated by x_n as

$$(x_n) = \frac{\mathfrak{a}_n}{\mathfrak{d}_n},$$

where \mathfrak{a}_n and \mathfrak{d}_n are coprime integral ideals.

- For n as above, let $\mathcal{S}_n = \mathcal{S}_n(P) = \{\mathfrak{p} \in \mathcal{P}_K : \mathfrak{p} | \mathfrak{d}_n\}$. By assumption on P , we have $\mathcal{S}_1 = \emptyset$.

- For $\ell \in \mathcal{P}_{\mathbb{Q}}$, define a_{ℓ} to be the smallest positive number such that $\mathfrak{d}_{\ell^{a_{\ell}}}$ has at least t primitive divisors. (By Theorem 3.1, applied with $p = 2$ for $\ell \neq 2$ and with $p = 3$ for $\ell = 2$, we have that a_{ℓ} exists and $a_{\ell} = 1$ for all but finitely many ℓ .)
- Let $\mathcal{L} = \{\ell \in \mathcal{P}_{\mathbb{Q}} : a_{\ell} > 1\}$ and $L = \prod_{\ell \in \mathcal{L}} \ell^{a_{\ell}-1}$.
- For $k = 1, \dots, t$ define $\mathfrak{p}_n^{(k)}$ to be the k th largest primitive prime divisor of \mathfrak{d}_n (if it exists). (Order the primitive prime divisors according to their norm, and break ties for prime ideals $\mathfrak{p}_1, \mathfrak{p}_2$ of the same norm according to Section 2: compute the corresponding $\alpha_{\mathfrak{p}_1}, \alpha_{\mathfrak{p}_2}$ and see which one comes first under some ordering of the field.)
- For a prime ℓ , define

$$\mu_{\ell} = \sup_{X \in \mathbb{Z}_{\geq 2}} \frac{\#\{\mathfrak{p} \in \mathcal{S}_{\ell} : \mathbf{N}\mathfrak{p} \leq X\}}{\#\{\mathfrak{p} \in \mathcal{P}_K : \mathbf{N}\mathfrak{p} \leq X\}}.$$

- Let \mathcal{M}_K be the set of all normalized absolute values of K .
- Let $\mathcal{M}_{K,\infty} \subset \mathcal{M}_K$ be the set of all archimedean absolute values of K .

5. On densities of some sets of primes

The main result of this section is the proposition below.

Proposition 5.1. *The natural density of the set $\mathcal{Q}(E) = \{\mathfrak{q}_{\ell}, \ell \in \mathbb{Z}_{>0}\}$, where \mathfrak{q}_{ℓ} is any primitive divisor of $[\ell]P$ (see Notation 4.1), is zero.*

In [15], it was shown that the set $\{\mathfrak{p}_{\ell}, \ell \in \mathbb{Z}_{>0}\}$, where \mathfrak{p}_{ℓ} is the *largest* primitive divisor of P_{ℓ} , is equal to zero. Below we modify this proof and show that the primitive divisor does not have to be the largest in order for the density to be zero. The key result we need from [15] is stated below.

Lemma 5.1. *For $n \in \mathbb{Z}_{>0}$, let $\omega(n)$ be the number of distinct prime factors of n . For any $t \geq 1$, the density of $\mathcal{Z}(E, t) = \{\mathfrak{p} : \omega(\#E(\mathbb{F}_{\mathfrak{p}})) < t\}$ is 0. (See Lemma 3.12 of [15].)*

As in [15] and [14] we also need the following result and an observation.

Theorem 5.1 (Hasse). $\#E(\mathbb{F}_{\mathfrak{p}}) \leq \mathbf{N}\mathfrak{p} + 1 + 2\sqrt{\mathbf{N}\mathfrak{p}}$.

Remark 5.1. If \mathfrak{p} is a prime at which E has a good reduction and such that \mathfrak{p} is a primitive divisor of ℓP , then $\ell | \#E(\mathbb{F}_{\mathfrak{p}})$. Note that since there are only finitely many primes at which E has a bad reduction, we can ignore these primes when calculating the density.

We now prove Proposition 5.1.

Proof. We choose $\varepsilon > 0$ and show that the upper natural density of $\mathcal{Q}(E)$ is less than ε . By the Prime Number Theorem, for some positive constants $C_{\mathbb{Q}}, C_K$ we have

$$\begin{aligned} \#\{p \in \mathcal{P}_{\mathbb{Q}} : p \leq X\} &= O(X/\log X) < \frac{C_{\mathbb{Q}}X}{\log X}, \\ \#\{\mathfrak{p} \in \mathcal{P}_K : \mathbf{N}\mathfrak{p} \leq X\} &= O(X/\log X) > \frac{C_KX}{\log X}. \end{aligned}$$

Choose $t \in \mathbb{Z}_{>1}$ so that

$$2^{4-t} < \frac{C_K \varepsilon}{4C_{\mathbb{Q}}}$$

and choose $X \in \mathbb{R}_{>0}$ large enough so that

$$\frac{\#\{\mathfrak{p} \in \mathcal{Z}(E, t), \mathbf{N}\mathfrak{p} \leq X\}}{\#\{\mathfrak{p} \in \mathcal{P}_K, \mathbf{N}\mathfrak{p} \leq X\}} < \varepsilon/2,$$

and

$$\frac{|\log C_K + \log \varepsilon - \log 4C_{\mathbb{Q}}|}{\log X} < 1.$$

Let $\overline{\mathcal{Z}(E, t)}$ be the complement of $\mathcal{Z}(E, t)$ in \mathcal{P}_K . Let $\mathfrak{p} \in \overline{\mathcal{Z}(E, t)}$ and assume $\mathfrak{p} = \mathfrak{q}_{\ell}$ for some positive integer ℓ . In this case,

$$\ell 2^t < \#E(\mathbb{F}_{\mathfrak{p}}) < \mathbf{N}\mathfrak{p} + 1 + 2\sqrt{\mathbf{N}\mathfrak{p}} < 4\mathbf{N}\mathfrak{p}$$

and therefore

$$\ell < 2^{4-t} \mathbf{N}\mathfrak{p} \leq \frac{C_K \mathbf{N}\mathfrak{p} \varepsilon}{4C_{\mathbb{Q}}}.$$

Thus for every $\mathfrak{p} \in \mathcal{Q}(E) \cap \overline{\mathcal{Z}(E, t)}$ there exists a unique rational prime $\ell < \frac{C_K \mathbf{N}\mathfrak{p} \varepsilon}{4C_{\mathbb{Q}}}$. Consider now the following ratio:

$$\begin{aligned} & \frac{\#\{\mathfrak{p} \in \mathcal{Q}(E) : \mathbf{N}\mathfrak{p} \leq X\}}{\#\{\mathfrak{p} \in \mathcal{P}_K : \mathbf{N}\mathfrak{p} \leq X\}} \\ &= \frac{\#\{\mathfrak{p} \in \mathcal{Q}(E) \cap \mathcal{Z}(E, t) : \mathbf{N}\mathfrak{p} \leq X\}}{\#\{\mathfrak{p} \in \mathcal{P}_K : \mathbf{N}\mathfrak{p} \leq X\}} + \frac{\#\{\mathfrak{p} \in \mathcal{Q}(E) \cap \overline{\mathcal{Z}(E, t)} : \mathbf{N}\mathfrak{p} \leq X\}}{\#\{\mathfrak{p} \in \mathcal{P}_K : \mathbf{N}\mathfrak{p} \leq X\}} \\ &\leq \varepsilon/2 + \frac{\#\{\ell \in \mathcal{P}_{\mathbb{Q}} : \ell \leq \frac{C_K \varepsilon X}{4C_{\mathbb{Q}}}\}}{\#\{\mathfrak{p} \in \mathcal{P}_K : \mathbf{N}\mathfrak{p} \leq X\}} \leq \varepsilon/2 + \frac{\frac{C_{\mathbb{Q}} C_K \varepsilon X}{4C_{\mathbb{Q}} \log(C_K \varepsilon X / 4C_{\mathbb{Q}})}}{\frac{C_K X}{\log X}} \\ &= \varepsilon/2 + \frac{\varepsilon \log X}{4(\log C_K + \log \varepsilon + \log X - \log 4C_{\mathbb{Q}})} < \varepsilon. \end{aligned}$$

□

Now we show that it is rare that \mathcal{S}_{ℓ} has a large fraction of the small primes.

Lemma 5.2. *For any $\varepsilon > 0$, the density of $\{\ell : \mu_{\ell} > \varepsilon\}$ is 0.*

Proof. The statement of this lemma is identical to the statement of Lemma 3.8 of [15] except for the fact that in our case \mathcal{S}_{ℓ} can contain primes of S_{bad} . However by Lemma 3.1, only finitely many ℓ can be affected by the inclusion of S_{bad} primes and therefore the density result is unaffected. □

The next lemma is Lemma 3.6 of [15] which we restate here without a proof.

Lemma 5.3. *Let $\vec{\alpha} \in \mathbb{R}^n$, let I be an open neighborhood of 0 in $\mathbb{R}^n/\mathbb{Z}^n$, and let $d \in \mathbb{Z}_{\geq 1}$. Then the set of primes $\ell \equiv 1 \pmod{d}$ such that $(\ell - 1)\vec{\alpha} \pmod{1}$ is in I has positive lower density.*

6. Infinite Diophantine discrete sets

In this section, we construct t distinct sequences of primes from which we will construct the sets S_1, \dots, S_t . We start with a lemma which will enable us to show that the sequences we construct are computable.

Lemma 6.1. *Let $\mathfrak{p}_\ell^{(k)}$ and μ_ℓ be as in Notation 4.1.*

- (1) *For all $k = 1, \dots, t$, the mapping $\ell \mapsto \mathfrak{p}_\ell^{(k)}$ is computable.*
- (2) *The mapping $\ell \mapsto \mu_\ell$ is computable.*

Proof.

- (1) Given $k, \ell \in \mathbb{Z}_{>0}$ we can effectively compute the coordinates of $x_\ell = x(\ell(P))$ and determine the factorization of \mathfrak{d}_ℓ as discussed in the Section 2. By considering the prime factorization of $\mathfrak{d}_1, \dots, \mathfrak{d}_{\ell-1}$ we can determine which primes occurring in \mathfrak{d}_ℓ are in fact primitive divisors, compute their norms and determine $\mathfrak{p}_\ell^{(k)}$.
- (2) First of all, as above, for any $\ell > 0$ we can effectively determine all the primes in S_ℓ and compute their norm. Secondly, once X in the definition of μ_ℓ is greater than the norm of $\mathfrak{p}_\ell^{(1)}$, the value of the ratio can only decline. Thus to compute μ_ℓ it sufficient to calculate the ratio for finitely many values of X only. Therefore, μ_ℓ can be computed effectively.

□

By [27, Corollary VI.5.1.1] and [29, Corollary V.2.3.1] there is an isomorphism of real Lie groups $\prod_{v \in \mathcal{M}_{K,\infty}} E(K_v) \simeq (\mathbb{R}/\mathbb{Z})^N \times (\mathbb{Z}/2\mathbb{Z})^{N'}$ for some $N \geq 1$ and $N' \geq 0$. Fix such an isomorphism, and embed $E(K)$ diagonally in $\prod_{v \in \mathcal{M}_{K,\infty}} E(K_v)$. Since $P = zQ$ with z even, the point P maps to an element $\vec{\alpha} \in (\mathbb{R}/\mathbb{Z})^N$.

Now we construct the sequences $\{\ell_{1,r}, \ell_{2,r}, \dots\}$ for $r = 1, \dots, t$. To do this we describe how to define $\ell_{i,r}$ using a set $V_{i,r}, i \in \mathbb{Z}_{>0}, r = 1, \dots, t$ of previously defined elements of the sequences. More specifically we let $V_{1,1} = \emptyset$. For $i > 1$ we set

$$V_{i,1} = \{\ell_{1,1}, \dots, \ell_{1,t}, \dots, \ell_{i-1,1}, \dots, \ell_{i-1,t}\},$$

and for $i \geq 1, 1 < r \leq t$, we set

$$V_{i,r} = \{\ell_{1,1}, \dots, \ell_{1,t}, \dots, \ell_{i,1}, \dots, \ell_{i,r-1}\}.$$

Let $\ell_{i,r}$ be the smallest prime outside \mathcal{L} and exceeding the bound implicit in Theorem 3.1 such that all of the following hold:

- (1) $\ell_{i,r} > \ell$, for all $\ell \in V_{i,r}$,
- (2) $\mu_{\ell_{i,r}} \leq 2^{-i}$,
- (3) $\mathbf{Np}_{\ell_{i,r}}^{(r)} > 2^i$, for all $\ell \in V_{i,r} \cup \{\ell_{i,r}\} \cup \mathcal{L}$,
- (4) $\ell_{i,r} \equiv 1 \pmod{i!}$, and
- (5) $|x_{\ell_{i,r}-1}|_v > i$, for all $v \in \mathcal{M}_{K,\infty}$.

We also choose $\ell_{1,1} > 3$.

Proposition 6.1. *The sequences $\{\ell_{1,r}, \ell_{2,r}, \dots\}$ are well defined and computable for $r = 1, \dots, t$.*

Proof. Condition (5) is equivalent to the requirement that $(\ell-1)\bar{\alpha}$ lie in a certain open neighborhood of 0 in $(\mathbb{R}/\mathbb{Z})^N$, since the Lie group isomorphism maps neighborhoods of O to neighborhoods of 0. Thus by Lemma 5.3, the set of primes satisfying (4) and (5) has positive lower density. By Lemma 5.2, (2) fails for a set of density 0. Therefore it will suffice to show that (1) and (3) are satisfied by all sufficiently large ℓ_i .

For fixed ℓ , the primes $\mathfrak{p}_{\ell\ell_{i,r}}^{(r)}$ for varying values of $\ell_{i,r}$ are distinct since $\mathfrak{p}_{\ell\ell_{i,r}}^{(r)}$ is the r -th largest primitive prime divisor of $\mathfrak{d}_{\ell\ell_{i,r}}$. So eventually their norms are greater than 2^i . The same holds for $\mathfrak{p}_{\ell\ell_{i,r}}^{(r)}$ for fixed $\ell \in \mathcal{L}$. Thus by taking $\ell_{i,r}$ sufficiently large, we can make all the $\mathfrak{p}_{\ell\ell_{i,r}}^{(r)}$ for $\ell = \ell_{i,r}$ or $\ell \in \mathcal{L}$ or $\ell \in V_{i,r}$ have norm greater than 2^i . Thus the sequence is well defined.

Each $\ell_{i,r}$ can be computed by searching primes in increasing order until one is found satisfying the conditions: conditions (1)–(4) can be verified effectively by Lemma 6.1, and condition (5) can be tested effectively, since $|x_{\ell_{i,r}-1}|_v$ is an algebraic real number. \square

We now define the following subsets of \mathcal{P}_K :

- $\mathcal{T}_{1,r} = \bigcup_{i \geq 1} \mathcal{S}_{\ell_{i,r}}, r = 1, \dots, t$;
- $\mathcal{T}_{2,r}^a$ is the set of $\mathfrak{p}_{\ell}^{(r)}$ for $\ell \notin (\{\ell_{1,r}, \ell_{2,r}, \dots\} \cup \mathcal{L})$, together with $\mathfrak{p}_{\ell_{a\ell}}^{(r)}$ for $\ell \in \mathcal{L}$;
- $\mathcal{T}_{2,r}^b = \{\mathfrak{p}_{\ell_{i,r}\ell_{j,r}}^{(r)} : 1 \leq j \leq i\}$;
- $\mathcal{T}_{2,r}^c = \{\mathfrak{p}_{\ell\ell_{i,r}}^{(r)} : \ell \in \mathcal{L}, i \geq 1\}$; and
- $\mathcal{T}_{2,r} = \mathcal{T}_{2,r}^a \cup \mathcal{T}_{2,r}^b \cup \mathcal{T}_{2,r}^c$.

By construction, all the terms considered above have t primitive divisors.

We now describe the important properties of these sequences.

Lemma 6.2.

- (1) *For each $r = 1, \dots, t$, the sets $\mathcal{T}_{1,r}$ and $\mathcal{T}_{2,r}$ are disjoint. If a subset $S_r \subset \mathcal{P}_K$ contains $\mathcal{T}_{1,r}$ and is disjoint from $\mathcal{T}_{2,r}$, then $\mathcal{E}_r := \tilde{E}(\mathcal{O}_{K,S_r}) \cap zE(K)$ is the union of*

$$\{\pm \ell_{i,r} P : i \geq 1\}$$

and some subset of the finite set $\{sP : s \mid \prod_{\ell \in \mathcal{L}} \ell^{a_\ell - 1}\}$.

- (2) *For any $j \in \{1, 2\}$ and $r, s \in \{1, \dots, t\}$ such that $r \neq s$ the sets $\mathcal{T}_{j,r}$ and $\mathcal{T}_{j,s}$ are disjoint.*
- (3) *For any $i \in \{1, 2\}$ and $r \in \{1, \dots, t\}$ the set $\mathcal{T}_{i,r}$ is computable.*

Proof.

- (1) The proof of this assertion is the same as the proof of Lemma 3.10 of [15]. The proof is not affected by the fact that we do not invert primes in S_{bad} , since in our case $S_1 = \emptyset$ also.

(2) First we assume that $j = 1$. In this case

$$\mathcal{T}_{1,r} \cap \mathcal{T}_{1,s} = \left(\bigcup_{i \geq 1} S_{\ell_{i,r}} \right) \cap \left(\bigcup_{i \geq 1} S_{\ell_{i,s}} \right) = \emptyset$$

since for any $r \neq s$ we have that $S_{\ell_{i_1,r}} \cap S_{\ell_{i_2,s}} = S_{(\ell_{i_1,r}, \ell_{i_2,s})} = \emptyset$ as $\{\ell_{i,r}\} \cap \{\ell_{i,s}\} = \emptyset$. Next let $j = 2$ and consider $\mathcal{T}_{2,r} \cap \mathcal{T}_{2,s}$. Since the set $\mathcal{T}_{2,r}$ consists of the r th largest primitive prime divisors of certain terms in the divisibility sequence \mathfrak{d}_n , and $\mathcal{T}_{2,s}$ consists of the s th largest primitive prime divisors of terms in the divisibility sequence, the definition of being a primitive divisor immediately implies that these sets can never have any nontrivial intersection when $r \neq s$.

(3) This assertion follows directly from the fact that each sequence:

$$\{\ell_{1,r}, \ell_{2,r}, \dots\}, 1 \leq r \leq t$$

is computable and from Lemma 6.1.

□

Proposition 6.2. *The natural density of $\mathcal{T}_{1,r}$ and $\mathcal{T}_{2,r}$ ($1 \leq r \leq t$) is zero.*

Proof. The proofs that $\mathcal{T}_{1,r}$, $\mathcal{T}_{2,r}^b$, and $\mathcal{T}_{2,r}^c$ have density 0 are identical to the proofs in Section 9 of [14]. The fact that $\mathcal{T}_{2,r}^a$ has density 0 follows from Proposition 5.1. □

Now we can construct infinite Diophantine subsets A_r of \mathcal{O}_{K,S_r} that are discrete in any topology of K . We first need the following lemma.

Lemma 6.3. *For each $v \in \mathcal{M}_K$ and $1 \leq r \leq t$ the sequence $\ell_{1,r}P, \ell_{2,r}P, \dots$ converges in $E(K_v)$ to P .*

Proof. This is Lemma 3.14 in [15]. □

We now have the following proposition.

Proposition 6.3. *Let S_r be as in Lemma 6.2 and let $A_r := \{x_{\ell_{1,r}}, x_{\ell_{2,r}}, \dots\}$. Then A_r is a Diophantine subset of \mathcal{O}_{K,S_r} . For any $v \in \mathcal{M}_K$, the set A_r is discrete when viewed as a subset of K_v .*

Proof. By Lemma 6.3, the elements of A_r form a convergent sequence in K_v whose limit x_1 is not in A_r . Hence A_r is discrete. By Lemma 6.2, part (1), $x(\mathcal{E}_r)$ is the union of the set A_r and a finite set. Since \mathcal{E}_r is Diophantine over \mathcal{O}_{K,S_r} , the set A_r is Diophantine over \mathcal{O}_{K,S_r} as well. □

In Section 8, we will use the sets A_1, \dots, A_t together with sets $\mathcal{T}_{1,1}, \dots, \mathcal{T}_{1,t}$ and $\mathcal{T}_{2,1}, \dots, \mathcal{T}_{2,t}$ to prove Theorem 1.1.

7. Constructing Diophantine models of \mathbb{Z} .

We will now modify the t sequences of primes constructed above so that in the resulting big rings Hilbert's Tenth Problem is undecidable.

Fix two primes $\mathfrak{p}, \mathfrak{q} \in \mathcal{P}_K$ of degree 1 that are primes of good reduction for E , and such that \mathfrak{p} and \mathfrak{q} do not ramify in K/\mathbb{Q} . Choose $\mathfrak{p}, \mathfrak{q}$ such that neither \mathfrak{p} nor \mathfrak{q} divides $y_1 = y(P)$, and such that the underlying primes $p, q \in \mathcal{P}_{\mathbb{Q}}$ are distinct and odd. Let $M = pq\#E(\mathbb{F}_{\mathfrak{p}})\#E(\mathbb{F}_{\mathfrak{q}})$.

We now define t sequences of primes $\{\ell_{i,r}\}, r = 1, \dots, t$ by using sets

$$V_{i,r}, i \in \mathbb{Z}_{>0}, r = 1, \dots, t$$

of previously defined elements of the sequences. More specifically we let $V_{1,1} = \emptyset$. For $i > 1$ we set

$$V_{i,1} = \{\ell_{1,1}, \dots, \ell_{1,t}, \dots, \ell_{i-1,1}, \dots, \ell_{i-1,t}\},$$

and for $i \geq 1, 1 < r \leq t$, we set

$$V_{i,r} = \{\ell_{1,1}, \dots, \ell_{1,t}, \dots, \ell_{i,1}, \dots, \ell_{i,r-1}\}.$$

Now let $\ell_{i,r}$ be the smallest prime outside \mathcal{L} and exceeding the bound implicit in Theorem 3.1 such that all of the following hold:

- (1) $\ell_{i,r} > \ell$ for all $\ell \in V_{i,r}$,
- (2) $\mu_{\ell_{i,r}} \leq 2^{-i}$,
- (3) $\mathbf{N}\mathfrak{p}_{\ell\ell_{i,r}}^{(r)} > 2^i$ for all $\ell \in V_{i,r} \cup \{\ell_{i,r}\} \cup \mathcal{L}$,
- (4) $\ell_{i,r} \equiv 1 \pmod{M}$,
- (5) the highest power of p dividing $(\ell_{i,r} - 1)/M$ is p^i , and
- (6) q divides $(\ell_{i,r} - 1)/M$ if and only if $i \in B$.

By Proposition 3.19 in [15] and by Proposition 6.1 we have:

Proposition 7.1. *The sequences $\{\ell_{1,r}, \ell_{2,r}, \dots\}$, $r = 1, \dots, t$, are well defined and computable.*

Lemma 7.1. *If $m \in \mathbb{Z}_{\geq 1}$, then*

$$\text{ord}_{\mathfrak{p}}(x_{mM+1} - x_1) = \text{ord}_{\mathfrak{p}}(x_{M+1} - x_1) + \text{ord}_p m.$$

Proof. This is Lemma 3.20 in [15]. □

We define the following subsets of \mathcal{P}_K :

- $\mathcal{T}_{1,r} = \bigcup_{i \geq 1} \mathcal{S}_{\ell_{i,r}}$;
- $\mathcal{T}_{2,r}^a$ is the set of $\mathfrak{p}_{\ell}^{(r)}$ for $\ell \notin (\{\ell_{1,r}, \ell_{2,r}, \dots\} \cup \mathcal{L})$, together with $\mathfrak{p}_{\ell_{a\ell}}^{(r)}$ for $\ell \in \mathcal{L}$;
- $\mathcal{T}_{2,r}^b = \{\mathfrak{p}_{\ell_{i,r}\ell_{j,r}}^{(r)} : 1 \leq j \leq i\}$;
- $\mathcal{T}_{2,r}^c = \{\mathfrak{p}_{\ell\ell_{i,r}}^{(r)} : \ell \in \mathcal{L}, i \geq 1\}$; and
- $\mathcal{T}_{2,r} = \mathcal{T}_{2,r}^a \cup \mathcal{T}_{2,r}^b \cup \mathcal{T}_{2,r}^c$.

As above we now have a version of Lemma 6.2.

Lemma 7.2.

- (1) For each $r = 1, \dots, t$, the sets $\mathcal{T}_{1,r}$ and $\mathcal{T}_{2,r}$ are disjoint. If a subset $S_r \subset \mathcal{P}_K$ contains $\mathcal{T}_{1,r}$ and is disjoint from $\mathcal{T}_{2,r}$, then $\mathcal{E}_r := \tilde{E}(\mathcal{O}_{K,S_r}) \cap zE(K)$ is the union of $\{\pm \ell_{i,r}P : i \geq 1\}$ and some subset of the finite set $\{sP : s \mid \prod_{\ell \in \mathcal{L}} \ell^{a_\ell - 1}\}$.
- (2) For any $j \in \{1, 2\}$ and $r, s \in \{1, \dots, t\}$ such that $r \neq s$ the sets $\mathcal{T}_{j,r}$ and $\mathcal{T}_{j,s}$ are disjoint.
- (3) For any $i \in \{1, 2\}$ and $r \in \{1, \dots, t\}$ the set $\mathcal{T}_{i,r}$ is computable.

We also have an analogous version of Proposition 5.1 and the proof is the same.

Proposition 7.2. *The natural density of $\mathcal{T}_{1,r}$ and $\mathcal{T}_{2,r}$ ($1 \leq r \leq t$) is zero.*

Now we can construct a Diophantine model of \mathbb{Z} in \mathcal{O}_{K,S_r} , where S_r is as in Lemma 7.2. We first need the following lemma.

Lemma 7.3. *Let $B = \{2^n + n^2 : n \in \mathbb{Z}_{\geq 1}\}$. Multiplication admits a positive existential definition in the structure $\mathcal{Z} := (\mathbb{Z}_{\geq 1}, 1, +, B)$. (Here B is considered as a unary predicate.) Hence the structure $(\mathbb{Z}, 0, 1, +, \cdot)$ admits a positive existential model in the structure \mathcal{Z} .*

Proof. This follows from Lemma 3.16 and Corollary 3.18 in [15]. \square

This lemma shows that instead of finding a Diophantine model of the ring \mathbb{Z} over $\mathcal{O}_{K,S}$, it will suffice to find a Diophantine model of \mathcal{Z} .

Proposition 7.3. *Let S_r be as in Lemma 7.2 and let $A_r := \{x_{\ell_{1,r}}, x_{\ell_{2,r}}, \dots\}$. Then A_r is a Diophantine model of \mathcal{Z} over \mathcal{O}_{K,S_r} via the bijection $\phi: \mathbb{Z}_{\geq 1} \rightarrow A$ taking i to $x_{\ell_{i,r}}$.*

Proof. The set A_r is Diophantine over \mathcal{O}_{K,S_r} by part (1) of Lemma 7.2.

We have

$$\begin{aligned} i \in B &\iff q \text{ divides } (\ell_{i,r} - 1)/M && \text{(by condition (4))} \\ &\iff \text{ord}_q(x_{\ell_{i,r}} - x_1) > \text{ord}_q(x_{M+1} - x_1) \end{aligned}$$

by Lemma 7.1 (with \mathfrak{q} in place of \mathfrak{p}). The latter inequality is a Diophantine condition on $x_{\ell_{i,r}}$. Thus the subset $\phi(B)$ of A_r is Diophantine over \mathcal{O}_{K,S_r} .

Finally, for $i \in \mathbb{Z}_{\geq 1}$, Lemma 7.1 and condition (3) imply $\text{ord}_{\mathfrak{p}}(x_{\ell_{i,r}} - x_1) = c + i$, where the integer $c = \text{ord}_{\mathfrak{p}}(x_{M+1} - x_1)$ is independent of i . Therefore, for $i, j, k \in \mathbb{Z}_{\geq 1}$, we have

$$i + j = k \iff \text{ord}_{\mathfrak{p}}(x_{\ell_{i,r}} - x_1) + \text{ord}_{\mathfrak{p}}(x_{\ell_{j,r}} - x_1) = \text{ord}_{\mathfrak{p}}(x_{\ell_{k,r}} - x_1) + c.$$

It follows that the graph of $+$ corresponds under ϕ to a subset of A_r^3 that is Diophantine over \mathcal{O}_{K,S_r} .

Thus A_r is a Diophantine model of \mathcal{Z} over \mathcal{O}_{K,S_r} . \square

8. Complementary rings

In this section, we complete the proofs of Theorems 1.1 and 1.2. First we need a general result about the existence of sets of primes of given densities. The result we will prove is contained in Proposition 8.2 below.

8.1. Sets of primes with prescribed densities. We start with describing the real numbers we consider as possible densities of our sets.

Definition 8.1 (Computable real numbers). A real number δ is called *computable* if there exists a computable sequence of rational numbers $r_n \in \mathbb{Q}$ such that

$$\lim_{n \rightarrow \infty} r_n = \delta.$$

It is easy to see that if $\alpha, \beta \neq 0$ are computable real numbers, then so is α/β . Next we observe that these are the only densities we should consider in the context of our problem.

Proposition 8.1. *Let K be a number field and let \mathcal{W}_K be a recursive set of primes of K having a natural density δ . In this case δ is a computable real number.*

Proof. Given our definition of a computable set \mathcal{A} of primes of a number field K (see Section 2), there exists a recursive procedure determining the size of the set $\{\mathfrak{p} \in \mathcal{A} : \mathbf{N}\mathfrak{p} \leq n\}$ uniformly in n , and therefore there exists a recursive function $g(n)$ computing $r_n = \frac{\#\{\mathfrak{p} \in \mathcal{A} : \mathbf{N}\mathfrak{p} \leq n\}}{\#\{\mathfrak{p} \in \mathcal{P}_K : \mathbf{N}\mathfrak{p} \leq n\}}$. If δ is the natural density of \mathcal{A} , then $\lim_{n \rightarrow \infty} r_n = \delta$ and therefore δ must be a computable real number. \square

We will now describe an (effective) procedure which, given a computable positive real number δ and a computable set of primes \mathcal{Z}_K of a number field K of natural density $\gamma \geq \delta$, constructs a computable set of primes $\mathcal{A}_K \subseteq \mathcal{Z}_K$ of natural density δ .

Without loss of generality we may assume that $\gamma > \delta$. Otherwise we set $\mathcal{A}_K = \mathcal{Z}_K$. Since \mathcal{Z}_K is computable, Proposition 8.1 implies that γ is computable. Let $\{d_i\}, \{g_i\}$ be computable sequences of rational numbers approximating δ and γ , respectively. Without loss of generality we can assume that $g_i \neq 0$ for all $i \in \mathbb{Z}_{>0}$. Then $\{r_i\} = \{d_i/g_i\}$ is a computable sequence of rational numbers approximating $\alpha := \delta/\gamma$. Without loss of generality we may also assume that all elements of the sequence $\{r_n\}$ are positive and strictly less than one.

Let $\{N_i\}$ be an increasing sequence of positive integers such that each $N_i = \mathbf{N}\mathfrak{p}_K$ for some prime \mathfrak{p}_K of K in \mathcal{Z}_K . Assume also that every positive integer $M \geq N_1$ occurring as a norm of a K -prime from \mathcal{Z}_K is an element of the sequence. Let $n = [K : \mathbb{Q}]$ and notice that n is the maximum number of K -primes that can have the same norm.

We now define recursively two sequences of sets of primes $\{\mathcal{A}_i\}$ and $\{\mathcal{Z}_i\}$. We denote by a_i and z_i the cardinality of $|\mathcal{A}_i|$ and $|\mathcal{Z}_i|$, respectively (for $i \geq 1$). Recall that $\{r_i\}$ is the sequence of rational numbers (strictly less than one) approximating $\alpha = \delta/\gamma$.

- (1) Set $\mathcal{A}_1 = \mathcal{Z}_1 = \{\mathfrak{p} \in \mathcal{Z}_K : \mathbf{N}\mathfrak{p} = N_1\}$.
- (2) If $\frac{a_i}{z_i} < r_i$, then set $\mathcal{A}_{i+1} = \mathcal{A}_i \cup \{\mathfrak{p} \in \mathcal{Z}_K : \mathbf{N}\mathfrak{p} = N_{i+1}\}$. Otherwise, set $\mathcal{A}_{i+1} = \mathcal{A}_i$.
- (3) For all $i \in \mathbb{Z}_{>0}$ set

$$\mathcal{Z}_{i+1} = \{\mathfrak{p} \in \mathcal{Z}_K : N_1 \leq \mathbf{N}\mathfrak{p} \leq N_{i+1}\}.$$

We have $\mathcal{Z}_K = \bigcup_{i \in \mathbb{Z}_{>0}} \mathcal{Z}_i$. We will now prove that the set $\mathcal{A}_K := \bigcup_{i \in \mathbb{Z}_{>0}} \mathcal{A}_i$ has density δ . To do this we need several lemmas. It is clear from the construction that \mathcal{A}_K is a computable set of primes.

Lemma 8.1.

- (1) If $\frac{a_i}{z_i} \geq r_i$ for some index i , then there is a positive integer k such that $\frac{a_{i+k}}{z_{i+k}} < r_i$.
- (2) If $\frac{a_i}{z_i} < r_i$ for some index i , then $\frac{a_{i+k}}{z_{i+k}} \geq r_i$ for some positive integer k .

Proof. The proofs for both assertions of the lemma will proceed by contradiction. Assume that the first assertion is false i.e., assume that there exists an $i \in \mathbb{Z}_{>0}$ such that $\frac{a_i}{z_i} \geq r_i$ and for all $k \in \mathbb{Z}_{>0}$ we have $\frac{a_{i+k}}{z_{i+k}} \geq r_{i+k}$. By step (2) in the above construction, the last inequality implies that $\mathcal{A}_{i+k} = \mathcal{A}_i$ and $a_{i+k} = a_i$ for all $k \in \mathbb{Z}_{>0}$. However, $\lim_{k \rightarrow \infty} z_{i+k} = \infty$ and therefore $\lim_{k \rightarrow \infty} \frac{a_{i+k}}{z_{i+k}} = 0$, while $\lim_{k \rightarrow \infty} r_{i+k} > 0$, which is a contradiction. Hence the first assertion of the lemma is true.

Assume now that the second assertion of the lemma is false i.e., that there exists an $i \in \mathbb{Z}_{>0}$ such that $\frac{a_i}{z_i} < r_i$ and such that

$$(8.1) \quad \forall k \in \mathbb{Z}_{>0} \quad \frac{a_{i+k}}{z_{i+k}} < r_{i+k}.$$

By step (2) of the above construction, (8.1) implies that

$$\forall k \in \mathbb{Z}_{>0} \quad \mathcal{A}_{i+k} = \mathcal{A}_{i+k-1} \cup \{\mathfrak{p} \in \mathcal{Z}_K : \mathbf{N}\mathfrak{p} = N_{i+1}\}.$$

This implies that $a_{i+k} \geq z_{i+k} - c$ for all $k \in \mathbb{Z}_{>0}$ for some fixed nonnegative integer c . At the same time, since the r_j 's are less than 1, (8.1) implies that $z_{i+k} > a_{i+k}$ for all $k \in \mathbb{Z}_{>0}$. Thus

$$\lim_{k \rightarrow \infty} \frac{a_{i+k}}{z_{i+k}} = \lim_{k \rightarrow \infty} \frac{z_{i+k}}{z_{i+k}} = 1 > \alpha = \lim_{k \rightarrow \infty} r_{i+k},$$

and therefore (8.1) cannot hold. \square

We now define two sequences of positive integers that we will use below.

Notation 8.1.

- Let $j_1 = 1$ and for $i \in \mathbb{Z}_{>1}$ define j_i to be the smallest positive integer greater than j_{i-1} such that $\frac{a_{j_i \pm 1}}{z_{j_i \pm 1}} < \frac{a_{j_i}}{z_{j_i}}$. (In other words, $\frac{a_{j_i}}{z_{j_i}}$ is a “local maximum”.)
- Let $k_0 = 0$ and for $i \in \mathbb{Z}_{>0}$ define k_i to be the smallest positive integer greater than k_{i-1} such that $\frac{a_{k_i \pm 1}}{z_{k_i \pm 1}} > \frac{a_{k_i}}{z_{k_i}}$. (In other words, $\frac{a_{k_i}}{z_{k_i}}$ is a “local minimum”.)

Remark 8.1. By construction of the sets $\mathcal{A}_i, \mathcal{Z}_i$ we have $z_i > a_i$ for all $i > 1$. Hence if $\mathcal{A}_i \subsetneq \mathcal{A}_{i+1}$, then $\frac{a_{i+1}}{z_{i+1}} = \frac{a_{i+m}}{z_{i+m}} > \frac{a_i}{z_i}$, where m is the number of primes of norm N_{i+1} in \mathcal{Z}_K . On the other hand, if $\mathcal{A}_i = \mathcal{A}_{i+1}$ then $\frac{a_{i+1}}{z_{i+1}} = \frac{a_i}{z_{i+m}} < \frac{a_i}{z_i}$. From Lemma 8.1 we can conclude that both j_i and k_i are defined for all $i \in \mathbb{Z}_{>0}$.

By construction of the sets \mathcal{A}_i we have $\frac{a_{j_i-1}}{z_{j_i-1}} < r_{j_i-1}$ and $\frac{a_{j_i}}{z_{j_i}} \geq r_{j_i}$ for all $i > 1$. Similarly, we have $\frac{a_{k_i-1}}{z_{k_i-1}} \geq r_{k_i-1}$ and $\frac{a_{k_i}}{z_{k_i}} < r_{k_i}$ for all $i > 0$.

We now show some properties of the sequences $\{k_i\}$ and $\{j_i\}$.

Lemma 8.2.

- (1) For all $i \in \mathbb{Z}_{\geq 0}$ we have $k_i < j_{i+1} < k_{i+1}$.

(2) For all $\ell \in \mathbb{Z}_{>0}$ there exists $i \in \mathbb{Z}_{>0}$ such that either

$$j_i \leq \ell \leq k_i \text{ and } \frac{a_{j_i}}{z_{j_i}} \geq \frac{a_\ell}{z_\ell} \geq \frac{a_{k_i}}{z_{k_i}}$$

or

$$k_i \leq \ell \leq j_{i+1} \text{ and } \frac{a_{k_i}}{z_{k_i}} \leq \frac{a_\ell}{z_\ell} \leq \frac{a_{j_{i+1}}}{z_{j_{i+1}}}.$$

Proof. By Lemma 8.1 and Remark 8.1 maxima and minima alternate in the sequence $\{a_i/r_i\}$. Further, by definition of $\mathcal{A}_1 = \mathcal{Z}_1$, it is clear that $j_1 = 1$ produces a local maximum in the sequence. \square

We now show that the local maxima and minima converge to $\alpha = \delta/\gamma = \lim_{i \rightarrow \infty} r_i$.

Lemma 8.3. $\lim_{i \rightarrow \infty} \frac{a_{j_i}}{z_{j_i}} = \alpha$ and $\lim_{i \rightarrow \infty} \frac{a_{k_i}}{z_{k_i}} = \alpha$.

Proof. We show that for any $\varepsilon > 0$ there exists a positive integer M such that for $i > M$ we have that $|\alpha - \frac{a_{j_i}}{z_{j_i}}| < \varepsilon$. The proof of the analogous statement with k_i substituted for j_i is similar. Fix $\mu < \varepsilon/4$. Let $I \in \mathbb{Z}_{>0}$ be large enough so that for all integers $s > I$ we have $|r_s - \alpha| < \mu < \varepsilon/4$ and $z_s > \frac{4n(1-\alpha)}{\varepsilon}$. (Recall that $n = [K : \mathbb{Q}]$.) Fix a positive integer $s > I$ and pick an i such that $j_i > s + 1$. By Remark 8.1

$$(8.2) \quad \frac{a_{j_i}}{z_{j_i}} = \frac{a_{j_{i-1}} + m}{z_{j_{i-1}} + m} \geq r_{j_i} > \alpha - \mu,$$

where as above, m is the number of K -primes in \mathcal{Z}_K with K -norm equal to N_{j_i} . By Remark 8.1 we also have $\frac{a_{j_{i-1}}}{z_{j_{i-1}}} < r_{j_{i-1}} < \alpha + \mu$. Thus

$$(8.3) \quad \frac{a_{j_i}}{z_{j_i}} - \alpha = \frac{a_{j_i}}{z_{j_i}} - \frac{a_{j_{i-1}}}{z_{j_{i-1}}} + \frac{a_{j_{i-1}}}{z_{j_{i-1}}} - \alpha =$$

$$(8.4) \quad \frac{a_{j_{i-1}}z_{j_{i-1}} + mz_{j_{i-1}} - a_{j_{i-1}}z_{j_{i-1}} - ma_{j_{i-1}}}{(z_{j_{i-1}} + m)z_{j_{i-1}}} + \frac{a_{j_{i-1}}}{z_{j_{i-1}}} - \alpha =$$

$$(8.5) \quad \frac{m(z_{j_{i-1}} - a_{j_{i-1}})}{(z_{j_{i-1}} + m)z_{j_{i-1}}} + \frac{a_{j_{i-1}}}{z_{j_{i-1}}} - \alpha =$$

$$(8.6) \quad \frac{m(1 - \frac{a_{j_{i-1}} + m}{z_{j_{i-1}} + m})}{z_{j_{i-1}}} + \frac{a_{j_{i-1}}}{z_{j_{i-1}}} - \alpha <$$

$$(8.7) \quad \frac{m(1 - \alpha + \mu)}{z_{j_{i-1}}} + \mu \leq \frac{n(1 - \alpha + \mu)}{z_{j_{i-1}}} + \mu < \varepsilon.$$

Combining (8.2) and (8.3)–(8.7) we conclude that

$$\left| \frac{a_{j_i}}{z_{j_i}} - \alpha \right| < \varepsilon.$$

Now we can prove that the natural density of \mathcal{A}_K is δ . \square

Corollary 8.1. *The natural density of \mathcal{A}_K is δ i.e.,*

$$\lim_{X \rightarrow \infty} \frac{\#\{\mathfrak{p}_K \in \mathcal{A}_K, N\mathfrak{p}_K \leq X\}}{\#\{\mathfrak{p}_K \in \mathcal{P}_K, N\mathfrak{p}_K \leq X\}} = \delta.$$

Proof. From Lemmas 8.2 and 8.3 it follows that $\lim_{i \rightarrow \infty} \frac{a_i}{z_i} = \alpha$. Now let X be a positive real number greater than N_1 . Then

$$N_i \leq X \leq N_{i+1}$$

for some positive integer i . (Recall that the N_i 's are norms of primes of K that appeared in the construction of the set \mathcal{A}_K .) Since a_i and z_i were defined to be the cardinality of \mathcal{A}_i and \mathcal{Z}_i , respectively, we have

$$\begin{aligned}\#\{\mathfrak{p}_K \in \mathcal{A}_K, \mathbf{N}\mathfrak{p}_K \leq X\} &= a_i, \\ \#\{\mathfrak{p}_K \in \mathcal{Z}_K, \mathbf{N}\mathfrak{p}_K \leq X\} &= z_i.\end{aligned}$$

This implies that $\lim_{X \rightarrow \infty} \frac{\#\{\mathfrak{p}_K \in \mathcal{A}_K, \mathbf{N}\mathfrak{p}_K \leq X\}}{\#\{\mathfrak{p}_K \in \mathcal{Z}_K, \mathbf{N}\mathfrak{p}_K \leq X\}} = \lim_{i \rightarrow \infty} \frac{a_i}{z_i} = \alpha$.

The statement of the corollary now follows from the fact that

$$\begin{aligned}& \lim_{X \rightarrow \infty} \frac{\#\{\mathfrak{p}_K \in \mathcal{A}_K, \mathbf{N}\mathfrak{p}_K \leq X\}}{\#\{\mathfrak{p}_K \in \mathcal{P}_K, \mathbf{N}\mathfrak{p}_K \leq X\}} \\ &= \lim_{X \rightarrow \infty} \left(\frac{\#\{\mathfrak{p}_K \in \mathcal{A}_K, \mathbf{N}\mathfrak{p}_K \leq X\}}{\#\{\mathfrak{p}_K \in \mathcal{P}_K, \mathbf{N}\mathfrak{p}_K \leq X\}} \cdot \frac{\#\{\mathfrak{p}_K \in \mathcal{Z}_K, \mathbf{N}\mathfrak{p}_K \leq X\}}{\#\{\mathfrak{p}_K \in \mathcal{Z}_K, \mathbf{N}\mathfrak{p}_K \leq X\}} \right) \\ &= \lim_{X \rightarrow \infty} \frac{\#\{\mathfrak{p}_K \in \mathcal{A}_K, \mathbf{N}\mathfrak{p}_K \leq X\}}{\#\{\mathfrak{p}_K \in \mathcal{Z}_K, \mathbf{N}\mathfrak{p}_K \leq X\}} \cdot \lim_{X \rightarrow \infty} \frac{\#\{\mathfrak{p}_K \in \mathcal{Z}_K, \mathbf{N}\mathfrak{p}_K \leq X\}}{\#\{\mathfrak{p}_K \in \mathcal{P}_K, \mathbf{N}\mathfrak{p}_K \leq X\}} = \alpha\gamma = \delta\end{aligned}$$

□

Finally we have the following proposition.

Proposition 8.2. *If $\delta_1, \dots, \delta_t$ is a finite set of nonnegative computable real numbers adding up to one, then there exist a partition of \mathcal{P}_K into computable sets W_1, \dots, W_t of densities $\delta_1, \dots, \delta_t$, respectively.*

Proof. Without loss of generality we can assume that all the densities are positive. We proceed in $t - 1$ steps. First set $\mathcal{Z}_K = \mathcal{P}_K$ and construct a computable set W_1 of density δ_1 . Observe that $\mathcal{P}_K - W_1$ is computable of density $1 - \delta_1 \geq \delta_2$. Now set $\mathcal{Z}_K = \mathcal{P}_K - W_1$ and construct W_2, W_3 , etc. □

Remark 8.2. The construction above shows in fact that there exists a partition of \mathcal{P}_K into sets of any densities adding to 1. However, if we do not require that the densities are computable, the resulting sets of primes may be uncomputable.

Now we can prove Theorems 1.1 and 1.2.

8.2. The proofs of Theorems 1.1 and 1.2. Let $\delta_1, \dots, \delta_t$ be nonnegative computable real numbers adding up to one. Let W_1, \dots, W_t be a partition of primes of K , where the natural density of each W_i is δ_i . Such a partition exists by Proposition 8.2. For the case of Theorem 1.1, let $\mathcal{T}_{1,r}, \mathcal{T}_{2,r}, r = 1, \dots, t$ be as defined in Section 6 and for the case of Theorem 1.2, let $\mathcal{T}_{1,r}, \mathcal{T}_{2,r}, r = 1, \dots, t$ be as defined as in Section 7. For $i = 1, \dots, t$ define

$$S_i = (W_i \cup \mathcal{T}_{1,i} \cup \mathcal{T}_{2,j}) - (\mathcal{T}_{2,i} \cup \bigcup_{r \neq i} \mathcal{T}_{1,r}),$$

where $j \in \{1, \dots, t\}$ is such that $j \equiv i - 1 \pmod{t}$. We claim the following:

- (1) *The natural density of S_i exists and is equal to δ_i .* This is true because by Propositions 5.1 and 6.2, for any i, j the natural density of $\mathcal{T}_{i,j}$ is 0.
- (2) *Each S_i contains all the primes of $\mathcal{T}_{1,i}$ and omits the primes of $\mathcal{T}_{2,i}$.* To see that this assertion is true, observe that we explicitly add $\mathcal{T}_{1,i}$ and remove $\mathcal{T}_{2,i}$, and by Propositions 6.2 and 7.2, we have that $\mathcal{T}_{2,i} \cap \mathcal{T}_{2,j} = \emptyset$ for $i \neq j$. Thus, adding $\mathcal{T}_{2,j}$ does not introduce any primes of $\mathcal{T}_{2,i}$ back. Further from the same propositions removing $\bigcup_{r \neq i} \mathcal{T}_{1,r}$ will not remove any primes of $\mathcal{T}_{1,i}$.
- (3) *S_1, \dots, S_t are a partition of \mathcal{P}_K .* First we show that $S_i \cap S_r = \emptyset$ for $i \neq r$. Since W_i and W_r are disjoint, the common elements can arise only from the primes which were added in, i.e., an intersection can arise from

$$(8.8) \quad (\mathcal{T}_{1,i} \cup \mathcal{T}_{2,j}) \cap (\mathcal{T}_{1,r} \cup \mathcal{T}_{2,l}),$$

where $j \equiv i - 1 \pmod{t}$, and $l \equiv r - 1 \pmod{t}$ so that $l \neq j$. By construction, all the primes of $\mathcal{T}_{1,r}$ are removed from S_i and all the primes of $\mathcal{T}_{1,i}$ are removed from S_r . Hence the only primes from (8.8) which can possibly be in $S_i \cap S_r$ are in $\mathcal{T}_{2,j} \cap \mathcal{T}_{2,l}$. This intersection is empty, however, by Propositions 6.2 and 7.2. Finally we show that $\bigcup_{i=1}^t S_i = \mathcal{P}_K$. As above we start with the fact that $\bigcup_{i=1}^t W_i = \mathcal{P}_K$ and note that we only have to follow the primes removed from W_i in the process of constructing S_i :

$$\mathcal{T}_{2,i} \cup \bigcup_{r \neq i} \mathcal{T}_{1,r}.$$

We have shown in Part 1 of this proposition that for $r = 1, \dots, t$, $\mathcal{T}_{1,r} \subset S_r$ and therefore the primes in the union $\bigcup_{r \neq i} \mathcal{T}_{1,r}$ are accounted for. That leaves the primes of

$$\mathcal{T}_{2,i} - \bigcup_{r \neq i} \mathcal{T}_{1,r} = \mathcal{T}_{2,i} - \bigcup_{r=1}^t \mathcal{T}_{1,r},$$

where the equality holds because $\mathcal{T}_{1,i} \cap \mathcal{T}_{2,i} = \emptyset$. When S_i is constructed, this set is moved to S_j , $j \equiv i + 1 \pmod{t}$ and observe that since $\mathcal{T}_{2,i} \cap \mathcal{T}_{2,j} = \emptyset$, the primes of $\mathcal{T}_{2,i} - \bigcup_{r=1}^t \mathcal{T}_{1,r}$ are not removed from S_j .

Acknowledgments

Our thanks go to the London Mathematical Society for the very enjoyable visit by the third author to the University of East Anglia. Also, to whichever barman invented the Marguerita. We also thank Bjorn Poonen who suggested the use of sets of nonnegative computable real densities rather than rational densities.

K.E. was partially supported by National Science Foundation grants DMS-0801123, DMS-1056703 and a Sloan Research Fellowship. A.S. was partially supported by National Science Foundation grant DMS-0650927 and grant 13419 from the John Templeton Foundation.

References

- [1] Yu. Bilu, G. Hanrot and P. M. Voutier, *Existence of primitive divisors of Lucas and Lehmer numbers*, J. Reine Angew. Math. **539** (2001), 75–122.
- [2] W. Calvert, V. Harizanov and A. Shlapentokh, *Turing degrees of isomorphism types of algebraic objects*, J. London Math. Soc. (2) **75**(2) (2007), 273–286.
- [3] J. Cheon and S. Hahn, *The orders of the reductions of a point in the Mordell–Weil group of an elliptic curve*, Acta. Arith. **88**(3) (1999), 219–222.
- [4] G. Cornelissen and K. Zahidi, *Topology of diophantine sets: remarks on Mazur's conjectures*, in 'The Hilbert's tenth problem: relations with arithmetic and algebraic geometry' (Ghent, 1999), Amer. Math. Soc., Providence, RI, 2000, 253–260.
- [5] M. Davis, H. Putnam and J. Robinson, *The decision problem for exponential diophantine equations*, Ann. of Math. (2), **74** (1961), 425–436.
- [6] K. Eisenträger and G. Everest, *Descent on elliptic curves and Hilbert's tenth problem*, Proc. Amer. Math. Soc. **137**(6) (2009), 1951–1959.
- [7] G. Everest, A. J. van der Poorten, I. E. Shparlinski and T. Ward, *Recurrence sequences*, Math. Surveys and Monographs, **104**, Amer. Math. Soc., Providence, RI, 2003.
- [8] Yu. V. Matijasevič, *The diophantineness of enumerable sets*, Dokl. Akad. Nauk SSSR, **191** (1970), 279–282.
- [9] B. Mazur, *The topology of rational points*, Experiment. Math. **1**(1) (1992), 35–45.
- [10] B. Mazur, *Speculations about the topology of rational points: an update*, Astérisque (1995), no. 228, pp. 4, 165–182, Columbia University Number Theory Seminar (New York, 1992).
- [11] B. Mazur, *Open problems regarding rational points on curves and varieties*, in 'Galois representations in arithmetic algebraic geometry' (Durham, 1996), London Math. Soc. Lecture Note Ser. **254**, Cambridge Univ. Press, Cambridge, 1998, 239–265.
- [12] B. Mazur and K. Rubin, *Ranks of twists of elliptic curves and Hilbert's tenth problem*, Invent. Math., **181**(3) (2010), 541–575.
- [13] S. Perlega, *Additional results to a theorem of Eisenträger and Everest*, Arch. der Mathematik, **97**(2) (2011), 141–149.
- [14] B. Poonen, *Hilbert's tenth problem and Mazur's conjecture for large subrings of \mathbb{Q}* , J. Amer. Math. Soc. **16**(4) (2003), 981–990.
- [15] B. Poonen and A. Shlapentokh, *Diophantine definability of infinite discrete nonarchimedean sets and diophantine models over large subrings of number fields*, J. Reine Angew. Math. **588** (2005), 27–47.
- [16] M. Pohst and H. Zassenhaus, *Algorithmic algebraic number theory is encyclopedia of mathematics and its applications*, **30**, Cambridge University Press, 1997.
- [17] A. Schinzel, *On primitive prime factors of $a^n - b^n$* , Proc. Cambridge Phils. Soc. **58** (1962), 555–562.
- [18] A. Schinzel, *Primitive divisors of the expression $A^n - B^n$ in algebraic number fields*, J. Reine Angew. Math. **268/69** (1974), 27–33.
- [19] A. Schinzel, *An extension of the theorem on primitive divisors in algebraic number fields*, Math. Comp. **61**(203) (1993), 441–444.
- [20] A. Shlapentokh, *Diophantine definability over some rings of algebraic numbers with infinite number of primes allowed in the denominator*, Invent. Math. **129**(3) (1997), 489–507.
- [21] A. Shlapentokh, *Defining integrality at prime sets of high density in number fields*, Duke Math. J. **101**(1) (2000), 117–134.
- [22] A. Shlapentokh, *Diophantine definability and decidability in large subrings of totally real number fields and their totally complex extensions of degree 2*, J. Number Theory **95**(2) (2002), 227–252.
- [23] A. Shlapentokh, *A ring version of Mazur's conjecture on topology of rational points*, Int. Math. Res. Not. (7) (2003), 411–423.
- [24] A. Shlapentokh, *On diophantine definability and decidability in some infinite totally real extensions of \mathbb{Q}* , Trans. Amer. Math. Soc. **356**(8) (2004), 3189–3207.
- [25] A. Shlapentokh, *Hilbert's tenth problem: diophantine classes and extensions to global fields*, Cambridge University Press, 2006.
- [26] A. Shlapentokh, *Diophantine definability and decidability in the extensions of degree 2 of totally real fields*, J. Algebra **313**(2) (2007), 846–896.

- [27] J. H. Silverman, *The arithmetic of elliptic curves*, Springer, New York, 1986.
- [28] J. H. Silverman, *Weierich's criterion and the ABC-conjecture*, J. Number Theory, **30** (1988), 226–237.
- [29] J. H. Silverman, *Advanced topics in the arithmetic of elliptic curves*, Springer, New York, 1994.
- [30] K. Zsigmondy, *Zur theorie der potenzreste*, Monatsh. Math. **3** (1892), 265–284.

DEPARTMENT OF MATHEMATICS, THE PENNSYLVANIA STATE UNIVERSITY, UNIVERSITY PARK,
PA 16802, USA

E-mail address: `eisentra@math.psu.edu`

SCHOOL OF MATHEMATICS, UNIVERSITY OF EAST ANGLIA, NORWICH NR4 7TJ, UK

E-mail address: `g.everest@uea.ac.uk`

DEPARTMENT OF MATHEMATICS, EAST CAROLINA UNIVERSITY, GREENVILLE, NC 27858, USA

E-mail address: `shlapentokha@ecu.edu`