

The A-polynomial and holonomy perturbations

JIANFENG LIN

Dunfield-Garoufalidis and Boyer-Zhang proved that the A-polynomial of a non-trivial knot in S^3 is non-trivial. In this paper, we use holonomy perturbations to prove the non-triviality of the A-polynomial for a non-trivial, null-homotopic knot in an irreducible 3-manifold. Also, we give a strong constraint on the A-polynomial of a knot in the 3-sphere.

1. Introduction

In [4], an algebraic curve D_N was associated to a compact 3-manifold N with a single torus boundary. We briefly review the construction: consider the variety $\chi(\pi_1(N))$ of characters of $SL(2, \mathbb{C})$ representations of $\pi_1(N)$. We can restrict a representation to the boundary group $\pi_1(\partial N)$, which gives us a map $r: \chi(\pi_1(N)) \rightarrow \chi(\pi_1(T^2))$. Choose a basis $B = \{\mathfrak{M}, \mathfrak{L}\}$ for the boundary group. Given two nonzero complex numbers m, l , we can define a $SL(2, \mathbb{C})$ representation $\gamma_{(m,l)}$ of $\pi_1(T^2)$:

$$(1) \quad \gamma_{(m,l)}(\mathfrak{M}) = \begin{pmatrix} m & 0 \\ 0 & m^{-1} \end{pmatrix}, \quad \gamma_{(m,l)}(\mathfrak{L}) = \begin{pmatrix} l & 0 \\ 0 & l^{-1} \end{pmatrix}.$$

So we have a natural map $h: \mathbb{C}^* \times \mathbb{C}^* \rightarrow \chi(\pi_1(T^2))$ defined by $h(m, l) := [\gamma_{(m,l)}] \in \chi(\pi_1(T^2))$.

Now we consider $h^{-1}(\overline{\text{Im}(r)})$. The following lemma is proved in [5].

Lemma 1.1. *This algebraic set $h^{-1}(\overline{\text{Im}(r)}) \subset \mathbb{C}^* \times \mathbb{C}^*$ has no 2-dimensional irreducible components.*

The algebraic curve D_N is defined to be the union of the 1-dimensional components of $h^{-1}(\overline{\text{Im}(r)})$. Each irreducible component of D_N is defined by a two variable irreducible polynomial. We multiply all these polynomials to get the A-polynomial of N , which we denote by $A_{N,B}$. (It depends on the choice of the basis B .)

If K is a null-homologous knot in a three manifold Y , we can take N to be the knot complement. There is a natural basis \tilde{B} for the boundary group: the meridian and the longitude (see Fact 2.8). The A-polynomial A_K of the knot K is defined to be $A_{N, \tilde{B}}$.

In [5] and [2], Dunfield-Garoufalidis and Boyer-Zhang independently proved that:

Theorem 1.2 (Dunfield-Garoufalidis, Boyer-Zhang [5],[2]). *For a non-trivial knot K in S^3 , we have $A_K \neq l - 1$.*

Remark 1.3. The reducible representations of the knot group always give a factor $l - 1$ of the A-polynomial. The A-polynomial of a trivial knot is exactly $l - 1$. Therefore, we say a knot K has non-trivial A-polynomial if $A_K \neq l - 1$.

Both Dunfield-Garoufalidis's and Boyer-Zhang's proofs make use of the following theorem.

Theorem 1.4 (Kronheimer-Mrowka [11]). *For a non-trivial knot K in S^3 , let Y_r ($r \in \mathbb{Q}$) be the manifold obtained by doing r -surgery along K . If $|r| \leq 2$, then $\pi_1(Y_r)$ admits some $SU(2)$ representation with non-cyclic image.*

This theorem is proved using techniques from $SU(2)$ gauge theory, namely holonomy perturbations. Holonomy perturbations were used by Floer in [3] to prove the surgery exact triangle. In fact, using holonomy perturbations, we can prove Theorem 1.2 directly. Moreover, we can prove something stronger.

The following are the main theorems of this paper.

Theorem 1.5. *Suppose K is a null-homologous, non-trivial knot in an orientable, closed, irreducible 3-manifold Y . If $A_K = l - 1$, then there exists a homomorphism $\rho : \pi_1(Y) \rightarrow SU(2)$ such that $\rho([K]) = -1 \in SU(2)$.*

Theorem 1.6. *For a null-homotopic, non-trivial knot K in any orientable, closed, irreducible 3-manifold Y , we have $A_K \neq l - 1$.*

In light of these two theorems, we raise the following question.

Question 1.7. Does any null-homologous knot in an irreducible 3-manifold have non-trivial A-polynomial?

If we restrict to knots in S^3 , Kronheimer and Mrowka's argument in [11] implies that the zero set of A-polynomial intersects the unit torus in some arcs. We give the precise statement:

Theorem 1.8. *Let K be a non-trivial knot in S^3 . Then given any complex number l with $|l| = 1$, we can find another unit length complex number m such that $A_K(l, m) = 0$.*

Remark 1.9. Theorem 1.8 gives strong constraints on the A-polynomial: for a generic polynomial, its zero set intersects the unit torus at isolated points. This is ruled out by Theorem 1.8.

In Section 2, we review some preliminaries and basic constructions related to holonomy perturbations and the non-vanishing theorem of critical points. In Section 3, we prove the main theorems and give some examples.

2. Preliminaries

In this section, we review some of the constructions related to holonomy perturbations. Most of the details can be found in [11] and [3].

2.1. Holonomy perturbation

Let M be a closed, oriented 3-manifold with $b_1(M) > 0$. Consider a rank 2 unitary bundle E over M with non-torsion $c_1(E)$. Let \mathfrak{g}_E be the bundle whose sections are traceless, skew-hermitian endomorphisms of E . Let \mathcal{A} be the affine space of $SO(3)$ connections of \mathfrak{g}_E . Let \mathcal{G} be the group of gauge transformations on E with determinant 1.

Fix a reference connection A_0 on \mathfrak{g}_E . For any connection A on \mathfrak{g}_E , denoting $A - A_0$ by ω , we have the Chern-Simons functional:

$$CS : \mathcal{A} \rightarrow \mathbb{R}$$

$$CS(A) = \frac{1}{4} \int_{X_0} \text{Tr} \left(2\omega \wedge F_{A_0} + \omega \wedge d\omega + \frac{2}{3} \omega \wedge \omega \wedge \omega \right)$$

The critical points of the Chern-Simons functional are the flat connections.

Floer defined a class of perturbations of the Chern-Simons functional as follows. We take a function $\phi : SU(2) \rightarrow \mathbb{R}$ which is invariant under conjugation. It is uniquely determined by the even, 2π -periodic function $f(x) :=$

$\phi \begin{pmatrix} e^{ix} & 0 \\ 0 & e^{-ix} \end{pmatrix}$. Let D be a compact 2-manifold with boundary. Consider an embedding of $D \times S^1$ in M such that \mathfrak{g}_E is trivial over it. Fix a trivialization of \mathfrak{g}_E over $D \times S^1$ and take a 2-form μ which is supported in the interior of D and has integral 1. Using the trivialization, we can lift A to a connection \bar{A} on the trivialized $SU(2)$ bundle \tilde{P} over $D \times S^1$. Consider:

$$\begin{aligned} \Phi : \mathcal{A} &\rightarrow \mathbb{R} \\ (2) \quad \Phi(A) &= \int_{p \in D} \phi(\text{Hol}_{\{p\} \times S^1}(\bar{A})) \mu(p) \end{aligned}$$

In this paper, we specialize D to be a disk D^2 or an annulus H . In the case of the disk, let $z_0 \in \partial D^2$ be a base point. We define the curves α and β to be $\partial D^2 \times \{0\}$ and $\{z_0\} \times S^1$ respectively. In the case of the annulus, we denote the two components of ∂H by c_1, c_2 and choose base points $z_i \in c_i$. Then for $i = 1, 2$, we define $\alpha_i, \beta_i \subset \partial(H \times S^1)$ to be $c_i \times \{0\}$ and $\{z_i\} \times S^1$ respectively.

Now we consider the perturbed Chern-Simons functional: $CS + \Phi : \mathcal{A} \rightarrow \mathbb{R}$.

The critical points can be completely described by the following lemmas. The first was proved by Floer. The second follows from similar arguments. For completeness, we sketch the proof for the second lemma. See [3] for details.

Lemma 2.1 (Floer [3]). *If $D \cong D^2$ and $A \in \mathcal{A}$ is a critical point of $CS + \Phi$, then:*

- 1) A is flat on $M - (D^2 \times S^1)$
- 2) After choosing a new trivialization of \tilde{P} , $\text{Hol}_\alpha(\bar{A}) = \begin{pmatrix} e^{i\tau} & 0 \\ 0 & e^{-i\tau} \end{pmatrix}$, $\text{Hol}_\beta(\bar{A}) = \begin{pmatrix} e^{i\nu} & 0 \\ 0 & e^{-i\nu} \end{pmatrix}$ and $\tau = f'(\nu) + 2\pi\mathbb{Z}$

Lemma 2.2. *If $D \cong H$ and $A \in \mathcal{A}$ is a critical point of $CS + \Phi$, then:*

- 1) A is flat on $M - (H \times S^1)$
- 2) The $SU(2)$ connection \bar{A} is reducible and we can choose a new trivialization of \tilde{P} such that:

$$\begin{aligned} \text{Hol}_{\alpha_1}(\bar{A}) &= \begin{pmatrix} e^{i\tau_1} & 0 \\ 0 & e^{-i\tau_1} \end{pmatrix}, & \text{Hol}_{\alpha_2}(\bar{A}) &= \begin{pmatrix} e^{i\tau_2} & 0 \\ 0 & e^{-i\tau_2} \end{pmatrix} \\ \text{Hol}_{\beta_1}(\bar{A}) &= \text{Hol}_{\beta_2}(\bar{A}) = \begin{pmatrix} e^{i\nu} & 0 \\ 0 & e^{-i\nu} \end{pmatrix} \end{aligned}$$

Moreover, we have:

$$(3) \quad \tau_1 - \tau_2 = f'(\nu) + 2\pi\mathbb{Z}.$$

Proof. Let P be the trivialized $SU(2)$ bundle over $H \times S^1$ and \mathfrak{g}_E be the vector bundle associated to the adjoint representation. For $p \in H$, denote $\{p\} \times S^1$ by β_p . For a point $a = (p, q) \in H \times S^1$, we consider the holonomy of \bar{A} along β_p . Let us denote it by $T_{\bar{A}}(a)$. This is an automorphism of the fiber of P over a . Thus $T_{\bar{A}}$ defines a section of $\text{Aut}(P)$. Since ϕ is conjugation invariant, it defines a function on $\text{Aut}(P)$. Let $\phi'(T_{\bar{A}})$ denotes the gradient of this function, evaluated at the section $T_{\bar{A}}$. One can check that $\phi'(T_{\bar{A}})$ defines a section of \mathfrak{g}_E . Now A is a critical point of $CS + \Phi$. It is proved in [3] that:

$$(4) \quad F_{\bar{A}} = \phi'(T_{\bar{A}})\mu.$$

Using this, we can deduce that the holonomy along β_p does not depend on p and the covariant derivative $\nabla_{\bar{A}}(T_{\bar{A}}) = 0$ (see [3]). If $T_{\bar{A}}(a) = \pm 1$ for any $a \in H \times S^1$, then $\phi'(T_{\bar{A}}) \equiv 0$. By Equation (4), \bar{A} is flat. Since $\pi_1(H \times S^1)$ is abelian, we see that \bar{A} is reducible. Suppose $T_{\bar{A}}(a) \neq \pm 1$ for some $a \in H \times S^1$. Since $\nabla_{\bar{A}}(T_{\bar{A}}) = 0$, we have $T_{\bar{A}}(a) \neq \pm 1$ for any $a \in H \times S^1$. Thus the existence of this section tells us that \bar{A} is also reducible in this case.

Because \bar{A} is reducible, we can choose a suitable trivialization of \tilde{P} such that $\bar{A} = \sigma \cdot \begin{pmatrix} i & 0 \\ 0 & -i \end{pmatrix}$ for some one form $\sigma \in \Omega^1(H \times S^1)$. Since the holonomy along β_p does not depend on p , we can assume that $\text{Hol}_{\beta_p}(\bar{A}) = \begin{pmatrix} e^{i\nu} & 0 \\ 0 & e^{-i\nu} \end{pmatrix}$ for any p . In particular, we have $\text{Hol}_{\beta_i}(\bar{A}) = \begin{pmatrix} e^{i\nu} & 0 \\ 0 & e^{-i\nu} \end{pmatrix}$ for $i = 1, 2$. Then Formula (4) becomes:

$$(5) \quad d\sigma = f'(\nu)\mu.$$

By Stokes' theorem, we have $\int_{\alpha_1} \sigma - \int_{\alpha_2} \sigma = \int_{H \times \{0\}} d\sigma = f'(\nu)$. It is easy to see that $\tau_i = \int_{\alpha_i} \sigma$. Therefore, we proved the lemma. \square

The following theorem was first proved in [11] and [12].

Theorem 2.3 (Kronheimer-Mrowka [11],[12]). *Let M, E, \mathfrak{g}_E be defined as before. If M admits an oriented, smooth taut foliation and is not $S^2 \times S^1$, then for any holonomy perturbation Φ , the perturbed Chern-Simons functional $CS + \Phi$ over E has at least one critical point.*

This highly non-trivial theorem is a corollary of several theorems. We only sketch the proof here. For details, see [11] and [12]. If M admits

a smooth, orientable taut foliation, then by the work by Eliashberg and Thurston in [6] and [7], M can be embedded in an admissible symplectic 4-manifold X . [11] proved that X can be chosen to satisfy several good conditions which imply, by Feehan and Leness's work (see [12] and [8]), that it satisfies Witten's conjecture relating Seiberg-Witten invariants and Donaldson invariants. Because X is symplectic, its Seiberg-Witten invariants are non-trivial by Taubes's non-vanishing Theorem in [13]. From Witten's conjecture, it follows that the Donaldson invariants D_X^v are non-trivial for any line bundle v on X . Then by a standard stretching argument, the perturbed Chern-Simons functional has at least one critical point.

The following two theorems are proved by Gabai.

Theorem 2.4 (Gabai [9]). *Let M be a closed, orientable, irreducible 3-manifold. If $b_1(M) > 0$, then M admits an orientable continuous taut foliation. Moreover, the taut foliation can be chosen to be smooth if $H_2(M, \mathbb{Z})$ is not generated by embedded tori or spheres.*

Theorem 2.5 (Gabai [9]). *Suppose $K \subset S^3$ is a non-trivial knot and Y_0 is obtained by doing 0-surgery along K . Then Y_0 admits an orientable taut foliation and is not $S^2 \times S^1$.*

Remark 2.6. When the genus of $K \subset S^3$ is 1, Gabai's taut foliation on Y_0 may not be smooth. But we can still embed M in an admissible symplectic 4-manifold. See the proof of Theorem 6.1 in [10].

Combining these theorems, we get:

Corollary 2.7. *Suppose M satisfies any one of the following two conditions:*

- *M is irreducible with $b_1(M) > 0$ and $H_2(M, \mathbb{Z})$ is not generated by embedded tori or spheres;*
- *M is the 0-surgery manifold for a non-trivial knot in S^3 .*

Then for any holonomy perturbation on M , the perturbed Chern-Simons functional has at least one critical point.

2.2. The $SU(2)$ -Pillowcase

Now suppose $K \subset Y$ is a null-homologous knot in an orientable closed 3-manifold Y . Let $N(K)$ denote an open tubular neighborhood of K . Unless

otherwise stated, all homology groups are considered with \mathbb{Z} coefficients. Using the Mayer-Vietoris sequence, it is easy to prove:

Fact 2.8. *The meridian $m \subset \partial(Y - N(K))$ is a primitive, nontorsion element in $H_1(Y - N(K))$. Also, we can find a unique longitude l such that $[l] = 0 \in H_1(Y - N(K))$ and $([m], [l])$ forms a basis for $H_1(\partial(Y - N(K)))$.*

Let $Y_r (r \in \mathbb{Q})$ be the manifold obtained by doing r -surgery along K . It is easy to see that $H_1(Y_0) = H_1(Y - N(K))$.

Suppose that the attached solid torus is $N_0 \subset Y_0$ with core c . Then $Y_0 - N_0 \cong Y - N(K)$. Now we consider a trivialized rank 2 unitary bundle E over $Y - N(K)$ (using basic obstruction theory, we can prove that any $SU(2)$ -bundle over $Y - N(K)$ is trivial since $\pi_1(SU(2)) = \pi_2(SU(2)) = 0$). Let \mathfrak{g}_E be defined as before. Let \mathcal{G} be the group of gauge transformations with determinant 1 on E . Consider a flat connection A on \mathfrak{g}_E . Using the trivialization, we can lift A to a connection on the trivialized $SU(2)$ bundle \tilde{P} over $Y - N(K)$. Denote this connection by \tilde{A} .

We have the following standard lemma:

Lemma 2.9. *By taking the holonomy of \tilde{A} , we get a 1-1 correspondence between the flat connections on \mathfrak{g}_E modulo the action of \mathcal{G} and the homomorphisms $\rho_A : \pi_1(Y - N(K)) \rightarrow SU(2)$ modulo conjugation.*

Because m and l are commutative as elements in $\pi_1(Y - N(K))$, we can gauge transform \tilde{A} so that $\text{Hol}_m(\tilde{A})$ and $\text{Hol}_l(\tilde{A})$ are both diagonal.

Thus we can assume that

$$(6) \quad \text{Hol}_m(\tilde{A}) = \begin{pmatrix} e^{i\theta_A} & 0 \\ 0 & e^{-i\theta_A} \end{pmatrix}, \quad \text{Hol}_l(\tilde{A}) = \begin{pmatrix} e^{i\eta_A} & 0 \\ 0 & e^{-i\eta_A} \end{pmatrix}.$$

Definition 2.10. We define the subset of $(\mathbb{R}/2\pi\mathbb{Z})^2$ as:

$$R_K := \{\pm(\theta_A, \eta_A) \mid A \text{ is a flat connection on } \mathfrak{g}_E|_{Y_0 - N_0}\}.$$

Remark 2.11. Points in the torus $(\mathbb{R}/2\pi\mathbb{Z})^2$ determine diagonal representations of $\pi_1(T^2)$ to $SU(2)$, although there is still a remaining conjugation action $(a, b) \mapsto (-a, -b)$. Therefore, $(\mathbb{R}/2\pi\mathbb{Z})^2$ is actually the double branch cover of the $SU(2)$ -pillowcase, which refers to the space $\text{Hom}(\pi_1(T^2), SU(2))/\text{conjugation}$. In this paper, we work with $\mathbb{Z}/2\mathbb{Z}$ equivariant arguments in the torus.

After checking the construction carefully and using Lemma 2.9, it's easy to see that R_K can be also defined as:

$$(7) \quad \left\{ (\theta, \eta) \mid \exists \rho : \pi_1(Y - K) \rightarrow SU(2) \right. \\ \left. \text{s.t. } \rho(m) = \begin{pmatrix} e^{i\theta} & 0 \\ 0 & e^{-i\theta} \end{pmatrix}; \rho(l) = \begin{pmatrix} e^{i\eta} & 0 \\ 0 & e^{-i\eta} \end{pmatrix} \right\}$$

Remark 2.12. Because $\pi_1(Y - K)$ is finitely generated and $SU(2)$ is compact, the $SU(2)$ representation space of $\pi_1(Y - K)$ is compact. Thus R_K is a closed subset of $(\mathbb{R}/2\pi\mathbb{Z})^2$.

Remark 2.13. If ρ is a reducible $SU(2)$ representation of $\pi_1(Y - K)$, then $\rho(l) = 1$. Thus the points in R_K off the line $\eta = 2k\pi$ give irreducible representations.

The case $Y \cong S^3$ of the following lemma is proved as Lemma 13 in [11]. Although the general proof is essentially the same, we give it here for completeness.

Lemma 2.14. *R_K is invariant under the translation by $(\pi, 0)$.*

Proof. By Fact 2.8, there exists $H_1(Y - N(K)) \rightarrow \mathbb{Z}_2$ mapping m to -1 and mapping l to 1 . Considering the composition of $\pi_1(Y - N(K)) \rightarrow H_1(Y - N(K)) \rightarrow \mathbb{Z}_2 \rightarrow SU(2)$, we get $\rho_0 : \pi_1(Y - N(K)) \rightarrow SU(2)$ with the image in the center. For $\rho : \pi_1(Y - N(K)) \rightarrow SU(2)$, we multiply it with ρ_0 to get another representation ρ' . We have $\rho(m) = -\rho'(m)$ and $\rho(l) = \rho'(l)$. Then we use Formula (7) to prove the lemma. \square

3. Proof of main theorem

3.1. The general case

Assume $K \subset Y$ is a null-homologous, non-trivial knot in an irreducible 3-manifold. If K is contained in a 3-ball in Y , then K has non-trivial Alexander polynomial by Theorem 1.2. Therefore, we just consider the case when K is not contained in a 3-ball. This implies that $Y - N(K)$ is irreducible and its boundary is incompressible. We take two copies of $Y - N(K)$ and denote them by $Y_j - N(K_j)$ ($j = 1, 2$). We glue them onto a thickened torus $T \times [-1, 1]$ using the same gluing map to get a irreducible closed manifold M . M

is just the double of the knot complement. It's easy to see that $b_1(M) > 0$ and the meridian $m \subset T \times \{0\}$ is a non-torsion element in $H_1(M)$.

Remark 3.1. We identified the torus T with $\partial(Y - N(K))$. Thus it makes sense to talk about the meridian $m \subset T$ and the longitude $l \subset T$ by the Fact 2.8.

Lemma 3.2. *The Poincaré dual of $[m] \in H_1(M)$ (denoted by $P.D.[m]$) can not be represented by union of embedded tori or spheres. Therefore, the manifold M satisfies the conditions of Corollary 2.7.*

Proof. Since M is irreducible, embedded spheres are all null-homologous. Thus we just consider the case $P.D.[m] = \sum [T_i]$ where T_i are embedded tori. We can assume each T_i intersect both m and $T \times \{0\}$ transversely. Since the intersection number of $P.D.[m]$ with m equals 1, some T_i must intersect m for odd times. Without loss of generality, we can assume this torus to be T_1 . Now suppose $T_1 \cap (T \times \{0\}) = \bigcup_{j=1}^n \gamma_j$ where γ_j are disjoint simple closed curves. We can assume none of these curves bound disks in T_1 . Otherwise, by choosing the innermost curve which bounds a disk, we are able to find some γ_i bounding a disk $D \subset T_1$ and the interior \mathring{D} does not intersect $T \times \{0\}$. Since $T \times \{0\}$ is incompressible, γ_i also bounds a disk $D' \subset T \times \{0\}$. Notice that $D \cup D'$ is null-homologous since M is irreducible. We can replace T_1 with $(T_1 \setminus D) \cup D'$ to eliminate the intersection γ_i .

Moreover, suppose some γ_i bounds a disk $D'' \in T \times \{0\}$. As before, we can assume that \mathring{D}'' does not intersect T_1 by choosing the innermost curve. Consider the surface $T'_1 = ((M \setminus (T \times [-1, 1])) \cap T_1) \cup (D'' \times \{\pm 1\})$. Notice that γ_i is a non-separating curve in T_1 since it does not bound a disk. We see that T'_1 is an embedded sphere representing the same (non-zero) homology class as T_1 , which is a contradiction with the fact that M is irreducible.

Now each γ_i is not null-homologous in either T_1 or $T \times \{0\}$. Since these curves don't intersect each other, they are parallel in both T_1 and $T \times \{0\}$. Notice that the components of $T_1 \setminus (\bigcup_{j=1}^n \gamma_j)$ are contained in the two components of $M \setminus (T \times \{0\})$ alternatively, which implies that n is even. Since each γ_i intersects m for the same times, the intersection number of T_1 with m is even. This is a contradiction. \square

Now we consider the rank 2 unitary E with $c_1(E)$ the Poincaré dual of $[m] \in H_1(M)$. We can get \mathfrak{g}_E by the following procedures:

- Construct trivialized $SO(3)$ bundles over $Y_j - N(K_j)$ ($j = 1, 2$) and $T \times [-1, 1]$. Denote them by \mathfrak{g}_1 , \mathfrak{g}_2 and \mathfrak{g}_0 respectively.

- Choose a map $f : T^2 \rightarrow SO(3)$ such that $f_* : \pi_1(\partial N_0) \rightarrow \pi_1(SO(3)) \cong \mathbb{Z}_2$ satisfies:

$$(8) \quad f_*([m]) = [0], f_*([l]) = [1].$$

- Glue $\mathfrak{g}_1, \mathfrak{g}_0$ along $T \times \{-1\}$ using the identity map and glue $\mathfrak{g}_0, \mathfrak{g}_2$ along $T \times \{+1\}$ using f .

In order to do holonomy perturbations, we embed $H_j \times S^1$ ($j = 1, 2$) into M by i_1, i_2 as follows:

$i_1(H_1 \times S^1) = T \times [-1, 0]$ and $i_1(* \times S^1)$ is parallel to $l \subset T^2 \times [-1, 0]$, which means that we use the holonomy along the longitude to do perturbation Φ_1 .

$i_2(H_2 \times S^1) = T \times [0, 1]$ and $i_2(* \times S^1)$ is parallel to $m \subset T^2 \times [0, 1]$, which means that we use the holonomy along the meridian to do perturbation Φ_2 .

Now we consider the perturbed Chern-Simons functional, $CS + \Phi_1 + \Phi_2 : \mathcal{A} \rightarrow \mathbb{R}$.

Notice that we already have trivializations of \mathfrak{g}_E when restricted to $Y_j - N(K_j)$ and $T^2 \times [-1, 1]$. Thus for a connection $A \in \mathcal{A}$, we can restrict A to these pieces and use the trivializations to lift it to trivialized $SU(2)$ bundles. We have six loops: $m \times \{j\} \subset T^2 \times [-1, 1]$ and $l \times \{j\} \subset T^2 \times [-1, 1]$, $j = -1, 0, 1$. Denote them by m_j, l_j .

If A is a critical point of $CS + \Phi_1 + \Phi_2$, denote by \tilde{A} the $SU(2)$ lift of $A|_{\mathfrak{g}_0}$.

Since we only do perturbations in the interior of $T \times [0, -1]$ and $T \times [0, 1]$, we have $\tilde{A}|_{T \times \{j\}}$ flat for $j = -1, 0, 1$. Taking the holonomy of $\tilde{A}|_{T \times \{j\}}$, we get a representation $\rho_j : \pi_1(T^2) \rightarrow SU(2)$.

Notation 3.3. For $(\theta, \eta) \in (\mathbb{R}/2\pi\mathbb{Z})^2$, we denote by $\rho_{(\theta, \eta)}$ the representation $\pi_1(T^2) \rightarrow SU(2)$ which maps m to $\begin{pmatrix} e^{i\theta} & 0 \\ 0 & e^{-i\theta} \end{pmatrix}$ and l to $\begin{pmatrix} e^{i\eta} & 0 \\ 0 & e^{-i\eta} \end{pmatrix}$. We write $\rho \sim \rho'$ if two representations ρ and ρ' differ by a conjugation. It is easy to see that $\rho_{(\theta, \eta)} \sim \rho_{(\theta', \eta')}$ if and only if $(\theta, \eta) = \pm(\theta', \eta')$.

Applying Lemma 2.2 to Φ_1 with $\alpha_1 = m_{-1}, \alpha_2 = m_0, \beta_1 = l_{-1}, \beta_2 = l_0$, we see that $\rho_{-1} \sim \rho_{(\theta_{-1}, \eta_{-1})}$ and $\rho_0 \sim \rho_{(\theta_0, \eta_0)}$ with:

$$(9) \quad \eta_{-1} = \eta_0 \text{ and } \theta_{-1} - \theta_0 = f'_1(\eta_0).$$

Applying Lemma 2.2 to Φ_2 with $\alpha_1 = l_0, \alpha_2 = l_1, \beta_1 = m_0, \beta_2 = m_1$, we see that $\rho_0 \sim \rho_{(\hat{\theta}_0, \hat{\eta}_0)}$ and $\rho_1 \sim \rho_{(\theta_1, \eta_1)}$ with:

$$(10) \quad \hat{\theta}_0 = \theta_1 \quad \text{and} \quad \hat{\eta}_0 - \eta_1 = f'_2(\theta_1).$$

Remark 3.4. Recall that we choose $\phi_1, \phi_2 : SU(2) \rightarrow \mathbb{R}$ to define Φ_1 and Φ_2 . They give us f_1, f_2 .

We can also use the trivialization of $\mathfrak{g}_j (j = 1, 2)$ to lift A to flat $SU(2)$ connections over $Y_j - N(K_j)$. Denote them by \tilde{A}_1 and \tilde{A}_2 . Because we glue $\mathfrak{g}_1, \mathfrak{g}_0$ using identity map, we have:

$$\text{Hol}_{m_{-1}}(\tilde{A}) = \text{Hol}_{m_{-1}}(\tilde{A}_1), \quad \text{Hol}_{l_{-1}}(\tilde{A}) = \text{Hol}_{l_{-1}}(\tilde{A}_1).$$

Since we glue \mathfrak{g}_0 and \mathfrak{g}_2 using f , their trivializations do not agree on their common boundary. Different trivializations of the $SO(3)$ bundle give different $SU(2)$ lifts of $A|_{T \times \{2\}}$. By Formula (8), it is easy to check:

$$\text{Hol}_{m_1}(\tilde{A}) = \text{Hol}_{m_1}(\tilde{A}_2), \quad \text{Hol}_{l_1}(\tilde{A}) = -\text{Hol}_{l_1}(\tilde{A}_2)$$

Notation 3.5. Let $S \subset (\mathbb{R}/2\mathbb{Z}\pi)^2$ be a subset. If h is a function with period 2π , we denote the set $\{(\theta, \eta + h(\theta)) \mid (\theta, \eta) \in S\}$ by $S + (*, h)$ and the set $\{(\theta + h(\eta), \eta) \mid (\theta, \eta) \in S\}$ by $S + (h, *)$. We also denote the set $\{(\theta + a, \eta + b) \mid (\theta, \eta) \in S\}$ by $S + (a, b)$ for constant a, b .

By Definition 2.10 and the discussion above, we see that

$$(11) \quad (\theta_{-1}, \eta_{-1}) \in R_K, (\theta_1, \eta_1) \in R_K + (0, \pi).$$

By Formula (9), we have $(\theta_0, \eta_0) \in R_K + (-f'_1, *)$. Since $\rho_{(\theta_0, \eta_0)} \sim \rho_0 \sim \rho_{(\hat{\theta}_0, \hat{\eta}_0)}$, we have $(\hat{\theta}_0, \hat{\eta}_0) = \pm(\theta_0, \eta_0)$. Notice that $R_K + (-f'_1, *)$ is symmetric under the reflection at $(0, 0)$. We have $(\hat{\theta}_0, \hat{\eta}_0) \in R_K + (-f'_1, *)$. By Formula (10), we see that $(\theta_1, \eta_1) \in (R_K + (-f'_1, *)) + (*, -f'_2)$. Thus (θ_1, η_1) is an intersection point of $(R_K + (-f'_1, *)) + (*, -f'_2)$ and $R_K + (0, \pi)$. We get the following important proposition.

Proposition 3.6. *Suppose that $(R_K + (-f'_1, *)) + (*, -f'_2)$ does not intersect $R_K + (0, \pi)$. Then the perturbed Chern-Simons functional $CS + \Phi_1 + \Phi_2$ has no critical point.*

Now we are ready to prove Theorem 1.5.




Figure 1.

Proof. Suppose K has trivial A-polynomial. Then $R_K \setminus \{\eta = 2k\pi\}$ has only finitely many points, by Lemma 1.1. If there exists no representation $\pi_1(Y) \rightarrow SU(2)$ mapping K to -1 , then $(0, \pi) \notin R_K$. By Lemma 2.14 $(k\pi, \pm\pi) \notin R_K$. So, we can choose an odd, 2π -periodic function g_1 such that the curve $\{g_1(\eta) = \theta, 0 < \eta < 2\pi\}$ does not intersect R_K . Then $R_K + (g_1, *)$ does not intersect $\{\theta = k\pi, 0 < \eta < 2\pi\}$. This can be done because there are only finitely many points to move away from the lines $\{\theta = k\pi, 0 < \eta < 2\pi\}$ and none of them have η component equal to π . We find another odd, 2π -periodic function g_2 such that $(R_K + (g_1, *)) + (*, g_2)$ does not intersect the translation $R_K + (0, \pi)$. This can be done because $R_K + (g_1, *)$ has only finitely many points outside the circle $\theta = 2k\pi$ and none of them have θ component equal to $k\pi$. $R_K + (0, \pi)$ has only finitely many points off the circle $\theta = \pi$. We can move these discrete points away from the one dimensional components. As long as $|g_2| < \frac{\pi}{2}$, the one dimensional component of $(R_K + (g_1, *)) + (*, g_2)$ and that of $R_K + (0, \pi)$ do not intersect.

Then we find even, 2π -periodic functions f_1, f_2 such that $f'_j = -g_j$ and use them to define perturbations Φ_j . By Proposition 3.6, the perturbed Chern-Simons functional $CS + \Phi_1 + \Phi_2$ has no critical point. However, since K is not contained in a 3-ball by Theorem 1.2, $CS + \Phi_1 + \Phi_2$ has at least one critical point by Lemma 3.2 and Corollary 2.7. This is a contradiction. \square

Theorem 1.6 is a trivial corollary of Theorem 1.5. Here are some other corollaries.

Remark 3.7. If K is null-homotopic, then it is easy to see that R_K has no points on the line $\theta = 0, 0 < \eta < 2\pi$. Therefore, Theorem 1.6 can be proved using the perturbation Φ_2 without the use of Φ_1 .

Corollary 3.8. *If K is a non-trivial, null-homologous knot in an irreducible, orientable 3-manifold M and $\pi_1(M)/(K)$ has no $SO(3)$ representation with non-cyclic image, then $A_K \neq l - 1$. Here (K) denotes the normal subgroup group generated by $[K] \in \pi_1(M)$. In particular, if $\pi_1(M)$ has no non-cyclic $SO(3)$ representation, then any null-homologous, non-trivial knot has non-trivial A-polynomial.*

Proof. Suppose K has trivial A-polynomial. Then there exists $\rho : \pi_1(M) \rightarrow SU(2)$ such that $\rho(K) = -1$. The representation ρ induces a $SO(3)$ -representation of $\pi_1(M)$ which factors through $\pi_1(M)/(K)$. By our assumption, this representation has cyclic image and so does ρ . Thus ρ factors through $H_1(M)$, which is impossible because K is null homologous and $\rho(K) \neq 1$. \square

Example 3.9. *If $K \subset M$ is a non-trivial knot in the Poincaré homology sphere and $2[K] \neq e \in \pi_1(M)$, then $A_K \neq l - 1$.*

Example 3.10. *Let $K \subset Y_1$ be a non-trivial knot in Y_1 , where Y_1 is the manifold obtained by doing 1-surgery along some knot K' in S^3 . If K is homotopic in Y_1 to the meridian of $S^3 - K'$ then $A_K \neq l - 1$.*

The following example is given by Dunfield:

Example 3.11. *Let Y be the $\frac{37}{2}$ -surgery manifold on $(-2, 3, 7)$ -pretzel knot. Then $\pi_1(Y)$ admits no non-cyclic $SO(3)$ representation. Therefore any non-trivial null-homologous knot $K \subset Y$ has non-trivial A-polynomial.*

3.2. The S^3 case

Now let's go back to S^3 and use the technique in [11] to prove Theorem 1.8. Let K be a non-trivial knot in S^3 . By Remark 1.3, we can assume that $A_K(m, l) = (l - 1)f(m, l)$. By Theorem 1.2, we have $f(m, l) \neq 1$. Denote the algebraic curve defined by $f(m, l)$ in $\mathbb{C} \oplus \mathbb{C}$ by C .

Consider its intersection with the unit torus $T^2 = \{(m, l) \mid |m| = |l| = 1\}$. It is a closed subset of the torus. The following fact is straightforward by Lemma 1.1 and Formula (7).

Lemma 3.12. *If $(\theta, \eta) \in R_K$ and $\eta \neq 2k\pi$, then either $(e^{i\theta}, e^{i\eta})$ is a 0-dimensional component of $h^{-1}(\overline{\text{Im}(r)})$ or $(e^{i\theta}, e^{i\eta}) \in C \cap T^2$.*

Now we can prove Theorem 1.8:

Proof. Suppose for some unit length $l_0 = e^{i\eta_0}$ ($\eta_0 \in [0, 2\pi)$), we can't find $m_0 = e^{i\theta_0}$ such that $f(m_0, l_0) = 0$. Since $C \cap T^2$ is a compact set, there exists $\epsilon > 0$ such that this holds for any $\eta'_0 \in [\eta_0 - \epsilon, \eta_0 + \epsilon]$. Thus we can assume that $\eta_0 \neq 0$ and the line $\{l = e^{i\eta_0}\} \subset \mathbb{C} \oplus \mathbb{C}$ does not pass through the 0-dimensional components of $h^{-1}(\overline{\text{Im}(r)})$. By Lemma 3.12, we see that R_K does not intersect the circle $\eta = \eta_0$. Thus the translation $R_K + (0, -\pi)$ (denote it by S_K) does not intersect the circle $\eta = \eta_0 - \pi$. By Definition 2.10, S_K is symmetric under the reflection at the origin. Thus S_K does not intersect the circle $\eta = \pi - \eta_0$. We assume that $\eta_0 \geq \pi$ for convenience and the other cases can be proved similarly. Now consider the following line segments:

- $c_1 = \{(\theta, \eta_0 - \pi) \mid \theta \in [0, \pi]\}$;
- $c_2 = \{(\theta, \pi - \eta_0) \mid \theta \in [-\pi, 0]\}$;
- $c_3 = \{(0, \eta) \mid \eta \in [\pi - \eta_0, \eta_0 - \pi]\}$;
- $c_4 = \{(\pi, \eta) \mid \eta \in [0, \eta_0 - \pi]\}$
- $c_5 = \{(-\pi, \eta) \mid \eta \in [\pi - \eta_0, 0]\}$




Figure 2.

By the discussion before, S_K does not intersect c_1 and c_2 . Since S^3 is simply connected, S_K does not intersect c_3, c_4, c_5 . We can join all these

segments to get a piecewise linear path from $(-\pi, 0)$ to $(\pi, 0)$ and passing through $(0, 0)$. By Remark 2.12, we can find a small neighborhood N of this path such that $S_K \cap N = \emptyset$. It is easy to see that we can find an odd, 2π -periodic function $g : [-\pi, \pi] \rightarrow (-\pi, \pi)$ such that the graph of g lies in N . Thus S_K does not intersect the graph of g . We can find an even function $f : [-\pi, \pi] \rightarrow \mathbb{R}$ with period 2π satisfying $f' = g$.

We consider the rank 2 unitary bundle E over Y_0 (recall that Y_0 is the 0-surgery manifold along K) with $c_1(E)$ the Poincaré dual of the meridian. We do a holonomy perturbation along the surgery solid torus in Y_0 using the even function f . By Lemma 2.1, the critical points of the perturbed Chern-Simons functional correspond to the intersection of S_K with the graph of g , which is empty. (We have S_K instead of R_K here, because E has non-trivial first Chern class.) So the perturbed Chern-Simons functional has no critical points. This contradicts Corollary 2.7. \square

This theorem has the following corollary, which was also proved by Boden in [1]:

Corollary 3.13. *For any non-trivial knot $K \subset S^3$, $\deg_M A(M, L) \neq 0$.*

Acknowledgements

The author wishes to thank Nathan Dunfield, Hans Boden and Yi Ni for valuable discussions and comments. The author is especially grateful to Ciprian Manolescu for inspiring conversations and helpful suggestions in writing this paper.

References

- [1] H. U. Boden, *Non-triviality of the m -degree of the A-polynomial*. Proceedings of the American Mathematical Society, (2014) 2173–2177.
- [2] S. Boyer and X. Zhang, *Every nontrivial knot in S^3 has nontrivial A-polynomial*. Proceedings of the American Mathematical Society, (2005) 2813–2815.
- [3] P. J. Braam and S. K. Donaldson, *Floers work on instanton homology, knots and surgery*. In: The Floer memorial volume, 195–256, Springer (1995).

- [4] D. Cooper, M. Culler, H. Gillet, D. D. Long and P. B. Shalen, *Plane curves associated to character varieties of 3-manifolds*. *Inventiones mathematicae*, **118** (1994), no. 1, 47–84.
- [5] N. M. Dunfield and S. Garoufalidis, *Non-triviality of the A -polynomial for knots in S^3* . *Algebraic & Geometric Topology*, **4** (2004), no. 2, 1145–1153.
- [6] Y. Eliashberg, *A few remarks about symplectic filling*. *Geometry & Topology*, **8** (2004), no. 1, 277–293.
- [7] Y. Eliashberg, W. C. Thurston and O. P.-I. Macdonald, *University Lecture Series*. Amer. Math. Soc (1998).
- [8] P. Feehan and T. G. Leness, *A general $SO(3)$ -monopole cobordism formula relating Donaldson and Seiberg-Witten invariants*. Preprint, [arXiv:math/0203047](https://arxiv.org/abs/math/0203047), (2002).
- [9] D. Gabai, *Foliations and the topology of 3-manifolds*. *Bulletin of the American Mathematical Society*, **8** (1983), no. 1, 77–80.
- [10] P. Kronheimer, T. Mrowka, P. Ozsváth and Z. Szabó, *Monopoles and lens space surgeries*. *Annals of mathematics*, (2007) 457–546.
- [11] P. B. Kronheimer and T. S. Mrowka, *Dehn surgery, the fundamental group and $SU(2)$* . Preprint, [arXiv:math/0312322](https://arxiv.org/abs/math/0312322), (2003).
- [12] P. B. Kronheimer and T. S. Mrowka, *Wittens conjecture and Property P*. *Geometry & Topology*, **8** (2004), no. 1, 295–310.
- [13] C. H. Taubes, *The Seiberg-Witten invariants and symplectic forms*. *Math. Res. Lett.*, **1** (1994), no. 6, 809–822.

DEPARTMENT OF MATHEMATICS
UNIVERSITY OF CALIFORNIA LOS ANGELES
BOX 951555, LOS ANGELES, CA 90095-1555, USA
E-mail address: juliuslin@math.ucla.edu

RECEIVED JULY 5, 2013