

On some nonadmissible smooth irreducible representations for GL_2

DANIEL LE

Let $p > 2$ be a prime. We give examples of smooth absolutely irreducible representations of $\mathrm{GL}_2(\mathbb{Q}_{p^3})$ over \mathbb{F}_{p^3} which are not admissible.

1. Introduction

Smooth representations of p -adic reductive groups arise naturally in the theory of automorphic forms. Smooth here means that every vector is invariant under an open subgroup. Classical finite-dimensionality results for automorphic forms imply *admissibility*: the invariants of the representation under any open subgroup is finite-dimensional. Both of these notions make sense for a base field of any characteristic. Representation theory over base fields of positive characteristic has attracted considerable attention in recent years because of its connection to congruences of automorphic forms and the modularity of Galois representations.

In the recent groundbreaking work [2], smooth, irreducible, admissible mod p representations of connected reductive p -adic groups are classified in terms of supercuspidal representations, closely mirroring the earlier theory in characteristic not equal to p . For a base field of characteristic different from p , it is known from [9, II.2.8] moreover that every smooth irreducible representation of a connected reductive p -adic group is admissible. [1, Question 1] asks whether a similar statement is true for mod p representations. As mentioned in *loc. cit.*, this question has an affirmative answer in some simple cases and when the group is $\mathrm{GL}_2(\mathbb{Q}_p)$ combining results of [3–5]. We provide a negative answer, at least when $p > 2$, already for GL_2 but over a larger field.

Theorem 1.1. *Let $p > 2$. There exists a smooth absolutely irreducible $\mathrm{GL}_2(\mathbb{Q}_{p^3})$ -representation over \mathbb{F}_{p^3} which is not admissible.*

It will be clear from the construction that there are infinitely many such representations. Moreover, similar constructions exist for unramified extensions of larger degree (see Remark 2.3), but we content ourselves with describing the simplest example. The above result is yet another example of a distinguishing feature of the mod p theory, namely that the theory is very sensitive to the field of definition of the group.

Admissibility is a desirable property, in part because it implies that the irreducible representation has a central character, admits Hecke eigenvalues for weights, and has an endomorphism ring of finite dimension over the base field. [1, Question 2, Question 8] ask whether irreducible mod p representations must have central characters and Hecke eigenvalues. The representations that we construct have central characters and Hecke eigenvalues (matching certain supersingular representations), and so we do not answer these questions. However, by restricting scalars for a representation we construct, we also prove the following.

Theorem 1.2. *There exists a smooth irreducible $\mathrm{GL}_2(\mathbb{Q}_{p^3})$ -representation over \mathbb{F}_{p^3} whose endomorphisms contain $\overline{\mathbb{F}}_p$.*

Of course, such a representation cannot be absolutely irreducible as the endomorphism ring over $\overline{\mathbb{F}}_p$ would contain $\overline{\mathbb{F}}_p \otimes_{\mathbb{F}_{p^3}} \overline{\mathbb{F}}_p$.

We now make brief remarks on the construction. Irreducible mod p representations are typically rather difficult to construct, much less nonadmissible ones. Global constructions coming from the theory of automorphic forms always give admissible representations and parabolic induction preserves admissibility. However, the Bruhat–Tits tree and the diagrams of [8] give a powerful method of constructing mod p representations of p -adic GL_2 with fixed K -socle where K is the maximal compact subgroup. [7] uses this close control of the K -socle to prove both irreducibility and admissibility for many representations that they construct. The main idea of this paper is that the control of the K -socle can also be used to prove irreducibility and *nonadmissibility*. We construct an infinite-dimensional diagram that gives rise to a nonadmissible $\mathrm{GL}_2(\mathbb{Q}_{p^3})$ -representation, and prove irreducibility using the methods of *ibid*.

Acknowledgments

We thank Marie-France Vigneras for encouraging the author to work on this problem. The intellectual debt owed to the work of Christophe Breuil and Vytautas Paškūnas will be clear to the reader. We thank Stefano Morra for

producing some diagrams in an earlier version. We thank Benjamin Schraen for suggesting that we could prove Theorem 1.2. We thank Florian Herzig, Yongquan Hu, Karol Koziol, and an anonymous referee for many helpful comments on an earlier version. The author was supported by the Simons Foundation under an AMS-Simons travel grant and by the National Science Foundation under the Mathematical Sciences Postdoctoral Research Fellowship No. 1703182. We thank the University College London, ENS de Lyon, and L’Institut Montpelliérain Alexander Grothendieck for providing hospitality and excellent working conditions while part of this work was carried out.

1.1. Notation

Let $p > 2$ and let q be p^f for a positive integer f . Fix an algebraic closure $\overline{\mathbb{F}}_p$ of \mathbb{F}_q . If V is an \mathbb{F}_q -vector space, let $V_{\overline{\mathbb{F}}_p}$ denote $V \otimes_{\mathbb{F}_q} \overline{\mathbb{F}}_p$.

Let G be $\mathrm{GL}_2(\mathbb{Q}_q)$, Z the center of G , K be $\mathrm{GL}_2(\mathbb{Z}_q)$, and I (resp. I_1) the preimage in K of the upper triangular matrices (resp. unipotent upper triangular matrices) in $\mathrm{GL}_2(\mathbb{F}_q)$ under the natural reduction map. Let $\Pi \in G$ be the matrix $\begin{pmatrix} 0 & 1 \\ p & 0 \end{pmatrix}$. Then Π normalizes I and the normalizer $N(I)$ of I is $IZ \sqcup IZ\Pi$. Moreover, we have an isomorphism

$$(1.1) \quad N(I) / \left\langle \begin{pmatrix} p & 0 \\ 0 & p \end{pmatrix} \right\rangle \xrightarrow{\sim} I \rtimes \mathbb{Z}/2$$

$$(1.2) \quad \Pi \mapsto (\mathrm{id}, 1).$$

For a character χ of IZ , let χ^s be the character of IZ given by precomposing χ by Π -conjugation. If V is an IZ -representation and χ a character of IZ , we let V^χ be the χ -isotypic part of V .

2. Diagrams

2.1. Diamond diagrams

A *diagram* is a triple (D_0, D_1, r) where D_0 is a smooth KZ -representation, D_1 is a smooth $N(I)$ -representation, and r is an IZ -equivariant map $D_1 \rightarrow D_0$. A diagram is a *basic 0-diagram* if r induces an isomorphism $D_1 \xrightarrow{\sim} D_0^{I_1}$.

Let $\rho : G_{\mathbb{Q}_q} \rightarrow \mathrm{GL}_2(\overline{\mathbb{F}}_p)$ be a generic continuous irreducible representation in the sense of [7, Definition 11.7] (such representations exist with the assumption that $p > 2$). Let $\mathcal{D}(\rho)$ be the set of Serre weights defined in [7, §11]. To ρ , [7, Theorem 13.8] attaches a family of basic 0-diagrams. We fix

for the rest of the paper a basic 0-diagram $(D_0(\rho), D_1(\rho), r)$ in this family which is defined over \mathbb{F}_q . That is $D_0(\rho)$ and $D_1(\rho)$ are finite dimensional KZ and $N(I)$ -representations over \mathbb{F}_q , respectively, and $(D_0(\rho)_{\overline{\mathbb{F}_p}}, D_1(\rho)_{\overline{\mathbb{F}_p}}, r)$ is a member of the family constructed in *loc. cit.* Then r identifies $D_1(\rho)$ with $D_0(\rho)^{I_1}$ as IZ -representations, which we will identify implicitly.

In fact, the isomorphism classes of $D_0(\rho)$ and $D_1(\rho)$ do not depend on the above choice (though r does). The K -representation $D_0(\rho)|_K$ satisfies the following properties:

- the K -action on $D_0(\rho)$ factors through $\mathrm{GL}_2(\mathbb{F}_q)$;
- there is a direct sum decomposition

$$D_0(\rho) = \oplus_{\sigma \in \mathcal{D}(\rho)} D_{0,\sigma}(\rho)$$

where the $\mathrm{GL}_2(\mathbb{F}_q)$ -socle of $D_{0,\sigma}(\rho)$ is σ for all $\sigma \in \mathcal{D}(\rho)$;

- the Jordan–Hölder factors of $D_0(\rho)$ are multiplicity free ([7, Theorem 13.8]), and $D_1(\rho)$ is a multiplicity free semisimple IZ -representation ([7, Lemma 14.1]).

Recall from [7, Lemma 11.4] and the paragraph thereafter that there is a bijection

$$\begin{aligned} 2^{\mathbb{Z}/f} &\rightarrow \mathcal{D}(\rho) \\ J &\mapsto \sigma_J \end{aligned}$$

Define an automorphism $\delta : 2^{\mathbb{Z}/f} \rightarrow 2^{\mathbb{Z}/f}$ by $j \in \delta(J)$ if and only if $j + 1 \in J$ (resp. $j + 1 \notin J$) for $j \neq 0$ (resp. for $j = 0$). This “shift then flip at $j = 0$ ” is denoted δ_i in [7, §15].

We introduce one final piece of notation. For $0 \leq s \leq q - 1$, let

$$S_s := \sum_{\lambda \in \mathbb{F}_q} \lambda^s \begin{pmatrix} [\lambda] & 1 \\ 1 & 0 \end{pmatrix} \in \mathbb{F}_q[K].$$

Proposition 2.1. *Let v be a nonzero element in $D_1(\rho)^\chi$. Then there is a unique $0 \leq s(\chi) \leq q - 1$ such that $S_{s(\chi)}(v)$ is a nonzero element of $(\mathrm{soc}_K D_0(\rho))^{I_1}$.*

Proof. Since $D_1(\rho)^\chi$ is one-dimensional, the K -representation generated by v has irreducible socle using the last two bulleted points above (cf. the proof of [6, Proposition 5.1(i)]). The result now follows from [7, Lemma 2.7]. \square

Define a linear map

$$S : D_1(\rho) \rightarrow (\text{soc}_K D_0(\rho))^{I_1}$$

which maps a nonzero (IZ, χ) -eigenvector v to $S_{s(\chi)}v$.

We recall the following result.

Proposition 2.2. *Let χ_J be the I -character of $\sigma_J^{I_1}$. Then $S \circ \Pi$ gives an isomorphism $D_1(\rho)^{\chi_J}$ to $D_1(\rho)^{\chi_{\delta(J)}}$ for all $J \in 2^{\mathbb{Z}/f}$.*

Proof. This follows from [7, Lemma 15.2] (see also the proof of [6, Proposition 5.1]). \square

2.2. An infinite diagram

In this section, we let f be 3.

Remark 2.3. When $f = 2$, $2^{\mathbb{Z}/f}$ is a single δ -orbit. When $f = 3$, $2^{\mathbb{Z}/f}$ consists of two δ -orbits, namely

$$\begin{aligned} \emptyset &\mapsto \{0\} \mapsto \{0, 2\} \mapsto \{0, 1, 2\} \mapsto \{1, 2\} \mapsto \{1\} \mapsto \emptyset \\ \{2\} &\mapsto \{0, 1\} \mapsto \{2\}. \end{aligned}$$

For $f > 3$, $2^{\mathbb{Z}/f}$ always contains more than one δ -orbit since the size of an orbit must divide the order of the automorphism δ , which is $2f$. It is the existence of more than one δ -orbit which allows us to make the construction in this section.

Let D_0 be the KZ -representation $\oplus_{i \in \mathbb{Z}} D_{0,i}$ where there is a fixed isomorphism $D_{0,i} \cong D_0(\rho)_{\overline{\mathbb{F}}_p}$. Let ι_i be the inclusion $D_0(\rho) \subset D_{0,i} \subset D_0$. For $v \in D_0(\rho)$, we denote $\iota_i(v)$ by v_i .

Let D_1 be $D_0^{I_1}$. Let $\lambda = (\lambda_i)_{i \in \mathbb{Z}}$ be in $\prod_{i \in \mathbb{Z}} \overline{\mathbb{F}}_p^\times$. For such a λ , we now define an action of $N(I)$ on D_1 such that Π^2 acts trivially. By (1.1), it suffices to define an involution on D_1 taking D_1^χ to $D_1^{\chi^s}$ for every character χ of IZ . We will denote this involution by $\tilde{\Pi}$.

Let χ_+ be the IZ -character of the space $\sigma_{\{1\}}^{I_1}$ and χ_- be the IZ -character of the space $\sigma_{\{0,1\}}^{I_1}$ (as usual Π^2 acts trivially).

Proposition 2.4. *There is an IZ -character χ_1 (resp. χ_2) such that both of the spaces $D_{0,\sigma_{\{2\}}}(\rho)^{\chi_1}$ and $D_{0,\sigma_\emptyset}(\rho)^{\chi_1^s}$ (resp. $D_{0,\sigma_{\{0,1\}}}(\rho)^{\chi_2}$ and $D_{0,\sigma_{\{0\}}}(\rho)^{\chi_2^s}$) are nonzero.*

Proof. This follows from an explicit check using [7, Corollary 14.10 and Lemma 15.2]. In the notation of [7, §11], we have that $\sigma_{\{2\}}$ corresponds to

$$(\lambda_0(r_0), \lambda_1(r_1), \lambda_2(r_2)) = (r_0, p - 2 - r_1, r_2 + 1)$$

and $\sigma_{\{0,1\}}$ corresponds to

$$(\lambda_0(r_0), \lambda_1(r_1), \lambda_2(r_2)) = (p - 1 - r_0, r_1 + 1, p - 2 - r_2).$$

Then χ_1 corresponds to

$$(\mu_0(\lambda_0(r_0)), \mu_1(\lambda_1(r_1)), \mu_2(\lambda_2(r_2))) = (p - 2 - r_0, p - 1 - r_1, r_2 + 1)$$

and χ_2 corresponds to

$$(\mu_0(\lambda_0(r_0)), \mu_1(\lambda_1(r_1)), \mu_2(\lambda_2(r_2))) = (p - r_0, r_1 + 1, r_2).$$

□

In fact, the characters χ_1 and χ_2 are uniquely described by the properties in Proposition 2.4, but we will not use this. As we will see, the only property that we will need is that χ_1 (resp. χ_2) is a character in $D_{0,\sigma_{\{2\}}}(\rho)^{I_1}$ (resp. $D_{0,\sigma_{\{0,1\}}}(\rho)^{I_2}$), which is not in $(\sigma_{\{2\}})^{I_1}$ (resp. $(\sigma_{\{0,1\}})^{I_1}$). The exact choices and formulas of Proposition 2.4 will not be important, and we include them only for the sake of concreteness.

If $v \in D_1(\rho)^\chi$ with

$$\chi \notin \{\chi_+, \chi_+^s, \chi_-, \chi_-^s, \chi_1, \chi_1^s\},$$

we define

$$\tilde{\Pi}(v_i) = (\Pi v)_i.$$

If $v \in D_1(\rho)^{\chi_+}$, then we define

$$\tilde{\Pi}(v_i) = (\Pi v)_{i+1}.$$

If $v \in D_1(\rho)^{\chi_-}$, then we define

$$\tilde{\Pi}(v_i) = (\Pi v)_{i-1}.$$

If $v \in D_1(\rho)^{\chi_1}$, then we define

$$\tilde{\Pi}(v_i) = \lambda_i(\Pi v)_i.$$

This now uniquely defines an $\overline{\mathbb{F}}_p$ -linear involution $\tilde{\Pi}$ of D_1 , and it takes D_1^χ to $D_1^{\chi^s}$ for every character χ of IZ as desired.

Let $D(\lambda)$ be the basic 0-diagram (D_0, D_1, can) with the above actions, where can denotes the canonical inclusion $D_1 \subset D_0$. We define an $\overline{\mathbb{F}}_p$ -linear map $\tilde{S} : D_1 \rightarrow (\text{soc}_K D_0)^{I_1}$ by the formula $\tilde{S}\iota_i = \iota_i S$, where S is as defined in §2.1.

3. The construction

For the purposes of notation, we review the proof of the following result, which is a special case of [7, Theorem 9.8], although we work over \mathbb{F}_q rather than $\overline{\mathbb{F}}_p$.

Theorem 3.1. *There exists a smooth G -representation τ over \mathbb{F}_q such that*

- *there is an injection of diagrams $(D_0(\rho), D_1(\rho), r) \subset (\tau|_{KZ}, \tau|_{N(I)}, \text{id})$;*
- *τ is generated as a G -representation by the image of $D_0(\rho)$; and*
- *the induced injection $\text{soc}_K D_0(\rho) \hookrightarrow \text{soc}_K \tau$ is an isomorphism.*

Proof. Let Ω be the K -injective envelope of $D_0(\rho)|_K$. We give Ω a KZ -action by demanding that Π^2 acts trivially. There is an idempotent $e \in \text{End}_I(\Omega)$ such that $e(\Omega)|_I$ is an I -injective envelope of $D_1(\rho)$. There is a decomposition of $e(\Omega)|_I$ as a direct sum

$$\bigoplus_{\chi} \Omega_{\chi},$$

where χ runs over the I -characters in $D_1(\rho)$ and Ω_{χ} is an I -injective envelope of the χ -isotypic part of $D_1(\rho)$. By [7, Lemma 9.5], there is an \mathbb{F}_q -linear map $e(\Omega) \rightarrow e(\Omega)$ which intertwines the action and Π -conjugate action of IZ , extends the action of Π on $D_1(\rho)$, and whose restriction to Ω_{χ} for each χ above gives a map

$$\Omega_{\chi} \rightarrow \Omega_{\chi^s}.$$

This gives an action of $N(I)$ on $e(\Omega)$. There is also an action of $N(I)$ on $(1 - e)(\Omega)$ by [7, Lemma 9.6]. This gives an action of $N(I)$ on Ω whose restriction to I is compatible with the action coming from KZ on Ω . By [8, Corollary 5.18], this gives an action of G on Ω . We then take τ to be the G -representation generated by $D_0(\rho)$. \square

Theorem 3.2. *There exists a smooth G -representation π over $\overline{\mathbb{F}}_p$ such that*

- *there is an injection of diagrams $D(\lambda) \subset (\pi|_{KZ}, \pi|_{N(I)}, \text{id})$;*
- *π is generated as a G -representation by the image of D_0 ;*

- the induced injection $\text{soc}_K D_0 \hookrightarrow \text{soc}_K \pi$ is an isomorphism; and
- if $\lambda \in \prod_{i \in \mathbb{Z}} \mathbb{F}_q^\times$, then π is defined over \mathbb{F}_q .

Proof. Let Ω be the K -injective envelope of $D_0(\rho)|_K$ as in the proof of Theorem 3.1. We give Ω a KZ -action by demanding that Π^2 acts trivially. Recall the definitions of $e \in \text{End}_I(\Omega)$ and Ω_χ from the proof of Theorem 3.1. Now let Ω_∞ be the KZ -representation $\bigoplus_{i \in \mathbb{Z}} \Omega_i$ where there is a fixed isomorphism $\Omega_i \cong \Omega_{\mathbb{F}_p}$. Let ι_i be the KZ -injection $\Omega \subset \Omega_i \subset \Omega_\infty$. To define an action of $N(I)$ on Ω_∞ , it suffices to define an involution, which we call $\tilde{\Pi}$, on Ω_∞ which intertwines the action and Π -conjugate action of IZ . For each $i \in \mathbb{Z}$, we define $\tilde{\Pi} \circ \iota_i|_{(1-e)(\Omega)}$ to be $\iota_i \circ \Pi|_{(1-e)(\Omega)}$. For $\chi \notin \{\chi_+, \chi_+^s, \chi_-, \chi_-^s, \chi_1, \chi_1^s\}$, we define $\tilde{\Pi} \circ \iota_i|_{\Omega_\chi}$ to be $\iota_i \circ \Pi|_{\Omega_\chi}$. We define $\tilde{\Pi} \circ \iota_i|_{\Omega_{\chi_+}}$ to be $\iota_{i+1} \circ \Pi|_{\Omega_{\chi_+}}$, $\tilde{\Pi} \circ \iota_i|_{\Omega_{\chi_-}}$ to be $\iota_{i-1} \circ \Pi|_{\Omega_{\chi_-}}$, and $\tilde{\Pi} \circ \iota_i|_{\Omega_{\chi_1}}$ to be $\iota_i \circ \lambda_i \Pi|_{\Omega_{\chi_1}}$. This completely determines the \mathbb{F}_p -linear involution $\tilde{\Pi}$. It is easy to see that the defined action of $N(I)$ on Ω_∞ extends the action of $N(I)$ on D_1 . By [8, Corollary 5.18], this gives an action of G on Ω_∞ . If $\lambda \in \prod_{i \in \mathbb{Z}} \mathbb{F}_q^\times$, then this action is defined over \mathbb{F}_q . Then if we let π be the G -subrepresentation of Ω_∞ generated by D_0 , π satisfies the required hypotheses. Indeed, we have that $\text{soc}_K \Omega_\infty = \text{soc}_K \pi = \text{soc}_K D_0$, and π is defined over \mathbb{F}_q if Ω_∞ is. \square

Let $D_{0,I}(\rho)$ and $D_{0,II}(\rho)$ be $D_{0,\sigma_{\{2\}}}(\rho) \oplus D_{0,\sigma_{\{0,1\}}}(\rho)$ and $\bigoplus_J D_{0,\sigma_J}(\rho)$, respectively, where the sum is over

$$J \in \{\emptyset, \{0\}, \{0, 2\}, \{0, 1, 2\}, \{1, 2\}, \{1\}\}.$$

(This partition $2^{\mathbb{Z}/3} = J \cup J^c$ corresponds to δ -orbits, see Remark 2.3.) We now recall the following special case of [7, Theorem 19.10(i)], since the arguments play a crucial role in the proof of Theorem 3.4.

Theorem 3.3. *Any G -representation τ satisfying the hypotheses in Theorem 3.1 is absolutely irreducible.*

Proof. Let $\tau' \subset \tau_{\mathbb{F}_p}$ be a nonzero G -subrepresentation. Since $\text{soc}_K \tau_{\mathbb{F}_p} \cong \text{soc}_K D_0(\rho)_{\mathbb{F}_p}$, there is a J such that $\text{Hom}_K(\sigma_J, \tau')$ is nonzero. Then by [7, Lemma 19.7], we have the inclusion $D_{0,\sigma_{\delta(J)}}(\rho)_{\mathbb{F}_p} \subset \tau'$. Repeating this, one obtains an inclusion of one of $D_{0,I}(\rho)_{\mathbb{F}_p}$ and $D_{0,II}(\rho)_{\mathbb{F}_p}$ in τ' . Then either $(\tau')^{I, \chi_1}$ or $(\tau')^{I, \chi_1^s}$ is nonzero. Applying Π , we see that they both must be nonzero so that $D_{0,\sigma_\emptyset}(\rho)_{\mathbb{F}_p}$ and $D_{0,\sigma_{\{2\}}}(\rho)_{\mathbb{F}_p}$ are both in τ' . Repeating the earlier argument, we have that $D_0(\rho)_{\mathbb{F}_p} \subset \tau'$. Since $\tau_{\mathbb{F}_p}$ is generated by $D_0(\rho)_{\mathbb{F}_p}$, we have that $\tau' = \tau_{\mathbb{F}_p}$. \square

The following is the main result of this section.

Theorem 3.4. *If $\lambda_0 \in \mathbb{F}_q$ and $\lambda_i \neq \lambda_0$ for all $i \neq 0$, then any G -representation π satisfying the hypotheses in Theorem 3.2 is irreducible over $\overline{\mathbb{F}}_p$. If moreover the \mathbb{F}_q -span of $(\lambda_i)_i$ is $\overline{\mathbb{F}}_p$, then π is irreducible as a G -representation over \mathbb{F}_q .*

Proof. Let π' be a nonzero G -subrepresentation of π seen as a representation over \mathbb{F}_q by restriction of scalars. Since $\text{soc}_K \pi' \subset D_0$, there exists $\sigma \in \mathcal{D}(\rho)$ such that $\text{Hom}_K(\sigma, \pi')$ is nonzero. Then there exists a $(c_i)_i$ in $\oplus_{i \in \mathbb{Z}} \overline{\mathbb{F}}_p$ such that

$$\left(\sum_i c_i \iota_i \right) (D_{0,\sigma}(\rho)) \cap \pi' \neq 0.$$

Lemma 3.5. *Suppose that $\sigma \in \mathcal{D}(\rho)$ and $(d_i)_i \in \oplus_{i \in \mathbb{Z}} \overline{\mathbb{F}}_p$ are elements such that*

$$\left(\sum_i d_i \iota_i \right) (D_{0,\sigma}(\rho)) \cap \pi' \neq 0.$$

Then for any $j \in \mathbb{Z}$,

$$\left(\sum_i d_i \iota_{i+j} \right) (D_0(\rho)) \subset \pi'.$$

Proof. We assume that σ is σ_\emptyset , as the other cases are similar. Then as in the proof of Theorem 3.3, we see from repeatedly applying $\widetilde{S}\widetilde{\Pi}$ that

$$\left(\sum_i d_i \iota_{i+j} \right) (D_{0,II}(\rho)) \subset \pi'$$

for $j > 0$. Since for each $j > 0$, we have that

$$\left(\sum_i d_i \iota_{i+j} \right) (D_{0,II}(\rho)^{\chi_2^s}) \subset \pi',$$

we have that

$$\left(\sum_i d_i \iota_{i+j} \right) (D_{0,I}(\rho)^{\chi_2}) \subset \pi'$$

for $j > 0$. Again repeatedly applying $\widetilde{S\Pi}$, we see that

$$\left(\sum_i d_i \iota_{i+j} \right) (D_{0,I}(\rho)) \subset \pi'$$

for all $j \in \mathbb{Z}$. Then since

$$\left(\sum_i d_i \iota_{i+j} \right) (D_{0,I}(\rho)^{\chi_2}) \subset \pi'$$

for all $j \in \mathbb{Z}$, we have that

$$\left(\sum_i d_i \iota_{i+j} \right) (D_{0,II}(\rho)^{\chi_2^s}) \subset \pi'$$

for all $j \in \mathbb{Z}$. We conclude that

$$\left(\sum_i d_i \iota_{i+j} \right) (D_{0,II}(\rho)) \subset \pi'$$

for all $j \in \mathbb{Z}$ by again repeatedly applying $\widetilde{S\Pi}$. \square

In the proof of the next lemma, we will use the following notation. For $(d_i)_i \in \oplus_{i \in \mathbb{Z}} \overline{\mathbb{F}}_p$, let $\#(d_i)_i$ be the cardinality of $\{i \in \mathbb{Z} | d_i \neq 0\}$.

Lemma 3.6. *There is a nonzero constant $c \in \overline{\mathbb{F}}_p$ such that $c\iota_0(D_{0,\sigma_{\{0\}}}(\rho)) \subset \pi'$.*

Proof. Fix nonzero elements $v^1 \in D_1(\rho)^{\chi_1}$ and $v^2 \in D_1(\rho)^{\chi_2}$. One checks that $(S\Pi)^2 v^1$ and $S\Pi v^2$ are nonzero elements in $\sigma_{\{0\}}^{I_1} \subset D_1(\rho)$ using the definition of χ_1 and χ_2 and Proposition 2.2. Thus, there exists a scalar $\mu \in \mathbb{F}_q^\times$ such that

$$(S\Pi)^2 v^1 = \mu S\Pi v^2.$$

Then by the definition of the action of Π on D_1 , we have that

$$(\widetilde{S\Pi})^2 v_i^1 = \lambda_i \mu \widetilde{S\Pi} v_i^2$$

for all $i \in \mathbb{Z}$.

By Lemma 3.5, there exists a nonzero $(c_i)_i$ in $\oplus_{i \in \mathbb{Z}} \overline{\mathbb{F}}_p$ such that

$$\left(\sum_i c_i \iota_i \right) D_0(\rho) \subset \pi'.$$

Assume that $\#(c_i)_i$ is minimal among such elements of $\oplus_{i \in \mathbb{Z}} \overline{\mathbb{F}}_p$. It suffices to show that $\#(c_i)_i = 1$ by Lemma 3.5. By Lemma 3.5, we can also assume that c_0 is nonzero.

Since $\sum_i c_i v_i^1$ and $\sum_i c_i v_i^2$ are in π' , then by the first paragraph, we have that

$$\sum_i c_i ((\tilde{S}\tilde{\Pi})^2 v_i^1 - \lambda_0 \mu \tilde{S}\tilde{\Pi} v_i^2) = \sum_i (\lambda_i - \lambda_0) c_i \mu \tilde{S}\tilde{\Pi} v_i^2$$

is in π' , using that $\lambda_0 \in \mathbb{F}_q$. We see from Lemma 3.5 that

$$\left(\sum_i c'_i \iota_i \right) D_{0, \sigma_{\{0\}}}(\rho) \cap \pi' \neq 0$$

for $c'_i = (\lambda_i - \lambda_0) c_i$. Since the $\lambda_i \neq \lambda_0$ for $i \neq 0$ and $c_0 \neq 0$, $\#(c'_i)_i = \#(c_i)_i - 1$. Since we assumed that $\#(c_i)_i$ is minimal, we must have that $\#(c_i)_i = 1$. \square

We now complete the proof of Theorem 3.4. By Lemma 3.5, it suffices to show that c in Lemma 3.6 can be taken to be any element of $\overline{\mathbb{F}}_p^\times$. If π' is a subrepresentation of π over $\overline{\mathbb{F}}_p$, this is clear. Now assume that the \mathbb{F}_q -span of $(\lambda_i)_i$ is $\overline{\mathbb{F}}_p$. By Lemma 3.5, $c\iota_j(D_0(\rho)^{X_1}) \subset \pi'$ for all $j \in \mathbb{Z}$. By applying $\tilde{\Pi}$ to $c\iota_j(D_0(\rho)^{X_1})$, we see that c can be taken to be $c\lambda_j$ for all $j \in \mathbb{Z}$. Since $(c\lambda_i)_i$ spans $\overline{\mathbb{F}}_p$ over \mathbb{F}_q , we are done. \square

Note that since D_0 is not admissible, any π as in Theorem 3.2 is not admissible. Taking $\lambda \in \prod_{i \in \mathbb{Z}} \mathbb{F}_q^\times$, Theorem 3.4 implies Theorem 1.1 by taking the \mathbb{F}_q -model of π constructed in Theorem 3.2. Since the endomorphisms of any such π must contain $\overline{\mathbb{F}}_p$, taking $(\lambda_i)_i$ to span $\overline{\mathbb{F}}_p$ over \mathbb{F}_q and restricting scalars of π to \mathbb{F}_q , Theorem 3.4 implies Theorem 1.2.

References

- [1] N. Abe, G. Henniart, F. Herzig, and M.-F. Vignéras, *Questions on mod p representations of reductive p -adic groups*.

- [2] N. Abe, G. Henniart, F. Herzig, and M.-F. Vignéras, *A classification of irreducible admissible mod p representations of p -adic reductive groups*, J. Amer. Math. Soc. **30** (2017), no. 2, 495–559.
- [3] L. Barthel and R. Livné, *Irreducible modular representations of GL_2 of a local field*, Duke Math. J. **75** (1994), no. 2, 261–292.
- [4] L. Berger, *Central characters for smooth irreducible modular representations of $\mathrm{GL}_2(\mathbf{Q}_p)$* , Rend. Semin. Mat. Univ. Padova **128** (2012), 1–6 (2013).
- [5] C. Breuil, *Sur quelques représentations modulaires et p -adiques de $\mathrm{GL}_2(\mathbf{Q}_p)$. I*, Compositio Math. **138** (2003), no. 2, 165–188.
- [6] C. Breuil, *Diagrammes de Diamond et (ϕ, Γ) -modules*, Israel J. Math. **182** (2011) 349–382.
- [7] C. Breuil and V. Paškūnas, *Towards a modulo p Langlands correspondence for GL_2* , Mem. Amer. Math. Soc. **216** (2012), no. 1016, vi+114.
- [8] V. Paškūnas, *Coefficient systems and supersingular representations of $\mathrm{GL}_2(F)$* , Mém. Soc. Math. Fr. (N.S.) (2004), no. 99, vi+84.
- [9] M.-F. Vignéras, *Représentations l -Modulaires d’un Groupe Réductif p -Adique avec $l \neq p$* , Vol. 137 of Progress in Mathematics, Birkhäuser Boston, Inc., Boston, MA (1996), ISBN 0-8176-3929-2.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF TORONTO
40 ST. GEORGE ST., TORONTO, ON CANADA M5S 2E4
E-mail address: le@math.toronto.edu

RECEIVED SEPTEMBER 29, 2018